



Development of EST-SSR Markers for *Primula ovalifolia* (Primulaceae) by Transcriptome Sequencing

Authors: Yuan, Shuai, Zeng, Gui, Shi, Miaomiao, and Zhang, Dianxiang

Source: Applications in Plant Sciences, 5(12)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1700100>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

DEVELOPMENT OF EST-SSR MARKERS FOR *PRIMULA OVALIFOLIA* (PRIMULACEAE) BY TRANSCRIPTOME SEQUENCING¹

SHUAI YUAN^{2,3}, GUI ZENG^{2,3}, MIAOMIAO SHI^{2,4}, AND DIANXIANG ZHANG²

²Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, People's Republic of China; and ³University of Chinese Academy of Sciences, Beijing 10049, People's Republic of China

- **Premise of the study:** Microsatellite primers were developed for *Primula ovalifolia*, a member of *Primula* section *Petiolares* (Primulaceae), to study the population genetics and species delimitation in this section.
- **Methods and Results:** A total of 4753 markers were successfully designated from 5139 putative simple sequence repeat loci. We isolated 38 expressed sequence tag-simple sequence repeat markers from 220 selected marker sites and tested polymorphism in three populations of *P. ovalifolia*, one of *P. tardiflora*, and one of *P. epilosa*. The number of alleles per locus ranged from one to 19, and the observed and expected levels of heterozygosity varied from 0 to 0.938 and 0 to 0.915, respectively. Most of the loci could be successfully cross-amplified in the two congeneric species.
- **Conclusions:** These markers will be useful for further population genetic analysis and gene flow estimation of *P. ovalifolia* and its relatives.

Key words: EST-SSR marker; interspecific transferability; *Primula ovalifolia*; Primulaceae; transcriptome.

Primula L. section *Petiolares* Pax (Primulaceae) is mainly distributed in the Hengduanshan-Himalaya Mountains with only a few members occurring in Kashmir, central China, and other regions (Hu and Kelso, 1996). *Primula ovalifolia* Franch. and *P. tardiflora* (C. M. Hu) C. M. Hu are two closely related species in the section. *Primula ovalifolia* is widely distributed in southwestern and adjacent central China, around the Sichuan Basin, mainly growing in shaded habitats in broad-leaved forests and ravines, with altitudes ranging from 600 to 2500 m. *Primula tardiflora* is morphologically similar to *P. ovalifolia* but with a more limited distribution, known only from a single locality in E'mei Mountain. It was first considered as a subspecies of *P. ovalifolia*, and later was regarded as distinct from *P. ovalifolia* by its higher altitude habitat, later floral phenology, and some neglected vegetative traits (Hu and Kelso, 1996). A phylogeographic study based on chloroplast DNA data suggested that *P. tardiflora* is genetically close to the E'mei Mountain population of *P. ovalifolia* (Xie et al., 2012). Development of highly polymorphic nuclear markers (e.g., simple sequence repeats [SSRs]) will help to delineate between the two species. In *Primula*, only a few genomic SSR markers have been developed for several species thus far, such as *P. vulgaris* Huds. (Van et al., 2006), *P. obconica* Hance (Yan

et al., 2010), *P. sieboldii* E. Morren (Ueno et al., 2011), *P. veris* L. (Bickler et al., 2013), *P. poissonii* Franch., and *P. wilsonii* Dunn (Zhang et al., 2013). Considering that these markers demonstrate low polymorphism and limited transferability, and that species of section *Petiolares* are phylogenetically distant from the above-named species, expressed sequence tag (EST)-SSRs specific to section *Petiolares* could provide useful tools for evolutionary and ecological studies in this section. In this study, we first obtained transcriptome data for *P. ovalifolia* using the Illumina platform and then designed marker pairs based on SSR loci. A subset of the markers was selected to investigate their polymorphism and transferability in congeneric species of *Primula*.

METHODS AND RESULTS

Transcriptome sequencing—Plants of *P. ovalifolia* were collected from Mount E'mei, Sichuan Province, China. Leaves from one individual were sampled, frozen immediately in liquid nitrogen, and stored at -80°C for RNA extraction and transcriptome sequencing. RNA was extracted using TRIzol Reagent (QIAGEN, Dusseldorf, Germany) and delivered to Genepioneer Technologies Corporation (Nanjing, China) for construction of cDNA libraries and sequencing. The cDNA libraries were sequenced on the Illumina HiSeq 2500 platform (Illumina, San Diego, California, USA) according to the manufacturer's recommendations. A total of 49,094,910 raw reads were obtained and deposited in the National Center for Biotechnology Information (NCBI) Short Read Archive under BioProject ID PRJNA379052 (accession no. SRP102475). Raw reads were first cleaned by trimming adapters and removing ambiguous reads ('N' > 10%) and low-quality reads (Phred score < 30). Clean reads were assembled into 142,468 transcripts using Trinity tools with default parameters (Haas et al., 2013) and were then clustered into 67,577 unigenes with TGICL version 2.1 (Pertea et al., 2003).

Development of EST-SSR markers—The EST-SSR loci were identified from unigenes longer than 1 kb using MicroSatellite identification tool

¹Manuscript received 23 August 2017; revision accepted 17 November 2017.

The authors are indebted to Mingzhi Li and Jing Deng for their kind help in the laboratory. This work was supported by the National Natural Science Foundation of China (grant no. 31400209, U1603231) and the Ministry of Science and Technology of China (grant no. 2013FY111200).

⁴Author for correspondence: mmshi@scbg.ac.cn

doi:10.3732/apps.1700100

Applications in Plant Sciences 2017 5(12): 1700100; <http://www.bioone.org/loi/apps> © 2017 Yuan et al. Published by the Botanical Society of America. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC-BY-NC-SA 4.0), which permits unrestricted noncommercial use and redistribution provided that the original author and source are credited and the new work is distributed under the same license as the original.

TABLE 1. Characteristics of 38 microsatellite loci identified in *Primula ovalifolia*.

Locus ^a	Primer sequences (5'–3')	Repeat motif	Allele size range (bp)	Fluorescent dye	GenBank accession no.
C11339	F: CGACTTCACTCCCACTGTTG R: GGTCAAAATCACCGGAAAGA	(CT) ₈	278–310	HEX	MF716482
C13959	F: TGGACGCCAATCTTTCTCAT R: TTGCATATCCCTCCCAATA	(AG) ₈	266–282	FAM	MF716483
C14309	F: ATTGCAAGTGTCTTTCGGC R: TCCCTTTGCTAAAAAGAAGGC	(TC) ₁₀	116–132	ROX	MF716484
C14710	F: TACACCGGTGGAAGAAGTC R: GGTGTTCTCTTAATCCCC	(AG) ₈	258–276	TAMRA	MF716485
C15228	F: TTTCCAATCCATGTCGTTC R: CAATTGACCCCAACAAACA	(AG) ₈	268–276	HEX	MF716486
C21614	F: GCGCCGTGACATAAATCATA R: GGTGGAGGTGTCGTAAGGA	(CT) ₉	180–184	HEX	MF716487
C23409	F: CAGTCAAATCACCAAGGGCT R: CAGCTTGTTCGATTGTTGGA	(CT) ₈	171–203	FAM	MF716488
C23644	F: GCGTAAGTAGTAGCGGTGGC R: CGCCCAATAACAAAACCCAG	(CT) ₈	238–268	TAMRA	MF716489
C23746	F: CCACTGCCTCCATTACCATT R: AACGTTCATTTTCAGGTGC	(AG) ₉	112–114	TAMRA	MF716490
C24233	F: CTGCAAAAACAGTCTTGGA R: GGGCAGTTTTGTGTCCATT	(CT) ₁₀	284–298	HEX	MF716491
C24268	F: ATGGCAAATTCGGATTCAAG R: ACACGCACGTCTCCTCTTCT	(AT) ₈	221–249	ROX	MF716492
C24676	F: CCTGCAAAACAGTTAGGCACA R: TTTTCGCTATTATCACCGCC	(GA) ₁₀	206–224	FAM	MF716493
C40984	F: AGGAGTGAGAGGGGTTGGT R: CACAACAATTAAGCAGACAAAAA	(CT) ₈	145–153	TAMRA	MF716494
C45417	F: GGGGAGCAGAGTAATAGG R: CTTGAAAGTGGCAAGGCAAT	(AT) ₁₀	166–252	ROX	MF716495
C46430	F: CCAAGCCACACCACACAT R: GAGGACGGAGAGTACGCAAG	(CT) ₈	285–305	ROX	MF716496
C47095	F: GTCTGATCATGGCAGTGGTG R: GATCGGACGGTGGAGAATAA	(GA) ₈	212	HEX	MF716497
C48258	F: GGTGAATCATCACCAATCC R: TGCCCAACATATGCCTTCT	(GA) ₉	176–212	TAMRA	MF716498
C48475	F: GGCCCAAAGGAAAGGATAA R: TGTGAGTGGAATTGGGAACA	(AG) ₈	139–173	FAM	MF716499
C50127	F: CCAGCGAGATTGTGATTGA R: CAGATGAACATGTACACACCTGC	(TC) ₈	172–184	TAMRA	MF716500
C51170	F: GTACTCATCCGGCACCATT R: AAAGCCGCAAGACCAGTAAA	(CT) ₁₀	294–322	ROX	MF716501
C53509	F: ACCATCCCAATTCCTTCTC R: GCAGCAGTGACGACTGGTAA	(TC) ₈	204–236	TAMRA	MF716502
C53824	F: CTCGATCTCCAAGGGCTAAA R: CCCCTCTCTGTGTCATGGAA	(GT) ₈	248–254	TAMRA	MF716503
C53825	F: CAACAACAGTTTTGGAGCA R: CCCTTGGGATCTCATCTTCA	(AG) ₉	282–320	HEX	MF716504
C53843	F: GGCAATAGTAGCCCAACA R: ATAACGCAACACCATCCAG	(GA) ₈	444–470	ROX	MF716505
C53920	F: AACATGAGATGCCTGCACAA R: TGGGTCTGCATGTGAAAGAA	(CT) ₉	184–192	TAMRA	MF716506
C53962	F: GGTACAAAAGAAAAACGAGCTGA R: GGTAGGCGCAGCACTATGTT	(CA) ₈	128–160	FAM	MF716507
C55683	F: TGATGAAAAGTTGGGCATGA R: CACCGTATCGTGTGGAGATG	(AT) ₉	156–182	HEX	MF716508
C55722	F: CCATCGGCCTCATAAGAAAA R: GTATGCTCTCCAGCTCCAC	(TA) ₈	206–220	TAMRA	MF716509
C57707	F: AGCAGCAAGAGCATTGGAGT R: TCATTGTTTCCAACCTCTCACAAA	(TC) ₈	225–239	ROX	MF716510
C58437	F: ACCGGTCTACACCATGACCT R: CACACAAGGCTTCTTTGCAG	(GA) ₈	252–256	ROX	MF716511
C58140	F: AACACAATCTCGTATACTATCCATCA R: GTAAATCTCGGCGTCGGTAA	(AG) ₆ ...(GA) ₉	280–304	HEX	MF716512
C59012	F: ATCGTCAACATCGTCGTGAG R: AGAGCGAGAAACCTCTTCCC	(AG) ₈	213–243	FAM	MF716513
C59078	F: CCGGCATTAAACACACTCAC R: CTACTGCTGCCGTGCATCTA	(TA) ₉	166–180	FAM	MF716514
C59155	F: TGCTTGCTTATTACCTGCC R: AATTGTTGGCGTTGGAAGAC	(AG) ₈	145–151	HEX	MF716515

TABLE 1. Continued.

Locus ^a	Primer sequences (5′–3′)	Repeat motif	Allele size range (bp)	Fluorescent dye	GenBank accession no.
C59702	F: TTAATCGTGACAGCCAGCAG R: GAGTCATCAATGCGAGGTGA	(CT) ₈	182–200	TAMRA	MF716516
C63472	F: ACGTGAAGCATGGTGCAATA R: CGGAAGCTTCTACTCGCCTA	(AT) ₉	206–234	FAM	MF716517
C65324	F: GCTCACCCTACCAACGATGT R: TCTCCACCGTCAAACCTACC	(GA) ₉	278–286	ROX	MF716518
C66329	F: CAAGGACCCGAATACTCCAA R: GATGGAATGGAAGGAGCAGA	(AT) ₉	152–170	HEX	MF716519

^aA touchdown PCR program with annealing temperature of 60–50°C was used for all loci.

(MISA) based on the Perl language (Thiel et al., 2003). We searched for SSRs with motifs ranging from mono- to hexanucleotides in size, and 4753 primer pairs were designated from 5139 putative loci using Primer3 web version 0.4.0 (Rozen and Skaletsky, 1999). A total of 220 markers comprising two nucleotides with at least eight contiguous repeat units were chosen for

screening, among which 102 primers produced clear bands with suitable fragment lengths (<500 bp) during the preliminary test with four individuals of *P. ovalifolia*.

These 102 loci were further tested with eight individuals of *P. ovalifolia*. PCR reactions were performed with three primers: a sequence-specific forward

TABLE 2. Results of initial primer screening in populations of *Primula* species.^a

Locus	<i>P. ovalifolia</i>									<i>P. tardiflora</i>			<i>P. epilosa</i>		
	OVA_EMS (n = 24)			OVA_HZG (n = 24)			OVA_BSH (n = 20)			TAR_EMS (n = 24)			EPI_PZ (n = 24)		
	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e	A	H _o	H _e
C11339	6	0.632	0.757	12	0.667	0.893	7	0.900	0.813*	3	0.136	0.210	1	0.000	0.000
C13959	4	0.313	0.619	3	0.250	0.507	3	0.056	0.517*	2	0.708	0.510	3	0.375	0.423
C14309	5	0.333	0.660	6	0.375	0.532	2	0.529	0.389	2	0.053	0.149	4	0.125	0.363*
C14710	6	0.833	0.779	7	0.708	0.764	8	0.857	0.847	3	0.813	0.679	3	0.565	0.569
C15228	5	0.579	0.731	5	0.300	0.278	3	0.300	0.296	—	—	—	1	0.000	0.000
C21614	3	0.231	0.495*	3	0.261	0.241	7	0.455	0.802	1	0.000	0.000	1	0.000	0.000
C23409	1	0.000	0.000	2	0.050	0.050	4	0.500	0.668	2	0.200	0.287	1	0.000	0.000
C23644	5	0.286	0.741*	12	0.739	0.887	4	0.842	0.698	3	0.667	0.519	2	0.083	0.082
C23746	1	0.000	0.000	2	0.125	0.120	2	0.000	0.278*	2	0.125	0.120	1	0.000	0.000
C24233	2	0.000	0.359	9	0.500	0.825*	3	0.049	0.050	2	0.375	0.311	2	0.261	0.232
C24268	10	0.500	0.870*	6	0.522	0.739	5	0.368	0.639*	6	0.429	0.606	1	0.000	0.000
C24676	3	0.087	0.405*	4	0.177	0.668*	4	0.938	0.666*	2	0.208	0.191	1	0.000	0.000
C40984	5	0.217	0.648*	2	0.895	0.508*	7	0.579	0.812*	6	0.333	0.559*	4	0.458	0.415
C45417	11	0.435	0.825	19	0.417	0.915	4	0.400	0.415*	6	0.250	0.509*	5	0.417	0.395
C46430	6	0.600	0.717	5	0.579	0.576	8	0.850	0.825	3	0.059	0.269	4	0.563	0.538
C47095	1	0.000	0.000	1	0.000	0.000	—	—	—	1	0.000	0.000	1	0.000	0.000
C48258	2	0.050	0.142	4	0.750	0.645	4	0.632	0.683*	1	0.000	0.000	4	0.652	0.654
C48475	5	0.278	0.819*	6	0.565	0.503	—	—	—	2	0.042	0.042	4	0.542	0.657
C50127	7	0.381	0.630	—	—	—	3	0.000	0.460*	3	0.750	0.573	4	0.318	0.698
C51170	6	0.750	0.761	11	0.542	0.872*	2	0.000	0.142*	3	0.476	0.441	—	—	—
C53509	11	0.632	0.871	6	0.263	0.713*	5	0.600	0.654	4	0.188	0.688*	—	—	—
C53824	7	0.778	0.764	13	0.667	0.884	—	—	—	5	0.304	0.442*	2	0.000	0.089*
C53825	1	0.000	0.000	2	0.526	0.398	3	0.368	0.547	—	—	—	2	0.375	0.311
C53843	8	0.765	0.743	10	0.870	0.876	5	0.800	0.643	3	0.435	0.445	5	0.539	0.634
C53920	3	0.044	0.086*	2	0.117	0.156	1	0.000	0.000	3	0.435	0.530	2	0.042	0.042
C53962	5	0.273	0.459	4	0.368	0.647	1	0.000	0.000	3	0.167	0.519*	2	0.046	0.206*
C55683	2	0.000	0.502	9	0.478	0.857*	3	0.063	0.576*	6	0.350	0.782*	4	0.154	0.665*
C55722	2	0.235	0.371	4	0.364	0.444	4	0.200	0.595*	1	0.000	0.000	6	0.667	0.660
C57707	8	0.810	0.829	8	0.609	0.837	3	0.083	0.344*	2	0.143	0.143	3	0.208	0.196
C58140	10	0.500	0.808*	9	0.500	0.825*	2	0.050	0.049	3	0.792	0.543*	—	—	—
C58437	3	0.684	0.522	2	0.067	0.067	2	0.600	0.495	2	0.478	0.476	—	—	—
C59012	2	0.053	0.053	3	0.191	0.180	3	0.200	0.184	3	0.762	0.638	1	0.000	0.000
C59078	2	0.263	0.309	—	—	—	1	0.000	0.000	3	0.385	0.631	2	0.048	0.048
C59155	3	0.046	0.09*	4	0.235	0.713	4	0.200	0.499*	1	0.000	0.000	1	0.000	0.000
C59702	6	0.563	0.794	6	0.409	0.538	3	0.111	0.106	4	0.200	0.407	2	0.208	0.311
C63472	4	0.154	0.542*	4	0.188	0.667*	5	0.375	0.734*	2	0.300	0.262	2	0.208	0.191
C65324	4	0.250	0.673*	—	—	—	3	0.789	0.517	4	0.292	0.645*	5	0.478	0.728*
C66329	1	0.000	0.000	3	0.042	0.451	1	0.000	0.000	6	0.524	0.436	1	0.000	0.000
Mean ^b	5.2	0.389	0.587	6.1	0.421	0.582	4.0	0.400	0.510	3.3	0.367	0.437	3.2	0.306	0.379

Note: — = unsuccessful amplification; A = number of alleles; H_e = expected heterozygosity; H_o = observed heterozygosity; n = number of individuals sampled.

^aLocality and voucher information are provided in Appendix 1.

^bMonomorphic loci are excluded.

*Significant deviation from Hardy–Weinberg equilibrium ($P < 0.05$).

primer with an M13(–21) tail at its 5' end, a sequence-specific reverse primer, and the universal fluorescent-labeled M13(–21) primer (FAM, ROX, HEX, or TAMRA; Invitrogen, Guangzhou, Guangdong, China) (Schuelke, 2000). Genomic DNA was extracted using a modified cetyltrimethylammonium bromide (CTAB) protocol (Doyle, 1991). The amplified 10- μ L mixture for SSRs included 5 μ L of Master Mix (Generay Biotech, Guangzhou, China), 0.4 mM of each primer pair, 3.2 μ L of deionized water, and 30–50 ng of genomic DNA. PCRs were run following a touchdown procedure with initial denaturation for 4 min at 94°C; followed by 10 cycles of 94°C for 35 s, 35 s at 60°C with an increment of –1°C per cycle, 45 s at 72°C; followed by 28 cycles of 94°C for 35 s, 35 s at 50°C, 45 s at 72°C; ending with an extra extension of 10 min at 72°C. PCR products were scanned by an ABI PRISM 3100 Genetic Analyzer using GeneScan 500 LIZ internal size standard (Invitrogen). Allele binning and calling were done using GeneMarker version 2.4.0 (SoftGenetics, State College, Pennsylvania, USA), and 38 primer pairs were selected for further polymorphism and transferability detection (Table 1). All of these SSR sequences have been deposited in GenBank (Table 1).

Polymorphism and transferability assessment—To assess the polymorphism level of these 38 loci, we genotyped 20–24 individuals in each of five populations from three species (Appendix 1). DNA extraction, PCR amplification, and length assessment of PCR products were performed following the procedures described above. Linkage disequilibrium among loci per population and deviation from Hardy–Weinberg equilibrium were tested using FSTAT version 2.9.3 (Goudet, 2001). We used GenAlEx 6.5 (Peakall and Smouse, 2012) to calculate the number of observed alleles per locus (A), expected heterozygosity (H_e), and observed heterozygosity (H_o).

No significant linkage disequilibrium was detected among loci after Bonferroni correction at $\alpha = 0.05$ confidence level, and some loci showed significant deviations from Hardy–Weinberg equilibrium (Table 2). The 38 EST-SSRs displayed varied genetic diversity in three populations of *P. ovalifolia* (Table 2). A , H_o , and H_e for each locus ranged from one to 19, 0 to 0.938, and 0 to 0.915, respectively (Table 2). Excluding monomorphic loci, the polymorphic EST-SSR markers showed an average A of 5.2, 6.1, and 4.0; H_e of 0.587, 0.582, and 0.51; and H_o of 0.389, 0.421, and 0.40, in each population, respectively (Table 2). Out of the 38 SSR markers, 36 loci were successfully amplified in *P. tardiflora* and 31 loci showed polymorphism, with A ranging from two to six (Table 2). Similarly, 34 loci were successfully amplified in *P. epilosa* Craib, among which 23 loci showed polymorphism, with A ranging from two to six (Table 2). Overall, most of the EST-SSR markers developed for *P. ovalifolia* could be successfully cross-amplified, leading to a high transferability in the two congeneric species.

CONCLUSIONS

We developed and characterized 38 EST-SSR markers based on transcriptome sequencing of *P. ovalifolia*, a widely distributed species of *Primula* section *Petiolaris*. These markers demonstrated high polymorphism in *P. ovalifolia*, with A ranging from one to 19, H_o from 0.000 to 0.938, and H_e from 0.000 to 0.915. Most of the markers could be successfully cross-amplified in congeneric species. These SSR makers are found to be useful tools for investigation of genetic structure and interspecific gene flow in this section.

LITERATURE CITED

- BICKLER, C., S. A'HARA, J. COTTRELL, L. ROGERS, AND J. BRIDLE. 2013. Characterisation of thirteen polymorphic microsatellite markers for cowslip (*Primula veris* L.) developed using a 454 sequencing approach. *Conservation Genetics Resources* 5: 1185–1187.
- DOYLE, J. 1991. DNA protocols for plants: CTAB total DNA isolation. In G. M. Hewitt and A. Johnston [eds.], *Molecular techniques in taxonomy*, 283–293. Springer-Verlag, Berlin, Germany.
- GOUDET, J. 2001. FSTAT (version 2.9.3): A program to estimate and test gene diversities and fixation indices. Institute of Ecology, Lausanne, Switzerland. Website <http://www2.unil.ch/popgen/softwares/fstat.htm> [accessed 27 November 2017].
- HAAS, B. J., A. PAPANICOLAOU, M. YASSOUR, M. GRABHERR, P. D. BLOOD, J. BOWDEN, M. B. COUGER, ET AL. 2013. De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols* 8: 1494–1512.
- HU, C. M., AND S. KELSO. 1996. Primulaceae. In Z. Y. Wu and P. H. Raven [eds.], *Flora of China*, vol. 15. Science Press, Beijing, China, and Missouri Botanical Garden Press, St. Louis, Missouri, USA.
- PEAKALL, R., AND P. SMOUSE. 2012. GenAlEx 6.5: Genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics (Oxford, England)* 28: 2537–2539.
- PERTEA, G., X. HUANG, F. LIANG, V. ANTONESCU, R. SULTANA, S. KARAMYCHEVA, Y. LEE, ET AL. 2003. TIGR Gene Indices clustering tools (TGICL): A software system for fast clustering of large EST datasets. *Bioinformatics (Oxford, England)* 19: 651–652.
- ROZEN, S., AND H. SKALETSKY. 1999. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: *Bioinformatics: Methods and protocols*, 365–386. Humana Press, Totowa, New Jersey, USA.
- SCHUELKE, M. 2000. An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology* 18: 233–234.
- THIEL, T., W. MICHALEK, R. VARSHNEY, AND A. GRANER. 2003. Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theoretical and Applied Genetics* 106: 411–422.
- UENO, S., Y. YOSHIDA, Y. TAGUCHI, T. UJINO-IHARA, N. KITAMOTO, M. HONJO, R. OHSAWA, ET AL. 2011. Generation of expressed sequence tags, development of microsatellite and single nucleotide polymorphism markers in *Primula sieboldii* E. Morren (Primulaceae) for analysis of genetic diversity in natural and horticultural populations. *Breeding Science* 61: 234–243.
- VAN, G. A., R. F. VAN, I. STIERS, T. SIERENS, J. BARKER, AND L. TRIEST. 2006. Isolation and characterization of microsatellite loci in primrose (*Primula vulgaris*). *Belgian Journal of Botany* 139: 261–264.
- XIE, X. F., H. F. YAN, F. Y. WANG, X. J. GE, C. M. HU, AND G. HAO. 2012. Chloroplast DNA phylogeography of *Primula ovalifolia* in central and adjacent southwestern China: Past gradual expansion and geographical isolation. *Journal of Systematics and Evolution* 50: 284–294.
- YAN, H., X. GE, C. M. HU, AND G. HAO. 2010. Isolation and characterization of microsatellite loci for the ornamental plant *Primula obconica* Hance (Primulaceae). *HortScience* 45: 314–315.
- ZHANG, L., H. F. YAN, W. WU, H. YU, AND X. J. GE. 2013. Comparative transcriptome analysis and marker development of two closely related Primrose species (*Primula poissonii* and *Primula wilsonii*). *BMC Genomics* 14: 329.

APPENDIX 1. Locality and voucher information for populations of *Primula ovalifolia*, *P. tardiflora*, and *P. epilosa* used in this study. Voucher specimens are deposited at the herbarium of South China Botanical Garden (IBSC), Guangzhou, Guangdong, China.

Species	Population code	Voucher no.	Location	Geographic coordinates	Altitude (m)	<i>n</i>
<i>Primula ovalifolia</i> Franch.	OVA_EMS	YS214	E'mei, China	29°32'55"N, 103°21'31"E	1702	24
	OVA_HZG	YS524	E'bian, China	29°02'42"N, 103°00'29"E	1793	24
	OVA_BSH	YS436	Pengzhou, China	31°14'09"N, 103°50'20"E	1900	20
<i>Primula tardiflora</i> (C. M. Hu) C. M. Hu	TAR_EMS	YS440	E'mei, China	29°32'49"N, 103°20'24"E	2448	24
<i>Primula epilosa</i> Craib	EPI_PZ	YS504	Pengzhou, China	31°11'32"N, 103°54'45"E	1300	24

Note: *n* = number of individuals sampled.