

Interactions Between *Xanthomonas* Species and *Arabidopsis thaliana*

Author: Buell, C. Robin

Source: The Arabidopsis Book, 2002(1)

Published By: The American Society of Plant Biologists

URL: <https://doi.org/10.1199/tab.0031>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First published on April 4, 2002: e0031. doi: 10.1199/tab.0031

Interactions between *Xanthomonas* Species and *Arabidopsis thaliana*

C. Robin Buell

The Institute for Genomic Research, 9712 Medical Center Drive, Rockville MD 20850, Phone: (301) 838 3558, Facsimile: (301) 838 0208, email: rbuell@tigr.org

Abbreviations: colony forming units (CFU), Columbia (Col-0), days post inoculation (dpi), hypersensitive response (HR), Landsberg *erecta* (Ler), pathogenesis-related protein 1 (PR-1), phenylalanine ammonia lyase (PAL), *Xanthomonas campestris* pv *campestris* (Xcc)

Key Words: *Xanthomonas*, hypersensitive response, tolerance, defense response, black rot

ABSTRACT

Arabidopsis has been well studied as a model plant for plant pathogen interactions. While a large portion of the literature has been devoted to interactions between *Arabidopsis* and *Pseudomonas* and *Peronospora* species, a small cadre of researchers have been making inroads on the response of *Arabidopsis* to *Xanthomonas*. Differential responses of *Arabidopsis* accessions to isolates of *Xanthomonas campestris* pv *campestris* include tolerance, a hypersensitive response, resistance without a hypersensitive response and disease which is characterized by chlorosis and necrosis. Loci that govern the recognition of *X. c. campestris* have been identified and are the focus of on-going positional cloning efforts. Signaling and other downstream molecules involved in manifestation of resistance to *Xanthomonas* have been investigated resulting in the identification of many components of the resistance response. Parallel to the characterization of the host response, molecular and genomic efforts focused on the pathogen have the potential to reveal the mechanisms by which this bacterium can invade and colonize host tissues.

Pathogens of *Arabidopsis*

Like other plant species, *Arabidopsis thaliana* is susceptible to only a limited number of pathogens including viruses, bacteria, fungi, nematodes and insect pests. Diseases resulting from these pathogens have been reported in the wild (Holub *et al.* 1994, 1995; Tsuji and Somerville 1992) suggesting both the pathogen and the host share an ecological niche and when the appropriate environmental conditions are present disease can occur. Diseases have also been observed in a laboratory setting where the host is deliberately exposed to the pathogen. Regardless of the setting, nature or the laboratory, *Arabidopsis* responds in a similar fashion as other higher plants when exposed to viral, prokaryotic, or eukaryotic pathogens.

Perhaps the more facile class of pathogen to work with in a laboratory setting are the bacterial pathogens. Bacteria have several advantages over the other classes of pathogens for pathological studies in that they can be cultured *in vitro* and have relatively rapid generation times (minutes not days). In addition, most bacterial plant pathogens elicit rapid host responses (hours to days) and have pathogenicity and avirulence factors that have been documented in other plant species thereby providing a foundation to begin work in *Arabidopsis*. Only a small number of bacterial species are pathogenic on *Arabidopsis*. The predominant bacterial pathogen utilized in *Arabidopsis* studies is *Pseudomonas syringae* and the reader is referred to other

chapters in this book describing *Arabidopsis* responses to this pathogen. Additional bacterial pathogens utilized in *Arabidopsis* research include *Erwinia* species which are causal agents of soft-rots, *Ralstonia* species which are causal agents of vascular wilts, and *Xanthomonas campestris* pathovars which are causal agents of blights and rots and are the focus of this chapter.

The Genus *Xanthomonas*

The genus *Xanthomonas* has been well described in several definitive publications (Starr 1981; Leyns *et al.*, 1984; Swings *et al.*, 1993) and the reader is referred to these for more details regarding this genus. In brief, *Xanthomonas* species are gram-negative rod-shaped aerobic bacteria with polar flagella and are found primarily in association with plants (diseased lesions, soil, plant debris) (Starr 1981). *Xanthomonas* species produce diagnostic pigments (termed xanthomonadins) and exude large amounts of xanthan gum, an extracellular polysaccharide which is used as a food additive (Williams 1980; Starr 1981; Swings *et al.*, 1993). *Xanthomonas* species, similar to other plant pathogens, can cause symptoms such as chlorosis, necrosis, cankers, and vascular wilts (Leyns *et al.*, 1984). Collectively, *Xanthomonas* isolates have been reported to cause disease on ~400 different monocotyledonous and dicotyledonous species (Leyns *et al.*, 1984), making *Xanthomonas* a significant plant pathogen. As will be discussed in more detail below, current mechanisms for control of *Xanthomonas* diseases rely extensively on breeding for resistance and/or cultural practices such as crop rotation and sanitation. An example of the extreme costs that can arise from lack of control of a *Xanthomonas* species, either through lack of breeding for resistance or ineffective control through cultural practices, is the recent epidemic of citrus canker in the United States (Brown 2001). Attempts to control citrus canker in Florida through an aggressive eradication program have cost over \$200 million to date, with no end in sight to control of the disease (Brown 2001).

Reports of *Arabidopsis*-infecting Isolates of *Xanthomonas*

With respect to infection of *Arabidopsis*, only isolates of *X. campestris* have been reported to cause disease. The species, *X. campestris*, is a large collection of isolates that

share similar biochemical and physiological phenotypes yet exhibit different host specificity and are separated into pathovars to reflect host specificity or host origin (Leyns *et al.*, 1984; Van den Mooter and Swings 1990). Table 1 lists the *X. campestris* pathovars that have been examined on *Arabidopsis* accessions. Both resistance and disease, as demonstrated by chlorosis and necrosis on inoculated leaves as well as systemic infections, have been reported on *Arabidopsis* accessions, suggesting that a full range of host responses occur in *Arabidopsis* (Tsuiji and Somerville 1988; Simpson and Johnson 1990; Davis *et al.*, 1991; Tsuiji *et al.*, 1991; Lummerzheim *et al.*, 1993; Parker *et al.*, 1993; Aufsatz and Grimm 1994).

Although a number of these pathovars can incite a response on *Arabidopsis*, the most prevalent pathovar utilized in *Xanthomonas*-*Arabidopsis* studies is the pathovar *campestris*. Various strains of this pathovar have demonstrated typical race-cultivar specificity indicative of the presence of a gene-for-gene relationship between the host and the pathogen. Thus, due to the abundance of literature with *X. campestris* pv *campestris* (Xcc), a majority of this chapter will be devoted to the interactions of *Arabidopsis* with various strains of Xcc.

Xcc and Black Rot of Crucifers

Xcc is the causal agent of black rot of crucifers (for review see Williams 1980). Xcc has a broad host range that includes a majority of members of the Cruciferae family. Numerous agronomically important species of crucifers including broccoli, brussels sprouts, cabbage, cauliflower, radish, and turnip are susceptible to Xcc. Xcc can also infect weed species within the Cruciferae (Schaad and Dianese 1981) making weed reservoirs of Xcc a factor in the control of black rot in crucifer production fields.

Xcc typically infects host plants through natural openings such as hydathodes, stomates, or wounds (for review see Williams 1980). Once the bacterium is in the vascular system, it can become systemic resulting in blackened veins and thus the name "black rot". Under favorable environmental conditions, the pathogen can become seed-borne (for review see Williams 1980). For commercial crucifers, black rot is a significant issue. The fact that the pathogen can be seed-borne has led to seed certification programs in developed countries. The certification process adds extensive cost and effort for commercial seed production.

Several lines of evidence have firmly established *Arabidopsis* as a natural host of Xcc. First, classic black rot

symptoms have been reported following infection of *Arabidopsis* with specific *Xcc* strains (Simpson and Johnson 1990; Tsuji *et al.*, 1991). Second, systemic infection occurs in highly susceptible accessions following challenge with *Xcc* (Figure 1; Buell and Somerville 1997). Third, a preference for hydathode infection, rather than stomatal infection, was observed in *Arabidopsis* leaves which is consistent with the entry route favored by *Xcc* in other cruciferous hosts (Hugouvieux *et al.*, 1998). Fourth, infections of *Arabidopsis* by *Xcc* do occur in nature as reported by Tsuji and Somerville (1992).

Pathogenicity Mechanisms of *Xanthomonas* species

There have been several excellent reviews on pathogenicity and virulence mechanisms in gram-negative plant pathogenic bacteria and the reader is referred to these articles for more in-depth discussions (Collmer 1998; Staskawicz *et al.*, 2001). With respect to *Xcc* pathogenicity on non-*Arabidopsis* crucifers, there are a number of publications by M. J. Daniels and colleagues and the reader is referred to these for a summary of *Xcc* virulence mechanisms (Daniels *et al.*, 1993; Dow and Daniels 1994). In brief, *Xcc* employs a vast array of degradative and regulatory mechanisms to parasitize its host. Extracellular polysaccharide production, cell wall degrading enzymes, proteases, as well as genes that encode for secretion of these products are involved in virulence on cruciferous hosts. In addition to these structural components, regulatory components of pathogenicity and virulence have been identified in *Xanthomonas* (for review see Daniels *et al.*, 1993; Dow and Daniels 1994).

Although there is a substantial amount of information cur-

rently available on pathogenicity mechanisms in *Xanthomonas*, the level of information is about to increase exponentially. The first genome of a plant pathogen completely sequenced was *Xylella fastidiosa* (Simpson *et al.*, 2000) which is the causal agent of citrus variegated chlorosis and is closely related to *Xanthomonas*. Not surprisingly, several components of virulence and pathogenicity in *Xcc* have orthologs in *X. fastidiosa*, including genes involved in xanthan gum (extracellular polysaccharide) production, regulation of pathogenicity factors, and the type II secretion system that is necessary for export of degradative enzymes (Dow and Daniels 2000; Simpson *et al.*, 2000).

Even more exciting is the new genome sequencing projects focused on *Xanthomonas* species. *X. axonopodis* pv *citri*, the causal agent of citrus canker, is being sequenced by the Organization for Nucleotide Sequencing and Analysis (ONSA), a consortium of laboratories in Brazil (Kamoun and Hogenhout 2001; <http://genoma4.iq.usp.br/Xanthomonas/>). This pathogen is closely related to *Xcc* and major insights into the virulence mechanisms of *Xanthomonas* will be revealed from this project. The other *Xanthomonas* genome project is focused on *Xcc* (Kamoun and Hogenhout 2001; http://genoma.fcav.unesp.br/xcc-campestris/home/xcc_menu.html). In addition, the genome of *Ralstonia solanacearum* (Kamoun and Hogenhout 2001; http://www.genoscope.cns.fr/externe/English/Projets/Projet_Y/Y.html) and *Pseudomonas syringae* pv *tomato* (<http://www.tigr.org>; <http://ppi.cornell.edu/>), both pathogens of *Arabidopsis*, are the focus of genome sequencing efforts and these genomic sequences will provide additional resources for the identification of pathogenicity mechanisms in *Xanthomonas*.

Genetic Basis for Resistance to *Xanthomonas*

Responses in *Arabidopsis* to *Xanthomonas* infections are dependent on genetic factors present in the host and in the pathogen. Several types of resistance responses have been documented in *Arabidopsis*-*Xanthomonas* interactions. These include tolerance, resistance mediated through a hypersensitive response (HR) and non-HR-mediated resistance. Genes involved in the resistance response can be classified into three classes: (1) those involved in the recognition of the pathogen (also known as R genes), (2) genes involved in signal transduction events, and (3) genes involved in the suppression of pathogen growth and development (defense response genes). Currently, five genes have been identified genetically that are hypothesized to be involved in recognition of the pathogen (Table 2; Tsuji *et al.*, 1991; Buell and Somerville

Table 1. Pathovars of *X. campestris* tested on *Arabidopsis*.

Pathovar	Reference
<i>aberrans</i>	Parker <i>et al.</i> , 1993
<i>armoraciae</i>	Davis <i>et al.</i> , 1991
<i>carotae</i>	Simpson & Johnson 1990
<i>campestris</i>	Aufsatz & Grimm 1994
	Davis <i>et al.</i> , 1991
	Lummerzhim <i>et al.</i> , 1993
	Parker <i>et al.</i> , 1993
	Simpson & Johnson 1990
	Tsuji & Somerville 1988
	Tsuji <i>et al.</i> , 1991
	Tsuji & Somerville 1992
<i>raphani</i>	Parker <i>et al.</i> , 1993
<i>vesicatoria</i>	Lummerzhim <i>et al.</i> , 1993
<i>vitans</i>	Lummerzhim <i>et al.</i> , 1993

1997; Godard *et al.*, 2000). These have been termed *RXC* for reaction to *Xanthomonas campestris* and are named *RXC1-5*. Additional genes have been identified in both the signal transduction pathway and in the suppression of pathogen growth/development and are discussed in more detail below (see also Table 3).

At the present, only a single corresponding gene for avirulence has been reported from a crucifer-infecting

Xanthomonas isolate. This avirulence gene, *avrXca*, was isolated from *X. c. raphani* and confers incompatibility in a large number of *Arabidopsis* accessions (Parker *et al.*, 1993). *avrXca* encodes a large protein (~67 kDa) that is possibly secreted from the bacterium and the upstream region of *avrXca* contains a putative *hrp* box suggesting *avrXca* is controlled by the *hrp* regulatory system. The *hrp* pathway is an essential component of pathogenicity and



Figure 1. Systemic infection of Xcc2D520 in differential *Arabidopsis* accessions. Select basal leaves (3-5) of 3-4 week old plants were infiltrated with Xcc2D520 ($\sim 10^7$ CFU ml $^{-1}$). Photographs were taken 23 d.p.i. The accessions from left to right are Col-0, Pr-0, and Ler.

virulence in bacterial plant pathogens and the reader is referred to these reviews for more information (Alfano and Collmer 1997; Cornelis and Van Gijsegem 2000). The lack of identification of corresponding avirulence gene(s) in *Xcc* isolates may reflect either technical difficulties in screening for such a gene or a lack of the gene in this pathogen. Genome sequencing efforts (see above) should provide valuable insight into this unresolved issue.

Tolerance to *Xcc*

Using the 2D520 strain of *Xcc*, Tsuji *et al.*, (1991) reported tolerance to the pathogen in the Columbia accession (Col-0). While the Pr-0 accession developed chlorosis and necrosis following infiltration with *Xcc*2D520, Col-0 remained asymptomatic. As the *in planta* bacterial levels were indistinguishable between these two accessions, the response in Col-0 was termed tolerance as limited bacterial growth was supported in Col-0 without the development of symptoms. Thus, an uncoupling of symptom formation and pathogen growth is seen in the interaction of Col-0 and Pr-0 with *Xcc* 2D520. Genetic mapping efforts indicated that a single dominant gene, termed *RXC1* (Tsuji *et al.*, 1991), confers tolerance. *RXC1* was been mapped to the lower arm of chromosome 2 (Buell and Somerville 1997).

The biochemical and physiological mechanism by which Col-0 is able to tolerate 2D520 is not well understood. This tolerance does not involve the synthesis of the phytoalexin, camalexin (Tsuji *et al.*, 1992). At the molecular level, examination of the mRNA accumulation levels of several genes involved in defense responses did not reveal an accumulation of mRNA unique to Col-0 and thus no genes could be correlated specifically with tolerance (Buell and Somerville 1995). Although an accumulation of pathogenesis-related protein 1 (PR-1) mRNA was observed, it was expressed at high levels in both Col-0 and Pr-0.

Non-HR Resistance to *Xcc*

Previous work with *Xcc*2D520 utilized the differential accessions Col-0 and Pr-0. A further survey of *Arabidopsis* accessions revealed additional susceptible accessions. Specifically, the accession, Landsberg *erecta* (Ler), is highly susceptible to *Xcc*2D520 (Tsuji and Somerville 1988; Buell and Somerville 1997). Inoculation of Ler results in severe chlorosis and necrosis which can spread systemically resulting in necrosis of the entire plant (Figure 1; Buell and Somerville 1997). A qualitative difference in the susceptible response is apparent between Pr-0 and Ler. Whereas Pr-0 lesions are primarily chlorotic in nature, Ler lesions are more necrotic with darkening of the vascular tissue that is absent from Pr-0 lesions (Figure 2). *In planta* bacterial levels in Ler are 10-100- fold higher than in Col-0 and in Pr-0, revealing a suppression of bacterial growth in Col-0 and Pr-0 in comparison to Ler (Buell and Somerville 1995, 1997). Thus, Col-0 exhibits two responses to *Xcc*2D520: tolerance to limited bacterial growth (as compared to the susceptible accession Pr-0) and resistance without a HR (when compared to the highly susceptible accession Ler).

Examination of the genetic basis for resistance to the 2D520 isolate indicated three loci, *RXC2*, *RXC3*, and *RXC4*, were involved in resistance (Buell and Somerville 1997). The major locus for resistance, *RXC2*, behaved as a single dominant gene whereas the other two loci, *RXC3* and *RXC4*, functioned in a digenic manner. These loci have been placed on the genetic map and map to chromosome 5 (*RXC2*, *RXC3*) and chromosome 2 (*RXC4*) (Buell and Somerville 1997). Although *RXC2* and *RXC3* map to distinct locations on chromosome 5, *RXC4* maps to the same region of chromosome 2 as *RXC1*. Due to limited recombination events, it is not clear whether *RXC1* and *RXC4* are the same gene or simply map to similar regions of the chromosome.

Table 2. Major loci involved in the recognition of *X. campestris*.

Locus	Phenotype	Map position	Reference
<i>RXC1</i>	Tolerance to <i>Xcc</i> 2D520	Chromosome 2	Tsuji <i>et al.</i> , 1991 Buell & Somerville 1997
<i>RXC2</i>	Non-HR resistance (monogenic) to <i>Xcc</i> 2D520	Chromosome 5	Buell & Somerville 1997
<i>RXC3</i>	Non-HR resistance (digenic) to <i>Xcc</i> 2D520	Chromosome 5	Buell & Somerville 1997
<i>RXC4</i>	Non-HR resistance (digenic) to <i>Xcc</i> 2D520	Chromosome 2	Buell & Somerville 1997
<i>RXC5</i>	HR to <i>Xcc</i> 147	Unknown	Godard <i>et al.</i> , 2000

Hypersensitive Response: Interactions with Xcc750

The Xcc750 isolate has been shown to induce a HR-like response on resistant accessions of *Arabidopsis* such as Col-0 while causing chlorosis on susceptible accessions such as Oy-0 (Aufsatz and Grimm 1994). The HR observed on Col-0 is coupled with a lack of significant bacterial growth, consistent with the HR observed in *Arabidopsis* with other pathogens such as *Pseudomonas*. Resistance in *Arabidopsis* is associated with high basal expression and an induction of expression of a low molecular weight protein termed ECS1 (formerly named CXc750; Aufsatz and Grimm 1994). Subsequent work using transgenic plants and polyclonal antibodies generated to ECS1 revealed it is associated with the cell wall (Aufsatz *et al.*, 1998). Although ECS1 had been tightly correlated with the resistance phenotype, definitive genetic and transgenic experiments have revealed ECS1 is not a resistance gene for Xcc750 but instead is linked to a locus involved in resistance to Xcc750.

Hypersensitive Response: Interactions with Xcc147

The response of *Arabidopsis* to the HR-inducing strain 147 of Xcc has been described in a series of papers by the Roby laboratory and other researchers. Xcc147 induces a HR on resistant accessions of *Arabidopsis* such as Col-0 (Lummerzheim *et al.*, 1993). At the molecular level, a correlation between mRNA accumulation of two defense genes, phenylalanine ammonia lyase (PAL) and β -1,3-glucanase, and the HR was observed (Lummerzheim *et al.*, 1993). Lummerzheim *et al.* (1993) also examined transcript accumulation of a basic class I chitinase and an ascorbate peroxidase and although they were up-regulated upon infection with Xcc147, they were also expressed in the compatible interaction with

Xcc8004 revealing a lack of specificity with the HR. There are four classes of chitinase (I to IV) and Gerhardt *et al.* (1997) were able to demonstrate that an extracellular class IV chitinase was expressed in *Arabidopsis* leaves following challenge with Xcc147. Unfortunately, class IV chitinase expression was not examined in leaves challenged with a compatible Xcc isolate and no conclusion can be made regarding the specificity of class IV chitinase expression in the HR. In healthy plants, the class IV chitinase was only expressed in siliques, suggesting a specific developmental pattern for this gene.

While the above studies with Xcc147 were performed using whole plant studies, a number of new genes involved in the HR to Xcc147 have been identified using cell suspension cultures. A cDNA clone encoding a putative sulfotransferase was identified from a library constructed from pathogen-challenged cell suspension cultures. The sulfotransferase, RaR047, is similar to flavonol sulfotransferases from *Flaveria* species (Lacomme and Roby 1996). RaR047 is developmentally regulated as mRNA could only be detected in cell cultures and in developing aerial tissues (Lacomme and Roby 1996). Expression of RaR047 was strongly induced in young seedlings by treatment with several elicitors of defense responses, including salicylic acid and jasmonic acid. The induction of RaR047 expression by elicitors is consistent with the induction of expression observed with pathogen treatment of *Arabidopsis* leaves (Lacomme and Roby 1996). Inoculation with either Xcc147 or an incompatible strain of *Pseudomonas syringae* pv *maculicola* (M2) resulted in accumulation of RaR047 mRNA. Compatible strains of Xcc (Xcc8004) and *P. s. maculicola* (M4) were also able to induce accumulation of RaR047 although the extent of accumulation was substantially less than that in incompatible interactions. Although there is a correlation of RaR047 expression with resistance (Lacomme and Roby 1996), the function of the encoded protein is unknown. It is speculated that RaR047 may function in synthesis of a molecule that is involved in signal transduction or that RaR047 functions directly in the suppression of pathogen growth (Lacomme and Roby 1996). Definitive biochemical studies will establish an enzymatic function for RaR047 and allow placement of this sulfotransferase in the defense response pathway.

In addition to sulfotransferase, cinnamoyl-CoA reductase, an enzyme involved in lignification, is associated with the development of the HR to Xcc147. Two cinnamoyl-CoA reductase genes, AtCCR1 and AtCCR2, exhibit different substrate specificities and do not exhibit coordinate regulation during development and pathogen challenge (Lauvergeat *et al.*, 2001). AtCCR1 is expressed throughout normal development and is expressed more in highly lignified tissues such as stems in comparison to leaf tissue. In contrast, AtCCR2 is weakly expressed in developing tissue. Following challenge with Xcc147, AtCCR2 and not




Figure 2. Comparison of symptom formation in differential *Arabidopsis* accessions following *X. c. campestris* infection. Leaves of 3-4 week old plants were infiltrated with a suspension of Xcc2D520 ($\sim 10^7$ CFU ml⁻¹). Photographs were taken at 6 d.p.i. (A) Col-0, (B) Pr-0, and (C) Ler.

AtCCR1, was highly upregulated. *AtCCR2* expression was correlated with the HR as *AtCCR2* expression was not detectable in compatible tissues. *AtCCR2* was also inducible by treatment with salicylic acid, further supporting its role in defense responses. Although these two genes share substantial similarity (81.6 % identity at the amino acid level) they clearly have distinct expression profiles *in planta*. With the completion of the *Arabidopsis* Genome Initiative (2000), it would be interesting to examine the regulatory regions of these genes for promoter sequences that may reflect the differential expression patterns.

An additional set of genes involved in the HR to Xcc147 was described by Lacomme and Roby in 1999. A total of 27 cDNA clones (*Athsr*) was identified by differential screening of a cDNA library that was constructed from cell suspension cells challenged with Xcc147. The 27 clones were then grouped into "cDNA clone families" based on cross-hybridization and sequencing results. The *Athsr2* cDNA family was the most heavily represented family with 16 clones identified. The *Athsr2* cDNA family encodes voltage-dependent anion channel proteins (Lacomme and Roby 1999) that are localized in the mitochondrion and function to transport small molecules across the mitochondrial membrane. Membrane integrity and control over solute movement across the membrane are central components in apoptotic cell death and voltage-dependent anion channel proteins have been well studied in mammalian apoptosis (for review see Green and Reed 1998; Boya *et al.*, 2001). The *Athsr3* cDNA family, represented by 6 clones from the screening, encodes an alternative oxidase (Lacomme and Roby 1999). In tobacco, Chivasa *et al.* (1997) demonstrated that alternative oxidase was a com-

ponent of the signal transduction pathway that leads to the HR following challenge with tobacco mosaic virus. While *Athsr2* and *Athsr3* were associated with the mitochondrion, *Athsr4* (represented by a single clone) encodes a protein with similarity to a Rab GDP-dissociation inhibitor protein. Rab proteins are involved in membrane/vesicle trafficking and Rab GDP-dissociation inhibitor proteins are intimately involved in the regulation of Rab proteins and ultimately in the regulation of membrane/vesicle trafficking (for a review on Rab proteins and their regulation see Stenmark and Olkkonen, 2001). The other 3 cDNA families (*Athsr5*, 6, 7), also represented by a single clone, did not have significant similarity to any entries in the database (Lacomme and Roby 1999).

The expression patterns of the *Athsr2-7* cDNAs were assessed in cell suspension cultures challenged with differential isolates of *Xanthomonas*: Xcc147 (avirulent), Xcc8004 (virulent), and Xcc 8B2 (control). All of the cDNAs were exclusively or preferentially expressed in cells challenged with Xcc147, confirming the success of the differential screening and suggesting a role for these genes in the HR (Lacomme and Roby 1999). Further research using a biochemical and a functional genomics approach will be essential for defining the roles of these genes in the HR.

The last gene identified from Xcc147-challenged cell suspension cells is the *AtMYB30* gene which encodes an orthologue of the myb oncogene (Daniel *et al.*, 1999; Lacomme and Roby 1999). MYB proteins are transcription factors and are involved in many cellular functions including plant defense responses (Yang and Klessig 1996). *AtMYB30* encodes a 323 amino acid protein and consistent with other MYB proteins contains MYB repeats in its

Table 3. Components of the signal transduction and defense response pathway in incompatible interactions between *Arabidopsis* and *X. c. pv campestris*

Component	Putative Identification	Function	Type of Incompatible Interaction	Reference
<i>AtCCR2</i>	Cinnamoyl-CoA reductase	Lignification	HR	Lauvergeat <i>et al.</i> , 2001
<i>AtMYB30</i>	MYB domain containing protein	Transcription Factor	HR	Daniel <i>et al.</i> , 1999
<i>Ap3.3a</i>	Centrin	Cytoskeleton	HR	Cordeiro <i>et al.</i> , 1998
<i>Athsr2 family</i>	Mitochondrial voltage-dependent anion channel proteins	Membrane potential and transport	HR	Lacomme & Roby 1999
<i>Athsr3 family</i>	Alternative oxidase	Cyanide insensitive respiration	HR	Lacomme & Roby 1999
<i>Athsr4</i>	Rab-GDP dissociation inhibitor protein	Regulation of membrane/vesicle trafficking	HR	Lacomme & Roby 1999
<i>Athsr5</i>	Unknown	Unknown	HR	Lacomme & Roby 1999
<i>Athsr6</i>	Unknown	Unknown	HR	Lacomme & Roby 1999
<i>Athsr7</i>	Unknown	Unknown	HR	Lacomme & Roby 1999
<i>BG2</i>	β -1,3-glucanase	Pathogen cell wall degradation	HR	Lummerzheim <i>et al.</i> , 1993
<i>COP2</i>	Unknown	Apical hook formation; light response	HR	Buell, unpublished
<i>ECS1</i>	Cell wall protein	Unknown	HR	Aufsatz & Grimm 1994; Aufsatz <i>et al.</i> , 1998
<i>HLS1</i>	Acetyltransferase	Apical hook formation	Non-HR	Buell 1998
<i>HXC2</i>	Unknown	Unknown	HR	Godard <i>et al.</i> , 2000
<i>PAL</i>	Phenylalanine ammonia lyase	Key enzyme in synthesis of secondary metabolites	HR	Lummerzheim <i>et al.</i> , 1993
<i>PR1</i>	Pathogenesis-related protein 1	Unknown; defense-associated	HR; Non-HR resistance	Buell & Somerville 1995
<i>RaR047</i>	Sulfotransferase	Unknown	HR	Lacomme & Roby 1996

N-terminus (Daniel *et al.*, 1999). At the developmental level, *AtMYB30* is expressed at extremely low levels and is detectable only in developing seedlings. Expression of *AtMYB30* was induced by challenge with avirulent *Xcc147* cells and avirulent *P. syringae* strains. Expression was correlated with phenotypic expression of the HR, as *AtMYB30* expression was not detectable in the susceptible accession Sf-2 or with virulent strains of *Xcc* or *P. syringae*. Expression of *AtMYB30* was also examined in various lesion-mimic mutants (*lsd*) which develop HR-like lesions in the absence of pathogens. In three *lsd* mutants (*lsd3*, *lsd4*, *lsd5*), a positive correlation between *AtMYB30* expression and lesion presence was found (Daniel *et al.*, 1999), further supporting the hypothesis that *AtMYB30* is a positive regulator of cell death.

Using differential display, Cordeiro *et al.*, (1998) were able to identify two novel genes involved in the early stages of the HR of *Arabidopsis* to *Xcc147*. One gene, *ap3.3a*, encodes a protein with similarity to centrin, a cytoskeletal protein. Similar to the other centrins, the *Arabidopsis ap3.3a* protein contains four regions that are hypothesized to be involved in Ca^{2+} -binding (Cordeiro *et al.*, 1998). Expression of *ap3.3a* was measured in compatible and incompatible interactions with *Xcc*. A rapid (1 hr) up-regulation of *ap3.3a* was observed in the incompatible interaction with *Xcc147* whereas accumulation of *ap3.3a* was not observed in the compatible interaction until 1-2 dpi. A second gene identified through differential display is the *ap4.3a* gene which encodes a protein with multiple kinase domains. *ap4.3a* is rapidly induced (15 min) upon challenge with *Xcc*, although specificity of the induction was not apparent between compatible and incompatible *Xcc* strains.

Mutational Approaches to Characterize Signal Transduction and Defense Pathway Components in *Arabidopsis*-*Xanthomonas* Interactions

In a complementary approach to differential screening of cDNA libraries, Roby and colleagues identified a mutant of *Arabidopsis* that is deficient in manifestation of the HR following challenge with *Xcc147*. In a screen of 20,700 M_2 plants, a single recessive mutant (*hxc-2*) was identified that is unable to mount a full HR (Godard *et al.*, 2000). Inoculation of *hxc-2* plants results initially in necrosis (48 hr) at the infection site which is followed by a chlorotic halo and ultimately, spreading chlorosis. In the *hxc-2* plants, bacterial growth is unrestricted and is indistinguishable from levels in the susceptible accession Sf-2. The *hxc-2*

mutant is able to mount an effective HR against avirulent strains of *P. syringae*, suggesting that the wild-type *HXC2* locus is specific for *Xcc*-mediated HR. At the biochemical level, the *hxc-2* mutants retain full responsiveness to salicylic acid which is an inducer of systemic acquired resistance (Godard *et al.*, 2000). However, accumulation of salicylic acid is impaired following challenge with *Xcc147*. At the molecular level, expression levels of several defense genes (*PAL-1*, *PR-1*, *PDF-1.2*) are altered in *hxc-2*, suggesting a role for *hxc-2* in the signal transduction pathway leading to resistance. Genetic mapping experiments place *HXC2* on chromosome 3 near the Major Recognition Gene Complex MRC-F (Godard *et al.*, 2000). Through allelism tests, it was demonstrated that *HXC2* is not the determinant of specificity to *Xcc147* and that a second gene, *RXC5*, governs recognition of the pathogen.

Ethylene has been implicated in numerous plant responses, including the contrasting responses of defense in incompatible interactions and symptom formation in compatible interactions (Abeles *et al.*, 1992). The availability of mutants in ethylene-associated developmental processes has provided valuable reagents to dissect the role of ethylene in plant-pathogen interactions (for a recent review on the ethylene response pathway see Bleecker and Kende 2000). Inoculation of the *ethylene insensitive2* (*ein2*) perception mutant with a virulent isolate of *Xcc* resulted in plants with reduced symptoms (Bent *et al.*, 1992), suggesting EIN2 is involved in symptom formation. *EIN2* has been cloned and encodes a novel protein (Alonso *et al.*, 1999). However, although EIN2 has been placed in the ethylene response pathway through epistasis tests, its function at the biochemical and molecular level is still unknown (Bleecker and Kende 2000). Another ethylene-related mutant, *hookless1* (*hls1-1*), is unable to form an apical hook during the triple response in germinating seedlings (Guzman and Ecker 1990). Infection of *hls1-1* with *Xcc2D520* results in disease symptoms and an increase in bacterial growth in comparison to the asymptomatic parental accession (Col-0), suggesting HLS1 functions in resistance, not tolerance, to *Xcc2D520* (Buell 1998; Buell, unpublished). The *HLS1* gene encodes an acetyltransferase-like protein yet a specific enzymatic function for the protein has not been determined either in apical hook formation or disease resistance (Lehman *et al.*, 1988). A second mutant in apical hook formation, *constitutive photomorphogenic 2* (*cop2*; Hou *et al.*, 1993), is also suppressed in resistance to *Xcc2D520* (Buell, C.R., unpublished) consistent with the phenotype observed in the *hls1-1* apical hook mutant. *COP2* has not been cloned but with the recent completion of the *Arabidopsis* genome (Arabidopsis Genome Initiative, 2000), this can be accomplished in the near future. Continued work focused on resolving the biochemical function of these proteins will be

valuable to dissecting their role in pathogen responses.

SUMMARY

In the dozen or so years since *Xanthomonas* was first reported as a pathogen of *Arabidopsis*, significant inroads into how this bacterium is able to parasitize this weed species have been made. A survey of *Xanthomonas* isolates has revealed that *Arabidopsis* is not limited to a single resistance mechanism to prevent pathogen growth and development. Instead, a range of responses is present in *Arabidopsis* to deal with *Xanthomonas* which may be reflective of the natural co-evolution of the host and pathogen. With the complete sequence of *Arabidopsis*, along with the ongoing functional genomic projects, we are now poised to identify all of the components involved in the response to this pathogen. Coupled with the pending complete genome sequence of several *Xanthomonas* species, especially *Xcc*, we will have a comprehensive resource to dissect the pathogen component of the interaction.

ACKNOWLEDGEMENTS

The critical review of this manuscript by Catherine Ronning and Elizabeth White are greatly appreciated. The assistance of Ama Kwamena-Poh and Elizabeth White in preparation of the manuscript is greatly appreciated. The work on *Xcc*2D520 was supported by a grant from the U. S. Department of Agriculture (98-35303-6666) to C. R. B.

REFERENCES

- Abeles F., Morgan P.W., and Saltveit, M.E.** (1992) Ethylene in Plant Biology, F. Abeles, P.W. Morgan, and M.E. Saltveit, eds, (Academic Press, San Diego).
- Alonso, J.M., Hirayama, T., Roman, G., Nourizadeh, S. and Ecker, J.R.** (1999). EIN2, a bifunctional transducer of ethylene and stress responses in *Arabidopsis*. *Science* **284**, 2148-2152.
- Arabidopsis Genome Initiative.** (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**, 796-815.
- Alfano, J. R. and Collmer, A.** (1997). The type III (Hrp) secretion pathway of plant pathogenic bacteria: Trafficking harpins, avr proteins, and death. *J. Bact.* **179**, 5655-5662.
- Aufsatz, W and Grimm, C.** (1994). A new pathogen-inducible gene of *Arabidopsis* is expressed in an eco-type-specific manner. *Plant Mol. Biology* **25**, 229-239.
- Aufsatz, W, Amry, D. and Grimm, C.** (1998). The *ECS1* gene of *Arabidopsis* encodes a plant cell wall-associated protein and is potentially linked to a locus influencing resistance to *Xanthomonas campestris*. *Plant Mol. Biology* **38**, 965-976.
- Bent, A.F., Innes, R.W., Ecker, J.R. and Staskawicz, B.** (1992). Disease development in ethylene-insensitive *Arabidopsis thaliana* infected with virulent and avirulent *Pseudomonas* and *Xanthomonas* pathogens. *Mol. Plant-Microbe Interact.* **5**, 372-378.
- Bleecker, A.B. and Kende, H.** (2000) Ethylene: A gaseous signal molecule in plants. *Annu. Rev. Cell Dev. Biol.*, **16**, 1-18.
- Boya, P., Roques, B., and Kroemer, G.** (2001) Viral and bacterial proteins regulating apoptosis at the mitochondrial level. *EMBO J.* **20**, 4325-4331.
- Brown, K.** (2001). Florida fights to stop citrus canker. *Science* **292**, 2275-2276.
- Buell, C. R. and Somerville, S.C.** (1995). Expression of defense-related and putative signaling genes during tolerant and susceptible interactions of *Arabidopsis* with *Xanthomonas campestris* pv. *campestris*. *Mol. Plant-Microbe Interact.* **8**, 435-443.
- Buell, C. R. and Somerville, S.C.** (1997). Use of *Arabidopsis* recombinant inbred lines reveals a monogenic and a novel digenic resistance mechanism to *Xanthomonas campestris* pv *campestris*. *Plant J.* **12**, 21-29.
- Buell, C. R.** (1998). *Arabidopsis*: A weed leading the field of plant-pathogen interactions. *Plant Physiol. Biochem.* **36**, 177-186.
- Chivasa, S., and Carr J.P.** (1998) Cyanide restores N gene-mediated resistance to tobacco mosaic virus in transgenic tobacco expressing salicylic acid hydroxylase. *Plant Cell* **10**, 1489-1498.
- Collmer, A.** (1998). Determinants of pathogenicity and avirulence in plant pathogenic bacteria. *Curr. Opin. Plant Biology* **1**, 329-335.
- Cordeiro, M.C., Piqueras, R., Oliveira, de D. and Castresana, C.** (1998). Characterization of early induced genes in *Arabidopsis thaliana* responding to bacterial inoculation: identification of centrin and of a novel protein with two regions related to kinase domains. *FEBS Letters* **434**, 387-393.
- Cornelius, G.R., and Van Gijsegem, F.** (2000). Assembly and function of type III secretory systems. *Annu. Rev. Microbiol.* **54**, 735-774.
- Daniel, X., Lacomme, C., Morel, J.B. and Roby, D.** (1999). A novel *myb* oncogene homologue in *Arabidopsis thaliana* related to hypersensitive cell death. *Plant J.* **20**, 57-66.

- Daniels, M.J., Barber, C.E., Dow, J.M., Han, B., Liddle, S.A., Newman, M.A., Parker, J.E., Soby, S.D. and Wilson, T.G.J.** (1993). Plant and bacterial genes involved in interactions between *Xanthomonas* and crucifers. In *Advances in Molecular Genetics of Plant-Microbe Interactions*, E.W. Nester and D.P.S.Verma, eds (Kluwer Academic Publishers, Netherlands), pp 423-433.
- Davis, K.R., Schott, E. and Ausubel, F.** (1991). Virulence of selected phytopathogenic pseudomonads in *Arabidopsis thaliana*. *Mol. Plant-Microbe Interact.* **4**, 477-488.
- Dow, J.M., and Daniels, M.J.** (1994). Pathogenicity determinants and global regulation of pathogenicity of *Xanthomonas campestris* pv *campestris*. *Curr. Top. Microbiol. Immunol.* **192**, 26-41.
- Dow, J.M. and Daniels, M.J.** (2000). *Xylella* genomics and bacterial pathogenicity to plants. *Yeast* **17**, 263-271.
- Gerhardt, L.B.A., Sachetto-Martins, G., Contarini, M.G., Sandroni, M., Ferreira, R.P., Lima, V.M., Cordeiro, M.C., Oliveira, D. and Margis-Pinheiro, M.** (1997). *Arabidopsis thaliana* class IV chitinase is early induced during the interaction with *Xanthomonas campestris*. *FEBS Letters* **419**, 69-75.
- Godard, F., Lummerzheim, M., Saindrenan, P., Balague, C. and Roby, D.** (2000). *hxc2*, an *Arabidopsis* mutant with an altered hypersensitive response to *Xanthomonas campestris* pv. *campestris*. *Plant J.* **24**, 749-761.
- Green D.R., and Reed, J.C.** (1998) Mitochondria and apoptosis. *Science* **281**, 1309-1312.
- Guzman, P. and Ecker, J.R.** (1990). Exploiting the triple response of *Arabidopsis* to identify ethylene-related mutants. *Plant Cell* **2**, 513-523.
- Holub, E.B., Beynon, J.L., and Crute, I.R.** (1994) Phenotypic and genotypic characterizations of interactions between isolates of *Peronospora parasitica* and accessions of *Arabidopsis thaliana*. *Molec. Plant Microbe Interactions*, **7**, 223-239.
- Holub, E.B., Brose, E., Tor, M., Clay, C., Crute, I.R., and Beynon, J.** (1995). Phenotypic and genotypic variation in the interaction between *Arabidopsis thaliana* and *Albugo candida*. *Molec. Plant Microbe Interactions*, **8**, 916-928.
- Hugouvieux, V., Barber, C.E. and Daniele, M.J.** (1998). Entry of *Xanthomonas campestris* pv. *campestris* into hydathodes of *Arabidopsis thaliana* leaves: A system for studying early infection events in bacterial pathogenesis. *Mol. Plant-Microbe Interact.* **11**, 537-543.
- Hou, Y., Arnim, A.G. and Deng, X.** (1993). A new class of *Arabidopsis* constitutive photomorphogenic genes involved in regulating cotyledon development. *Plant Cell* **5**, 329-339.
- Kamoun, S. and Hogenhout, S.** (2001). Agricultural Microbes Genome 2: First glimpses into the genomes of plant-associated microbes. *Plant Cell*, 451-458.
- Lacomme, C. and Roby, D.** (1996). Molecular cloning of a sulfotransferase in *Arabidopsis thaliana* and regulation during development and in response to infection with pathogenic bacteria. *Plant Mol. Biology* **30**, 995-1008.
- Lacomme, C. and Roby, D.** (1999). Identification of new early markers of the hypersensitive response in *Arabidopsis thaliana*. *FEBS Letters* **459**, 149-153.
- Lauvergeat, V., Lacomme, C., Lacombe, E., Lasserre, E., Roby, D. and Grima-Pettenati, J.** (2001). Two cinnamoyl-CoA reductase (CCR) genes from *Arabidopsis thaliana* are differently expressed during development and in response to infection with pathogenic bacteria. *Phytochemistry* **57**, 1187-1195.
- Lehman, A., Black, R. and Ecker, J.R.** (1996). *HOOKLESS1*, an ethylene response gene, is required for differential cell elongation in the *Arabidopsis* hypocotyl. *Cell* **85**, 183-194.
- Leyns, F., De Cleene, M., Swings, J and De Ley, J.** (1984). The host range of the genus *Xanthomonas*. *Botanical Review* **50**, 308-356.
- Lummerzheim, M., Oliviera, de D., Castresana, C., Miguens, F.C., Louzada, E., Roby, D., Van Montagu, M and Timmerman, B.** (1993). Identification of compatible and incompatible interactions between *Arabidopsis thaliana* and *Xanthomonas campestris* pv. *campestris* and characterization of the hypersensitive response. *Mol. Plant-Microbe Interact.* **6**, 532-544.
- Parker, J.E., Barber, C.E., Mi-Jiao, F. and Daniels, M.J.** (1993). Interaction of *Xanthomonas campestris* with *Arabidopsis thaliana*: Characterization of a gene from *X.c.* pv. *raphani* that confers avirulence to most *A. thaliana* accessions. *Mol. Plant-Microbe Interact.* **6**, 216-224.
- Schaad, N.W. and Dianese, J.C.** (1981). Cruciferous weeds as sources of inoculum of *Xanthomonas campestris* in black rot of crucifers. *Phytopathology* **71**, 1215-1220.
- Simpson, R.B. and Johnson, L.J.** (1990). *Arabidopsis thaliana* as a host for *Xanthomonas campestris* pv. *campestris*. *Mol. Plant-Microbe Interact.* **3**, 233-237.

- Simpson, A.J.G., Reinach, F.C., Arruda, P., Abreu, F.A., Acenicio, M., Alvenga, R., Alves, M.C., Araya, J.E., Baia, G.S., Baptista, C.S., Barros, M.H., Bonaccorsi, E.D., Bordin, S., Bove, J.M., Briones, M.R.S., Bueno, M.R.P., Camargo, A.A., Camargo, L.E.A., Carraro, D.M., Carrer, H., Colauto, N.B., Colombo, C., Costa, F.F., Costa, M.C.R., Costa-Neto, C.M., Coutinho, L.L., Cristofani, M., Dias-Neto, E., Docena, C., El-Dorri, H., Facincani, A.P., Ferreira, A.J.S., Ferreira, V.C.A., Ferro, J.A., Fraga, J.S., Franca, S.C., Franco, M.C., Frohme, M., Furlan, L.R., Garnier, M., Goldman, G.H., Goldman, M.H.S., Gomes, S.L., Gruber, A., Ho, P.L., Hoheisel, J.D., Junqueira, M.L., Kemper, E.L., Kitajima, J.P., Krieger, J.E., Kuramae, E.E., Laigret, F., Lambais, M.R., Leite, L.C.C., Lemos, E.G.M., Lemos, M.V.F., Lopes, S.A., Lopes, C.R., Machado, J.A., Machado, M.A., Madeira, A.M.B.N., Madeira, H.M.F., Marino, C.L., Marques, M.V., Martins, E.A.L., Martins, E.M.F., Matsukuma, A.Y., Menck, C.F.M., Miracca, E.C., Miyaki, C.Y., Monteiro-Vitorello, C.B., Moon, D.H., Nagai, M.A., Nascimento, A.L.T.O., Netto, L.E.S., Nhani Jr, A., Nobrega, F.G., Nunes, L.R., Oliveira, M.A., Oliveira, de M.C., Oliveira, de R.C., Palmieri, D.A., Paris, A., Peixoto, B.R., Pereira, G.A.G., Pereira Jr, H.A., Pesquero, J.B., Quaggio, R.B., Roberto, P.G., Rodrigues, V., M.A.J., Rosa, de, Rosa de Jr, V.e., Sa de, R.G., Santelli, R.V., Sawasaki, H.E., Silva da, A.C.R., Silva da, A.M., Silva da, F.R., Silva Jr, W.A., Silveira da, J.F., Silvestri, M.L.Z., Siqueira, W.J., Souza de, A.A., Souza de, A.P., Terenzi, M.F., Truffi, D., Tsai, S.M., Tuhako, M.H., Vallada, H., Van Sluys, M.A., Verjovski-Almeida, S., Vettore, A.L., Zago, M.A., Zatz, M., Meidanis, J. and Setubal, J.C. (2000). The genome sequence of the plant pathogen *Xylella fastidiosa*. *Nature* **406**, 151-157.
- Starr, M.P. (1981). The genus *Xanthomonas*. In *The Prokaryotes Vol 1*, M.P. Starr, H. Stolp, H.G. Truper, A. Balows, and H.G. Schlegel, eds (Springer Verlag, Berlin), pp 742-763.
- Staskawicz, B., Mudgett, M.B., Dangl, J. and Galan, J.E. (2001). Common and contrasting themes of plant and animal diseases. *Science* **292**, 2285-2289.
- Stenmark, H., and Oikkonen, V. M. (2001) The Rab GTPase family. *Genome Biology* **2**, 1-7.
- Swings, J., Vauterin, L. and Kersters, K. (1993). The bacterium *Xanthomonas*. In *Xanthomonas*, J.G. Swings and E.L. Civerolo, eds (Chapman and Hall, London) pp. 121-156
- Tsuji, J. and Somerville, S.C. (1988). *Xanthomonas campestris* pv. *campestris* induces chlorosis in *Arabidopsis thaliana*. *Arabidopsis Information Service* **26**, 1-8.
- Tsuji, J., Somerville, S.C. and Hammerschmidt, R. (1991). Identification of a gene in *Arabidopsis thaliana* that controls resistance to *Xanthomonas campestris* pv. *campestris*. *Physiol. Mol. Plant Pathology* **38**, 57-65.
- Tsuji, J. and Somerville, S.C. (1992). First report of the natural infection of *Arabidopsis thaliana* by *Xanthomonas campestris* pv. *campestris*. *Plant Disease* **76**, 539.
- Tsuji, J., Jackson, E.P., Gage, D.A., Hammerschmidt, R., and Somerville, S.C. (1992) Phytoalexin accumulation in *Arabidopsis thaliana* during the hypersensitive reaction to *Pseudomonas syringae* pv *syringae*. *Plant Physiol.* **98**, 1304-1309.
- Van den Mooter, M., and Swings, J (1990). Numerical analysis of 295 phenotypic features of 266 *Xanthomonas* strains and related strains and an improved taxonomy of the genus. *Int. J. of Systematic Bacteriology* **40**, 348-369.
- Williams, P.H. (1980). Black Rot: A continuing threat to world crucifers. *Plant Disease* **64**, 736-742.
- Yang, Y. and Klessig, D. F. (1996). Isolation and characterization of a tobacco mosaic virus-inducible *myb* oncogen homolog from tobacco. *Proc. Natl. Acad. Sci. USA* **93**, 1472-14977.