

## Correction

Authors: Wyatt, Sarah E., and Sanders, Heather L.

Source: BioScience, 59(10): 821

Published By: American Institute of Biological Sciences

URL: https://doi.org/10.1525/bio.2009.59.10.21

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.


## Correction

We have discovered that we made an error in our article "Leaf Evolution and Development: Advancing Technologies, Advancing Understanding" (*BioScience* 59: 17–26). In figure 4, we show a sample of the range of leaf morphologies represented in the pea mutant collection. Unfortunately, we inaccurately identified the mutants in some of the panels. The figure legend should read as follows: "Figure 4. A sample of the range of leaf morphologies represented in the pea mutant collection. (a) The *nana1* mutant has miniaturized leaves and fewer pairs of leaflets and tendrils, (b) the *afila/crispa* double mutant exhibits a range of abnormal morphologies, and (c) the *unifoliata* mutant and the *uni/afila* double mutant show terminal leaflets and truncated development. The *uni* mutants alone show reduced levels of dissection, while the double mutants reverse that slightly; (d) the upper two leaves are from wild-type plants and the lower are *uni-tac*, a less severe mutation at the *uni* locus. Images provided by Darleen DeMason, University of California, Riverside. Scale bar = 1 centimeter in (a) and (c), and 2 centimeters in (b) and (d)."

We sincerely regret the mistake.

SARAH E. WYATT HEATHER L. SANDERS Sarah E. Wyatt (wyatts@ohio.edu) is an associate professor at Ohio University in Athens, Ohio. Heather L. Sanders is with the Department of Plant Sciences at Oxford University, United Kingdom.

doi:10.1525/bio.2009.59.10.21



www.biosciencemag.org