

## **Worldwide Spread of *Pheidole teneriffana* (Hymenoptera: Formicidae)**

Author: Wetterer, James K.

Source: Florida Entomologist, 94(4) : 843-847

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.094.0417>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## WORLDWIDE SPREAD OF *PHEIDOLE TENERIFFANA* (HYMENOPTERA: FORMICIDAE)

JAMES K. WETTERER

Wilkes Honors College, Florida Atlantic University, 5353 Parkside Drive, Jupiter, FL 33458

### ABSTRACT

*Pheidole teneriffana* is an Old World ant species that has spread to other parts of the world through human commerce. To evaluate its known distribution, I compiled and mapped *P. teneriffana* records from >200 sites. I documented the earliest known *P. teneriffana* records for 43 geographic areas (countries, island groups, major islands, and US states), including 11 West Indian islands for which I found no previously published records: Antigua, Barbados, Barbuda, Curaçao, Dominica, Dominican Republic (Hispanola), Guadeloupe, Jamaica, Martinique, Montserrat, and St Lucia. *Pheidole teneriffana* is known from sites scattered across the greater Mediterranean region (North Africa, southern Europe, the Middle East, and neighboring islands), though it is unclear which parts of this area constitute its native range. *Pheidole teneriffana* is known from only a few Old World sites distant from the Mediterranean region, i.e., Ascension, China, England, St Helena, and South Africa. In the New World, where this species is certainly exotic, there are published *P. teneriffana* records only from California, Cuba, and Peru. The first records from 11 West Indian islands presented here, all from 2003 or later, suggest that *P. teneriffana* is actively spreading through this region. Almost all West Indian records of *P. teneriffana* come from beaches or urban areas. It seems doubtful that *P. teneriffana* will develop into a major pest species like its congener, the infamous *Pheidole megacephala*.

Key words: biogeography, biological invasion, exotic species, invasive species

### RESUMEN

*Pheidole teneriffana* es una especie de hormigas del Viejo Mundo que se ha extendido a otras partes del mundo a través del comercio humano. Para evaluar su distribución conocida, he compilado registros de *Pheidole teneriffana* de >200 sitios y los he puesto en un mapa. Documenté los registros de *P. teneriffana* mas antiguos conocidos de 43 zonas geográficas (países, grupos de islas, islas principales, y los estados de los EE.UU), incluyendo 11 islas de las Antillas de la cuales se encontró registros no estaban publicados anteriormente: Antigua, Barbados, Barbuda, Curazao, Dominica, República Dominicana (Hispanola), Guadalupe, Jamaica, Martinica, Montserrat y Santa Lucia. *Pheidole teneriffana* es conocida de sitios dispersos por toda la región Mediterránea (norte de África, el sur de Europa, el Medio Oriente y las islas vecinas), aunque no está claro qué partes de esta zona constituyen su área de distribución nativa. Se conoce la hormiga *Pheidole teneriffana* solamente de unos pocos lugares del Viejo Mundo lejos de la región Mediterránea, es decir, Ascensión, China, Inglaterra, Santa Elena y Sudáfrica. En el Nuevo Mundo, donde esta especie es ciertamente exótica, solo hay registros de *Pheidole teneriffana* publicados de California, Cuba y Perú. Los primeros registros de 11 islas de las Antillas que se presentan aqui, todos del año 2003 o después, indican que *P. teneriffana* esta dispersando activamente por esta región. Casi todos los registros de *P. teneriffana* de las Antillas provienen de las playas o zonas urbanas. Es ciertamente dudoso que *P. teneriffana* se convertirá en una importante especie plaga como su congénere, la infame *Pheidole megacephala*.

Numerous ant species have spread around the world through human commerce. For the most part, these “tramp” ants thrive only in disturbed environments and do not penetrate intact natural habitats. But as humans and their disturbance spread, so do tramp ants. The ecological importance of most tramp ants remains undocumented. Several species, however, such as *Pheidole megacephala* F., are known to have dramatic impacts (Wetterer 2007). For example, Zimmerman (1970) wrote that in Hawaii, “the endemic insect faunas of the lowlands of all the islands mostly have been

exterminated throughout the range of the voracious introduced predatory ant *Pheidole megacephala*.” In Australia, Young (2000) reported: “the rainforest at Howard Springs Nature Park is dominated by the coastal brown ant [*P. megacephala*], which has eliminated almost all species of native ants, other insect species, snails, spiders and centipedes.”

In addition to *P. megacephala*, only one other Old World *Pheidole* species has spread to the New World: *Pheidole teneriffana* Forel, a species similar in size and general appearance as *P. mega-*

*cephala*. Here, I compiled and mapped of *P. teneriffana* specimen records to evaluate its known worldwide distribution and speculate on its native range and possible impact.

Forel (1893) described *P. teneriffana* from Tenerife in the Canary Islands, off the northwest coast of Africa. *Pheidole teneriffana* is the senior synonym of *P. teneriffana taina* described from Cuba. Taylor (2010) speculated on-line that *P. teneriffana* is a junior synonym of *P. fervens*, though this view has not been accepted by other researchers (S. Cover, pers. comm.). Like most *Pheidole* species, *P. teneriffana* shows complete dimorphism (i.e., distinct minor and major workers with few, if any, intermediates) with majors having disproportionately large heads compared to minors. It is fairly simple to distinguish *P. teneriffana* from *P. megacephala*. In *P. teneriffana* majors, sculpturing covers the entire dorsal surface of the head, but in *P. megacephala*, sculpturing is confined to the anterior half of the head, (S. Cover, pers. comm.).

#### MATERIALS AND METHODS

Using published and unpublished records, I documented the worldwide range of *Pheidole teneriffana*. I obtained unpublished site records from museum specimens in the collection of the Museum of Comparative Zoology (MCZ) and the Smithsonian Institution (SI). In addition, I used on-line databases with collection information on specimens by Antweb ([www.antweb.org](http://www.antweb.org)), Asociación Ibérica de Mirmecología ([www.formicidae.org](http://www.formicidae.org)), the Global Biodiversity Information Facility ([www.gbif.org](http://www.gbif.org)), and Ants of Africa (Taylor 2010). Finally, I collected *P. teneriffana* specimens on islands of the West Indies.

I obtained geo-coordinates for collection sites from published references, specimen labels, maps, or geography web sites (e.g., [earth.google.com](http://earth.google.com), [www.tageo.com](http://www.tageo.com), and [www.fall-ingrain.com](http://www.fall-ingrain.com)). If a site record listed a geographic region rather than a "point locale," and I had no other record for this region, I used the coordinates of the largest town within the region or, in the case of small islands and natural areas, the center of the region. I did not map records of *P. teneriffana* found in newly imported goods or intercepted in transit by quarantine inspectors, e.g., a specimen intercepted in South Africa on goods from Kenya (G. Arnold; SI).

#### RESULTS

I collected *Pheidole teneriffana* at 20 sites on 10 West Indian islands: Antigua: St John's, city street (May 2007); Barbados: Bridgetown, city street (Jun 2006) and Oistins Beach (Nov 2003); Barbuda: Codrington, town (Jul 2007); Curaçao: Otrobanda, city street (Aug 2011), Punda, city

street (Jul 2004), and Punda, market (Aug 2011); Dominica: Roseau, urban waterfront (Jun 2004); Guadeloupe: Bas du Fort, shopping center (May 2008), Pointe-à-Pitre, by downtown store (Jun 2011), Sainte Anne, town center (May 2008), and Vieux Habitants, town (Jun 2008); Jamaica: Montego Bay, urban produce market (Dec 2010); Martinique: Fort-de-France, urban park (May 2008), Fort-de-France, city street (May 2008), and Robert, urban waterfront (Jul 2011); Montserrat: Old Road Bay, waterfront by an abandoned hotel (Jul 2007); St Lucia: Dennery, waterfront (Jul 2006), Micoud, waterfront (Jul 2006), and Pigeon Island, beach (Nov 2003). In addition, I obtained unpublished records from two other West Indian sites: Dominican Republic: Navarrette at bakery (Dec 2003; S. Cover) and St Lucia: Soufriere (Jan 2005; J. Endeman). I deposited voucher specimens from all sites at the MCZ.

I compiled *Pheidole teneriffana* specimen records from >200 sites worldwide (Fig. 1). I documented the earliest known *P. teneriffana* records for 43 geographic areas (countries, island groups, major islands, and US states; Tables 1-3) including 11 West Indian islands for which I found no previously published records: Antigua, Barbados, Barbuda, Curaçao, Dominica, Dominican Republic (Hispaniola), Guadeloupe, Jamaica, Martinique, Montserrat, and St Lucia.

Santschi (1919) listed *Pheidole teneriffana* (identified by G. Arnold), but not *P. megacephala*, from Samoa. Santschi (1928), however, listed *P. megacephala* from Samoa, but not *P. teneriffana*, indicating that he dismissed the earlier identification. Almost half of all mapped site records I found for *P. teneriffana* came from Egypt (e.g., Santschi 1908, Forel 1913, Alfieri 1931). Although Taylor (2010) regarded the records of *P. teneriffana* from Egypt to be *Pheidole providens*, a species otherwise known only from India, I mapped these as *P. teneriffana* nonetheless.

#### DISCUSSION

*Pheidole teneriffana* has widely scattered records from across the greater Mediterranean region (North Africa, Southern Europe, the Middle East, and neighboring islands), though it is unclear which parts of this region, if any, constitute its native range. Santschi (1918) proposed that *P. teneriffana* probably originated in upper Nile area, in what is now South Sudan, but there are no *P. teneriffana* specimens known from this region. Wilson (2003) wrote that *P. teneriffana* "is evidently a native of North Africa and possibly also the Canary Islands." Emery (1915) and Espadaler & Bernal (2003) considered *P. teneriffana* as exotic in the Canary Islands. Bytinski-Salz (1953) considered *P. teneriffana* as a recently arrived exotic in Israel, writing, "*Pheidole tenerif-*




Fig. 1. Worldwide distribution records of *Pheidole teneriffana*. All New World records are previously unpublished, except those from California, Cuba, and Peru.

*fana*, so far found east only till Egypt, reached Tel Aviv a few years ago.” Collingwood et al. (2004) wrote that *P. teneriffana* “is probably native throughout northern Africa and appears to be spreading over a wide front in the Middle East, Arabia and the Mediterranean countries.” If *P. teneriffana* is truly native across North Africa, it is remarkable how few records I found from any North African country other than Egypt: Western Sahara (0 records), Morocco (1 record), Algeria (0

TABLE 1. EARLIEST KNOWN RECORDS FOR *PHEIDOLE TENERIFFANA* FROM THE GREATER MEDITERRANEAN REGION: NORTH AFRICA, THE MIDDLE EAST, SOUTHERN EUROPE, AND NEIGHBORING ISLANDS.

|                  | Earliest record                                    |
|------------------|----------------------------------------------------|
| Canary Islands   | 1893 (Forel 1893)                                  |
| Eritrea          | ≤1901 (Emery 1901)                                 |
| Egypt            | ≤1908 (Santschi 1908)                              |
| Tunisia          | 1908 (Santschi 1908)                               |
| Turkey           | ≤1911 (Forel 1911)                                 |
| Sudan            | ≤1918 (Santschi 1918)                              |
| Libya            | ≤1924 (Emery 1924)                                 |
| Greece           | 1932 (Zimmermann 1934)                             |
| Syria            | ≤1934 (Menozzi 1934)                               |
| Israel           | ≤1949 (Bytinski-Salz 1953)                         |
| Saudi Arabia     | 1950 (Collingwood 1985)                            |
| Malta            | 1965 (Baroni Urbani 1968)                          |
| Spain            | 1981 (Acosta & Martínez 1983)                      |
| Balearic Islands | 1985 (de Haro et al. 1986)                         |
| Lebanon          | ≤1987 (Kugler 1988)                                |
| Italy            | 1987 (Mei 1995)                                    |
| Sicily           | 1990 (Mei 1995)                                    |
| UAE              | 1991 (Collingwood et al. 1997)                     |
| Morocco          | 1992 (Delabie 2007)                                |
| Yemen            | 1993 (Collingwood & Agosti 1996)                   |
| Iran             | ≤1995 (Alipanah et al. 1995 in Paknia et al. 2008) |
| Oman             | ≤1996 (Collingwood & Agosti 1996)                  |
| Kuwait           | ≤1996 (Collingwood & Agosti 1996)                  |

TABLE 2. EARLIEST KNOWN RECORDS FOR *PHEIDOLE TENERIFFANA* FROM SUB-SAHARAN AFRICA, EAST ASIA, NORTH EUROPE, AND SOUTH ATLANTIC ISLANDS.

|              | Earliest record             |
|--------------|-----------------------------|
| Kenya        | ≤1918 (Santschi 1918)       |
| South Africa | ≤1925 (Santschi 1925)       |
| China        | ≤1925 (Santschi 1925)       |
| England      | ≤1987 (Collingwood 1987)    |
| St Helena    | 2002 (Wetterer et al. 2007) |
| Ascension    | 2002 (Wetterer et al. 2007) |

records), Tunisia (2 records), Libya (1 record), and Sudan (1 record).

The earliest records of *P. teneriffana* come from the Canary Islands in the far west of the greater Mediterranean region and from Eritrea in the far southeast, followed by records from Egypt, Tunisia, and Turkey in between (Table 1). This chronology of the earliest known records does not favor any particular hypothesis concerning what part of the greater Mediterranean region may be the original native range of *P. teneriffana*. Suppositions on where *P. teneriffana* originated have been based on very little evidence. Genetic analyses may be necessary to evaluate the native range of this species. It seems unlikely that *P. teneriffana* is native to any of the isolated Old World sites where it has been collected distant from the Mediterranean region, i.e., Ascension, China, England, St Helena, and South Africa.

In the New World, where this species is certainly exotic, I found published *P. teneriffana* records only from California, Cuba, and Peru (Wilson 2003). The records presented here of *P. tenerif-*

*fana* from 11 West Indian islands, all from 2003 or later, suggest that *P. teneriffana* is actively spreading through this region. In the West Indies, I found *P. teneriffana* almost exclusively on beaches and at highly disturbed urban sites, particularly in waterfront areas. Santschi (1934), reporting this species from Alexandria, Egypt, similarly noted that *P. teneriffana* was rarely reported far from seaports. Both in Montego Bay, Jamaica and Pointe-à-Pitre, Guadeloupe, I found *P. teneriffana* nesting in the cracked concrete at the base of a doorway of a market. In Montego Bay, I found no other ants in the nearby dusty urban street, while in Pointe-à-Pitre, I found *P. megacephala* nesting at the bases of all the neighboring trees along the sidewalk outside the store. It may be that *P. teneriffana* is able to nest in microhabitats too dry for other species. Curiously, most Old World records of *P. teneriffana* are subtropical, but all New World records are tropical, except one from California (Fig. 1), and this one population appears to have been extirpated by another exotic ant, *Linepithema humile* (Gulmahamad 1999).

*Pheidole teneriffana* has remained relatively inconspicuous throughout most of its range, though it may become common in some urban areas. Collingwood et al. (1997) reported that in the United Arab Emirates, *P. teneriffana* was “very populous in irrigated gardens and along the coast where it appears to be spreading rapidly, and might replace local species.” In the Balearic Islands, Gómez & Espadaler (2006) reported that in Ibiza city, Ibiza, *P. teneriffana* “has become a common species in the gardens and trees and on sidewalks near the harbour. It seems to be expanding its range in Mallorca from El Arenal gardens - where it may become very abundant—to Son Veri to the East and La Lonja (Palma city) in the West,

TABLE 3. EARLIEST KNOWN RECORDS FOR *PHEIDOLE TENERIFFANA* FROM THE NEW WORLD. UNPUBLISHED RECORDS INCLUDE COLLECTOR, MUSEUM SOURCE, AND SITE.

|                 | Earliest record                                    |
|-----------------|----------------------------------------------------|
| Cuba            | 1930 (Aguayo 1932 as <i>P. teneriffana taina</i> ) |
| Peru            | 1967 (R. H. Crozier, MCZ): Ciudad Univ. San Marcos |
| California      | 1989 (Martinez 1992)                               |
| +St Lucia       | 2003 (J. K. Wetterer, MCZ): Pigeon Island          |
| +Barbados       | 2003 (J. K. Wetterer, MCZ): Oistins                |
| +Dominican Rep. | 2003 (S. P. Cover, MCZ): Navarrette                |
| +Dominica       | 2004 (J. K. Wetterer, MCZ): Roseau                 |
| +Curaçao        | 2004 (J. K. Wetterer, MCZ): Punda                  |
| +Antigua        | 2007 (J. K. Wetterer, MCZ): St John’s              |
| +Barbuda        | 2007 (J. K. Wetterer, MCZ): Codrington             |
| +Montserrat     | 2007 (J. K. Wetterer, MCZ): Old Road Bay           |
| +Martinique     | 2008 (J. K. Wetterer, MCZ): Fort-de-France         |
| +Guadeloupe     | 2008 (J. K. Wetterer, MCZ): Sainte Anne            |
| +Jamaica        | 2010 (J. K. Wetterer, MCZ): Montego Bay            |

MCZ = Museum of Comparative Zoology. + = No previously published records.



always in disturbed habitats.” Nonetheless, it seems unlikely that *P. teneriffana* will develop into a major pest species like *P. megacephala*. Future studies might examine why *P. teneriffana* and *P. megacephala*, two species that are morphologically very similar, have such different ecological impacts.

#### ACKNOWLEDGMENTS

I thank M. Wetterer for comments on this manuscript; S. Cover and L. Davis for ant identification; S. Cover and J. Endeman for providing unpublished records; S. Cover (MCZ) and T. Schultz (SI) for help with their respective ant collections; W. O'Brien for GIS help; D. P. Wojcik and S. D. Porter for compiling their valuable FORMIS bibliography; R. Pasos and W. Howerton of the FAU library for processing so many interlibrary loans; Florida Atlantic University and the National Science Foundation (DES-0515648) for financial support.

#### REFERENCES CITED

- ACOSTA, F. J., AND MARTÍNEZ, M. D. 1983. *Pheidole teneriffana* Forel, 1893. (Hym. Formicidae), nueva cita para la Península Ibérica. Bol. Asoc. Española Entomol. 7: 320.
- AGUAYO, C. G. 1932. Notes on the West Indian ants. Bull. Brooklyn Entomol. Soc. 27: 215-227.
- BARONI URBANI, C. 1968. Studi sulla mirmecofauna d'Italia. IV. La fauna mirmecologica delle isole Maltesi ed il suo significato ecologico e biogeografico. Ann. Mus. Civ. Sto. Natur. "Giacomo Doria" 77: 408-559.
- BYTINSKI-SALZ, H. 1953. The zoogeography of the ants in the Near East. Istanbul Univ. Fen Fakult. Mecmuasi. Seri B 19: 67-74.
- COLLINGWOOD, C. A. 1985. Hymenoptera: Fam. Formicidae of Saudi Arabia. Fauna Saudi Arabia 7: 230-302.
- COLLINGWOOD, C. A. 1987. Ant species in Yorkshire. Naturalist 112: 95-101.
- COLLINGWOOD, C. A., AND AGOSTI, D. 1996. Formicidae (Insects: Hymenoptera) of Saudi Arabia (Part 2). Fauna Saudi Arabia 15: 300-385.
- COLLINGWOOD, C. A., POHL, H., GÜSTEN, R., WRANIK, W., AND VAN HARTEN, A. 2004. The ants (Insecta: Hymenoptera: Formicidae) of the Socotra Archipelago. Fauna Arabia 20: 473-495.
- COLLINGWOOD, C. A., TIGAR, B. J., AND AGOSTI, D. 1997. Introduced ants in the United Arab Emirates. J. Arid Environ. 37: 505-512.
- de HARO, A., COLLINGWOOD, C. A., AND COMÍN, P. 1986. Prospección mirmecológica por Ibiza y Formentera (Balears). Orsis 2: 115-120.
- DELABIE, J. 2007. Présence de *Pheidole teneriffana* Forel, 1893, au Maroc (Hym., Formicidae, Myrmicinae). Bull. Soc. Entomol. France 112: 288.
- EMERY, C. 1901. Spicilegio mirmecologico. Boll. Soc. Entomol. Italiana 33: 57-63.
- EMERY, C. 1915. Les *Pheidole* du groupe *megacephala* (Formicidae). Rev. Zool. Africaine 4: 223-250.
- EMERY, C. 1924. Formiche della Cirenaica raccolte dal Dott. Enrico Festa e dal Prof. Filippo Silvestri. Boll. Soc. Entomol. Italiana 56: 6-11.
- ESPADALER, X., AND BERNAL, V. 2003. Exotic ants in the Canary Islands, Spain (Hymenoptera, Formicidae). Vieraea 31: 1-7.
- FOREL, A. 1893. Nouvelles fourmis d'Australie et des Canaries. Ann. Soc. Entomol. Belgique 37: 454-466.
- FOREL, A. 1911. Fourmis nouvelles ou intéressantes. Bull. Soc. Vaud. Sci. Natur. 47: 331-400.
- GÓMEZ, K., AND ESPADALER, X. 2006. Exotic ants (Hymenoptera: Formicidae) in the Balearic Islands. Myrmecol. Nachr. 8: 225-233.
- GULMAHAMAD, H. 1999. Extirpation of one exotic ant species by another in southern California. Pan-Pacific Entomol. 75: 227-229.
- KUGLER, J. 1988. The zoogeography of Israel. 9. The zoogeography of social insects of Israel and Sinai. Monog. Biolog. 62: 251-275.
- MARTÍNEZ, M. J. 1992. A new ant introduction for North America: *Pheidole teneriffana* (Forel) (Hymenoptera: Formicidae). Pan-Pacific Entomol. 68: 153-154.
- MEI, M. 1995. Arthropoda di Lampedusa, Linosa e Pantelleria (Canale di Sicilia, Mar Mediterraneo). Hymenoptera Formicidae (con diagnosi di due nuove specie). Il Natur. Siciliano 4: 753-772.
- MENOZZI, C. 1934. Reperti mirmecofaunistici raccolte dal Prof. L. di Caporriaco nelle oasi di Cufra e in altre località del deserto Libico. Att. Soc. Natur. Matemat. Modena 65[= (6)13]: 153-166.
- PAKNIA, O., RADCHENKO, A., ALIPANAH, H., AND PFEIFFER, M. 2008. A preliminary checklist of the ants (Hymenoptera: Formicidae) of Iran. Myrmecol. Nachr. 11: 151-159.
- SANTSCHI, F. 1908. Nouvelles fourmis de l'Afrique du Nord (Égypte, Canaries, Tunisie). Ann. Soc. Entomol. France 77: 517-534.
- SANTSCHI, F. 1918. Cinq notes myrmécologiques. Bull. Soc. Vaud. Sci. Natur. 52: 63-64.
- SANTSCHI, F. 1919. Cinq notes myrmécologiques. Bull. Soc. Vaud. Sci. Natur. 52: 325-350.
- SANTSCHI, F. 1925. Contribution à la faune myrmécologique de la Chine. Bull. Soc. Vaud. Sci. Natur. 56: 81-96.
- SANTSCHI, F. 1928. Formicidae (Formis). Insects Samoa 5:41-58.
- SANTSCHI, F. 1934. Fourmis d'une croisière. Bull. Ann. Soc. Entomol. Belgique 74: 273-282.
- TAYLOR, B. 2010. Ants of Africa. Tenth Edition. <<http://antbase.org/ants/africa>>, retrieved on 13 December 2010.
- WETTERER, J. K. 2007. Biology and impacts of Pacific islands invasive species. 3. The African big-headed ant, *Pheidole megacephala* (Hymenoptera: Formicidae). Pacific Sci. 61: 437-456.
- WETTERER, J. K., ESPADALER, X., ASHMOLE, P., CUTLER, C., AND ENDEMAN, J. 2007. Ants (Hymenoptera: Formicidae) of the South Atlantic islands of Ascension Island, St Helena, and Tristan da Cunha. Myrmecol. News 10: 29-37.
- WILSON, E. O. 2003. *Pheidole* on the New World: A dominant, hyperdiverse ant genus. Harvard University Press, Cambridge, Massachusetts. 794 pp.
- YOUNG, G. 2000. The coastal brown or big headed ant. Agnote 152: 1-4.
- ZIMMERMAN, E. C. 1970. Adaptive radiation in Hawaii with special reference to insects. Biotrop. 2: 32-38.
- ZIMMERMANN, S. 1934. Liste der von Prof. Dr. Fr. Werner im der Jahre 1932 auf den Ägäischen Inseln gesammelten Formiciden. Sitzungsber. Akad. Wiss. Wien Math. Naturwiss. Kl. Abt. I 1934: 66-69.