

## **New Data on *Conorhynchus hololeucus* (Coleoptera: Curculionidae)**

Author: Gültekin, Levent

Source: Florida Entomologist, 95(2) : 503-505

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.095.0241>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## NEW DATA ON *CONORHYNCHUS HOLOLEUCUS* (COLEOPTERA: CURCULIONIDAE)

LEVENT GÜLTEKİN

Atatürk University, Faculty of Agriculture, Department of Plant Protection, 25240 Erzurum, Turkey

E-mail: lgultekin@gmail.com

Trehala are unusual insect pupal chambers that are sweet and have been used as human food (Pierce 1915). Trehala are constructed on the outside of stems by larvae in the genus *Larinus* Dejean, 1821 (Coleoptera: Curculionidae: Lixinae). New data on 4 trehala constructing species of *Larinus* were recently presented by Gültekin (2008). Trehala-like capsule constructions have also been observed in several genera of Cleonini, including *Conorhynchus* Motschulsky, 1860, *Trichocleonus* Motschulsky, 1860, *Scaphomorphus* Motschulsky, 1860 and *Pachycerus* Schoenherr, 1823 (Volovnik 2010), and the Lixini genus *Hypolixus* Desbrochers, 1898 (Nikulina 1989). Larvae, pupae and new generation adults of *Hypolixus tigrinus* (Reitter 1890) were found in oval trehala-like capsules on the stem base of *Suaeda arcuata* Bunge (Caryophyllales: Amaranthaceae) and *S. paradoxa* Bunge (Nikulina 1989).

A recent ecological research project conducted in eastern Turkey revealed that larvae of *Conorhynchus hololeucus* (Pallas, 1781) (Coleoptera: Curculionidae: Lixinae) create structures similar to the trehala of *Larinus*. Their host plant in eastern Turkey is *Atriplex* sp. (Caryophyllales: Amaranthaceae) (Fig. 1), and according to Ter-Minassian (1988) host plants of *C. hololeucus* are *Atriplex tarica* L., *Salsola sclerantha* C. A. Meyer and *S. carinata* C. A. Meyer (Caryophyllales: Amaranthaceae) in former USSR. Adults feed on the leaves by chewing out semicircular areas. Usually 2-3 pairs of adults are present on one plant and they mate on the leaves (Fig. 2). The females lay eggs in the neck of the root of the host plant. The young larvae open superficial galleries downwards under the bark of the root. As each larva develops it makes an external trehala-like capsule, which it enlarges as development proceeds. The legless larva has a creamy white color with a brown head capsule (Fig. 3). The capsule may be separate (Fig. 4), or 4-5 capsules may be connected to each other on a single root of a host plant (Fig. 6). Pupation occurs in the capsule (Fig. 5) in late summer. The new generation of adults remains for some d in their capsules, and emerge in late summer-early autumn of the same year. Upon emergence, the adults search out a secluded place on the ground for hibernation. This weevil species produces one generation per year.

### Materials Examined

**TURKEY:** Erzurum Prov., Atatürk University Campus [N 39° 54'.266", E 41° 12'.962"] 1850m

asl, 28-VIII-2005, L. Gültekin, 1 ex.; Iğdır Prov., 5 km E of Karakoyunlu, 850 m asl, 2-VI-2002, L. Gültekin, 4 ex.; Bulakbaşı Vill. [N 39° 57'.072", E 44° 15'.299"], 848 m asl, 9-VII-2005, L. Gültekin, 11 ex.; 15 km E of Aralık, 880 m asl, 11-VI-2003, L. Gültekin, 13 ex.; Kars Prov., 40 km W of Kağızman, 1400 m asl, 1-VI-2002, L. Gültekin, 1 ex.; Konya Prov., 70 km SE of Aksaray [N 37° 46'.355", E 34° 14'.230"], 1043 m asl, 27-VI-2005, L. Gültekin, 1 ex.

### Distribution

The distribution of this weevil species includes Azerbaijan, Armenia, Russia, Iran, Kazakhstan, Mongolia (Csiki 1934; Ter-Minassian 1988) and new record for Turkey.

### Taxonomic Remarks

Recently, Alonso-Zarazaga & Lyal (2009) downgraded *Temnorhinus* Chevrolat, 1872 to a subgenus of *Conorhynchus*. According to this taxonomical act, *Conorhynchus* (*Temnorhinus*) *hololeucus* (Pallas, 1781) is a member of this genus and well suited.


The study was supported by the Collaborative Linkage Grant No. 978845 and NR-CLG-981318 of the NATO Life Science and Technology Programme.

### SUMMARY

Larvae of *Conorhynchus hololeucus* (Coleoptera: Curculionidae: Lixinae) construct a trehala-like capsule for the immature stages. The host plant is *Atriplex* sp. (Caryophyllales: Amaranthaceae), and this weevil species is a new record for Turkey. Adults feed on leaves; females lay eggs in root neck of host plants; larvae make an external trehala-like capsule on the root; pupation and adult emergence occur in the capsule. This weevil produces one generation per year.

### REFERENCES CITED

- ALONSO-ZARAZAGA, M. A., AND LYAL, C. H. C. 2009. On the nomenclature of some genera in the Cleonini (Coleoptera, Curculionidae, Lixinae). *Graellsia* 65(2): 241-242.
- CSIKI, E. 1934. *Coleopterorum Catalogus auspiciis et auxilio W. Junk editus a S. Schenking. Pars 134. Curculionidae: Subfam. Cleoninae.* Junk, Berlin, 152 pp.



Figs. 1-6. Host plant and biological stages of *Conorhynchus hololeucus* (Pallas). (1) *Atriplex* sp. with trehala-like capsule of *C. hololeucus* on the root; (2) feeding damage and mating pair of *C. hololeucus* on host plant; (3) larva; (4) individual capsule; (5) pupa in capsule; (6) multiple trehala-like capsules.

- GÜLTEKIN, L. 2008. Taxonomic review of the stem-inhabiting trehala-constructing *Larinus* Dejean, 1821 (Coleoptera: Curculionidae): New species, systematics and ecology. *Zootaxa* 1714: 1-18.
- NIKULINA, O. N. 1989. Contributions to the biology of weevils of the genus *Lixus* F. (Coleoptera, Curculionidae), developing in semishrubs and grasses of Tajikistan. *Entomol. Obozrenie* 68(3): 511-521.
- PIERCE, W. D. 1915. The uses of certain weevils and weevil products in food and medicine. *Proc. Entomol. Soc. Washington* 27: 151-154.
- TER-MINASSIAN, M. E. 1988. Zūki-dolgonosiki podsemejstva Cleoninae Fauny SSSR. Kornevye dolgonosiki (triba Cleonini). In Skarlat, O. A. (ed.), *Opredeliteli po Faune SSSR, izdavaemye Zoologiceskim Institutom Akademii Nauk SSSR* 155: 1-235.
- VOLOVNIK, S. V. 2010. Weevils Lixinae (Coleoptera, Curculionidae) as gall formers. *Entomol. Rev.* 90(5): 585-590.