

Hirsutella sp. (Hypocreales: Ophiocordycipitaceae) Affecting the Invasive Social Wasp *Vespula vulgaris* (Hymenoptera: Vespidae) in Southern Chile

Authors: Montalva, Cristian, Rojas, Eladio, Valenzuela, Eduardo, and Humber, Richard A.

Source: Florida Entomologist, 100(3) : 663-666

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0327>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Hirsutella sp. (Hypocreales: Ophiocordycipitaceae) affecting the invasive social wasp *Vespula vulgaris* (Hymenoptera: Vespidae) in southern Chile

Cristian Montalva^{1,*}, Eladio Rojas², Eduardo Valenzuela¹, and Richard A. Humber³

The genus *Vespula* (Hymenoptera: Vespidae) includes wasp species that have painful stings that cause irritating nuisances with impacts on human health and on a range of outdoor activities. These wasps are economically significant pests of such primary industries as beekeeping, forestry, and horticulture (Dymock et al. 1994; Beggs 2000, 2001; Gardner-Gee & Beggs 2012). The stings of the common wasp, *Vespula vulgaris* (L.) (Hymenoptera: Vespidae), are well documented to induce allergic reactions and, occasionally, fatal anaphylaxis (King et al. 1996; King & Spangfort 2000; Fitch et al. 2001; Tankersley & Ledford 2015). This wasp, which is native to Eurasia, has become a notorious pest in Argentina and New Zealand, where it can attain high population densities and cause major ecological consequences such as increased predation pressure on native insect communities (Yamane et al. 1980; Beggs 2001; Baz et al. 2010; Lester et al. 2014).

Vespula vulgaris was first detected in Chile during the summer of 2011, in the mountains of the Araucanía region (Barrera-Medina & Vidal 2013). It is now distributed from the Araucanía region through the southernmost Magallanes region. As a result of a continuous monitoring of Servicio Agrícola y Ganadero (SAG) in Los Lagos region (41.4717°S, 72.9367°W) for early detection of quarantine bark beetles in Chile, a single cadaver of a naturally fungal infected worker adult of *V. vulgaris* was collected during May 2015 (autumn in Chile). This insect was recovered in a funnel trap placed in the southern Chilean city of Puerto Montt (41.453914°S, 72.867700°W).

In the past, several natural enemies that could be useful control agents of this invasive *Vespula* species have been reported (Rose et al. 1999; Singh et al. 2010; Evison et al. 2012). Nonetheless, little is known of the biodiversity of entomopathogenic fungi on wasps in Chile apart from this report of a naturally occurring entomopathogenic fungus affecting *V. vulgaris* in southern Chile.

The identification of the fungus was done using a light microscope (Nikon® Eclipse E600; Nikon Corporation, Tokyo, Japan) and slide mounts in lactophenol-cotton blue were prepared as suggested by Humber (2012). Fifty measurements for each taxonomically significant fungal structure were made using images taken with a digital camera (Nikon® DS-Fi1; Nikon, Tokyo, Japan) and measured with Motic® Images Plus 2.0 software (Motic China Group Co., Ltd, Shenzhen, China). Attempts to cultivate the fungus involved transferring conidia to potato-dextrose-agar medium (PDA; Difco®, Becton, Dickinson and Company, Sparks, Maryland) in 60 × 15 mm Petri dishes. Dishes were sealed

with parafilm and incubated at 24 ± 1 °C and natural photophase. The development of fungi and of any contaminants was monitored daily for 15 d. However, none of the 15 attempts to cultivate this fungus on PDA were successful. This result suggested that the specimen might have suffered from some environmentally unfavorable conditions before its collection.

The dead, mycosed *V. vulgaris* adult presented a small number of relatively short, thick white synnemata whose overall appearance was exclusively typical for infections by *Hirsutella* (Hypocreales: Ophiocordycipitaceae) species (Fig. 1A, B) (Hodge 1998; Humber 2012). On the affected wasp, the fungus produced white mycelium and unbranched synnemata, 1,520 ± 340 µm long (overall range: 336–4,460 µm), emerging from the cuticle of the infected host (Fig. 1A, B). The synnemata were seen to be simple (although occasionally sparsely branched), dark colored, slender, leathery to brittle in texture, and bearing a discontinuous layer of conidiogenous cells with elongated and narrowed necks projecting from the synnemal surface. The conidiogenous cells were monophialidic, septate, scattered to moderately crowded, arose laterally from the synnemata, with cylindrical to ellipsoid bases, 25.2 ± 1.6 µm length × 1.5 ± 0.1 µm width (overall range: 16.4–44.1 µm length × 1.2–2.0 µm width) (Fig. 1C, D). The conidia were hyaline, smooth, one-celled and lemon-shaped, 8.5 ± 0.3 µm length × 3.7 ± 0.1 µm width (overall range: 7.1–10.6 µm length × 2.1–5.2 µm width), produced singly, only rarely seen in groups, and occasionally coated by an obvious mucoid sheath (Fig. 1E).

The trap containing the infected wasp was collected by Verónica Cruces (SAG-Puerto Montt), and the infected wasp was deposited in Jun 2016 in the Mycological Collection of the National Museum of Natural History of Chile, Parque Quinta Normal s/n, Santiago, Chile, as accession SGO 166649. The conidiogenous cells of this and other *Hirsutella* species are best described as phialidic, and form one to a few conidia tending strongly to be asymmetrical and often resembling the individual sections of an orange (Minter et al. 1983; Hodge 1998). The great majority of *Hirsutella* species are pathogens of insects; a few are pathogens of mites or spiders, and still fewer are pathogens of nematodes (Shimazu & Gockling 1997; Hodge KA, Plant Pathology and Plant-Microbe Biology Section, Cornell University, personal communication). The slime-embedded spores of many *Hirsutella* species are thought to be adapted for dispersal by contact with a passing invertebrate or in water drops (Evans 1989).

¹Universidad Austral de Chile, Facultad de Ciencias, Instituto de Bioquímica y Microbiología, Casilla 567, Valdivia, 5090000, Chile; E-mail: montalva.cristian@gmail.com (C. M.), evalenzu@uach.cl (E. V.)

²Servicio Agrícola y Ganadero, Laboratorio Regional, Ruta Puerto Octay U-55-V, Calle de Servicio, Osorno, 5290000, Chile; E-mail: eladio.rojas@sag.gob.cl (E. R.)

³United States Department of Agriculture (USDA) Agricultural Research Service (ARS) Emerging Pests and Pathogens Research Unit, Robert W. Holley Center for Agriculture and Health, Ithaca, NY, USA; E-mail: richard.humber@ars.usda.gov (R. A. H.)

*Corresponding author; E-mail: montalva.cristian@gmail.com (C. M.)




Fig. 1. *Vespula vulgaris* adult infected with *Hirsutella* sp. found in southern Chile. A. Mycotized wasp cadaver; B–C. synnema arising from the abdomen of the wasp; D. conidiogenous cells with cylindrical bases and an elongated, narrowed neck; E. conidiogenous cells and conidia. Horizontal scale bars represent 20 µm in C, D, and E.

The only *Hirsutella* species ever reported from the superfamily Vespoidea is *H. saussurei* (Cooke) Speare (Hypocreales: Ophiocordycitaceae). This species has been reported from the USA, Honduras, Panama, England, Papua New Guinea, Borneo, and Taiwan (Hodge 1998; Rose et al. 1999). *Hirsutella saussurei* was originally described as *Isaria saussurei* by Cooke (1892) from a *Polistes* sp. (Hymenoptera: Eumenidae) and transferred to *Hirsutella* by Speare (1920). Speare described the color of *H. saussurei* synnemata as brownish. That the synnemata of the Chilean *Hirsutella* species are whitish and smaller in size than noted by Speare (1920) suggests a reasonable possibility that these synnemata were only partially developed when the insect was collected but might have been both larger and darker if the specimen had remained in the field for a longer time.

The synnemata of the Chilean fungus differ from those of typical collections of *H. saussurei* (e.g., Speare 1920; Samson et al. 1988) by their comparatively smaller number, shorter length, comparatively greater thickness, and absence of lateral branchings of the synnemata. Speare (1920) described the conidiogenous cells of *H. saussurei* as

simple, sessile, with an inflated short basal portion tapering abruptly to a very long neck (35–70 µm), but the conidiogenous cells of the Chilean *Hirsutella* had a total length of 16.4 to 44.1 µm including the neck. As noted previously, all of these differences between the Chilean collection and other reported examples of *H. saussurei* may indicate that the fungus reported here may be a relatively immature collection of this species. Based on these results, we report the first incidence of an entomopathogenic fungus, *Hirsutella* sp. in this case, as a potential natural enemy of *V. vulgaris* in Chile. The *Hirsutella* found in the study might differ from *H. saussurei*, but more infected wasps from Chile need to be examined to determine whether the variant morphology of the fungus characterized here does, indeed, represent a developmentally early state of sporulation by *H. saussurei*.

The biodiversity of South American entomopathogenic fungi has still received surprisingly little attention. Aruta et al. (1974) and Aruta & Carrillo (1989) summarized the biodiversity of entomopathogenic fungi in Chile, and Sosa-Gomez et al. (2010) provided an extensive summary of these from Argentina and Brazil. Globally, the fungal ento-

mopathogens of *Vespula* species include *Aspergillus flavus* Link (Eurotiales: Trichocomaceae), *Beauveria bassiana* (Bals.-Criv.) Vuill. (Hypocreales: Cordycipitaceae), and *Beauveria brongniartii* (Sacc.) Petch (Cordycipitaceae) as well as a single *Hirsutella* species recorded to attack *Vespula germanica* (F.) (Hymenoptera: Vespidae). Only *B. bassiana* has been reported to infect *V. vulgaris* (Glare et al. 1993, 1996). This finding represents the first report of any wasp-pathogenic *Hirsutella* species from Chile.

We found only a single wasp infected with this *Hirsutella* sp., and we were unable to culture it. Future studies involving more infected individuals will be needed to obtain essential genomic sequence data to confirm our morphologically based identification. However, it must also be noted that no concerted genomic survey of a wide range of species of *Hirsutella* has yet been undertaken, so genome-based identifications for most of these species remain unattainable at this time. Further, the new rules of nomenclature that took effect in 2012 have led to the synonymization of most *Hirsutella* species to species in the very species-rich genus *Ophiocordyceps* (Hypocreales: Ophiocordycipitaceae). In this new classification, *H. saussurei* is now treated as a synonym for its known sexual (teleomorphic) state, *Ophiocordyceps humbertii* (C. P. Robin) G. H. Sung, J.-M. Hywel-Jones & Spatafora (Hypocreales: Ophiocordycipitaceae) (Sung et al. 2007).

The findings reported here contribute to our knowledge of the natural fungal enemies of *V. vulgaris* in Chile. Our aim is to use this fungus as a biological control agent; therefore, future collections and successful in vitro isolations of this species will be required. Further studies are needed to clarify the identification this fungus, to isolate conidia, and to produce infectious formulations before this pathogen could be applied in the field to limit the populations of this serious invasive pest.

The authors thank Servicio Agrícola y Ganadero for providing the specimen of *V. vulgaris*, and the Laboratorio Regional Osorno for supporting this study in their facilities. The authors also thank Christian Luz for the critical review of the manuscript. This study was supported by FONDECYT (Fondo Nacional de Desarrollo Científico y Tecnológico, Chile) project 1141066.

Summary

In Chile, the invasive and noxious pest *Vespula vulgaris* (L.) (Hymenoptera: Vespidae) was first reported in 2011 in the Araucanía region and is currently distributed between Araucanía and Magallanes regions. In Mar 2015 (autumn), during an ongoing monitoring of funnel traps by the Servicio Agrícola y Ganadero, a fungus-infected individual was collected. The fungus was identified morphologically as a species of *Hirsutella* (Hypocreales: Ophiocordycipitaceae). This is the first report of any *Hirsutella* species on *V. vulgaris* in Chile. No in vitro cultures were successfully established from the infected insect.

Key Words: invasive species; natural enemy; entomopathogenic fungus

Sumario

En Chile la plaga invasora y nociva *Vespula vulgaris* (L.) (Hymenoptera: Vespidae) se reportó por primera vez el año 2011 en la Región de la Araucanía y actualmente se distribuye entre las regiones de la Araucanía y Magallanes. En marzo del 2015 (otoño) durante un monitoreo que estaba en curso con trampas de embudo por el Servicio Agrícola y Ganadero, un individuo micotizado fue colectado. El hongo se identificó morfológicamente como una especie de *Hirsutella* (Hypocreales: Ophiocordycipitaceae). Este es el primer reporte de cualquier

Hirsutella en *V. vulgaris* en Chile. No se consiguió realizar cultivos in vitro desde el insecto infectado.

Palabras Clave: especie invasora; enemigo natural; hongo entomopatógeno

References Cited

- Aruta C, Carrillo R. 1989. Identificación de hongos del orden Entomophthorales en Chile. III. Agro Sur 17: 10–18.
- Aruta C, Carrillo R, González S. 1974. Determinación para Chile de hongos entomopatógenos del género *Entomophthora*. I. Agro Sur 2: 62–70.
- Barrera R, Vidal C. 2013. Primer reporte de *Vespula vulgaris* (Linnaeus, 1758) (Hymenoptera: Vespidae) en Chile. Boletín de la Sociedad Entomológica Aragonesa (S. E. A.) 52: 277–278.
- Baz A, Cifrián B, Martín-Vega D. 2010. Distribution of the german wasp (*Vespula germanica*) and the common wasp (*Vespula vulgaris*) (Hymenoptera: Vespidae) in natural habitats in Central Spain as shown by carrion-baited traps. Sociobiology 55: 871–881.
- Beggs JR. 2000. Impact and control of introduced *Vespula* wasps in New Zealand, pp. 404–409 In Austin A, Dowton M [eds.], Hymenoptera: Evolution, Biodiversity and Biological Control. CSIRO Publishing, Clayton South, Victoria, Australia.
- Beggs JR. 2001. The ecological consequences of social wasps (*Vespula* spp.) invading an ecosystem that has an abundant carbohydrate resource. Biological Conservation 99: 17–28.
- Cooke MC. 1892. Vegetable Wasps and Plant Worms. Society for Promoting Christian Knowledge, London, United Kingdom.
- Dymock JJ, Forgie SA, Ameratunga R. 1994. A survey of wasp sting injuries in urban Auckland from December to April in 1991/2 and 1992/3. New Zealand Medical Journal 107: 32–33.
- Evans HC. 1989. Mycopathogens of insects of epigeal and aerial habitats, pp. 205–238 In Wilding N, Collins NM, Hammond PM, Webber JF [eds.], Insect-Fungus Interactions. Academic Press, London, United Kingdom.
- Evison SEF, Roberts KE, Laurenson L, Pietravalle S, Hui J, Biesmeijer JC, Smith JE, Budge G, Hughes WHO. 2012. Pervasiveness of parasites in pollinators. PLoS One 7: e30641.
- Fitch CD, Hoffman DR, Schmidt M. 2001. Cloning of a paper wasp venom serine protease allergen. Journal of Allergy and Clinical Immunology 107: S221.
- Gardner-Gee R, Beggs JR. 2012. Invasive wasps, not birds, dominate in a temperate honeydew system. Austral Ecology 38: 346–354.
- Glare TR, O'Callaghan M, Wigley PJ. 1993. Checklist of naturally occurring entomopathogenic microbes and nematodes in New Zealand. New Zealand Journal of Zoology 20: 95–120.
- Glare TR, Harris RJ, Donovan BJ. 1996. *Aspergillus flavus* as a pathogen of wasps, *Vespula* spp., in New Zealand. New Zealand Journal of Zoology 23: 339–344.
- Hodge KT. 1998. Revisionary studies in *Hirsutella* (Anamorphic Hypocreales: Clavicipitaceae). PhD dissertation, Cornell University, Ithaca, New York.
- Humber RA. 2012. Identification of entomopathogenic fungi, pp. 151–187 In Lacey LA [ed.], Manual of Techniques in Invertebrate Pathology. Academic Press, London, United Kingdom.
- King TP, Spangfort MD. 2000. Structure and biology of stinging insect venom allergens. International Archives of Allergy and Immunology 123: 99–106.
- King TP, Lu G, Gonzalez M, Qian N, Soldatova L. 1996. Yellow jacket venom allergens, hyaluronidase and phospholipase: sequence similarity and antigenic cross-reactivity with their hornet and wasp homologs and possible implications for clinical allergy. Journal of Allergy and Clinical Immunology 98: 588–600.
- Lester PJ, Gruber MAM, Brenton-Rule EC, Archer M, Corley JC, Dvorak L, Masciocchi M, Van Oystaeyen A. 2014. Determining the origin of invasions and demonstrating a lack of enemy release from microsporidian pathogens in common wasps (*Vespula vulgaris*). Diversity and Distributions 20: 964–974.
- Minter DW, Brady BL, Hall RA. 1983. Five hyphomycetes isolated from eriophyid mites. Transactions of the British Mycological Society 81: 455–471.
- Rose EAF, Harris RJ, Glare TR. 1999. Possible pathogens of social wasps (Hymenoptera: Vespidae) and their potential as biological control agents. New Zealand Journal of Zoology 26: 179–190.
- Samson RA, Evans HC, Latgé JP. 1988. Atlas of Entomopathogenic Fungi. Springer Verlag, New York, New York.
- Shimazu M, Glockling SL. 1997. A new species of *Harposporium* with two spore types isolated from the larva of a cerambycid beetle. Mycological Research 101: 1371–1376.

- Singh R, Levitt AL, Rajotte EG, Holmes EC, Ostiguy N, van Engelsdorp W, Lipkin WI, de Pamphilis CW, Toth AL, Cox-Foster DL. 2010. RNA viruses in hymenopteran pollinators: evidence of inter-taxa virus transmission via pollen and potential impact on non-*Apis* hymenopteran species. PLoS One 5: e14357.
- Sosa-Gómez DR, López-Lastra CC, Humber RA. 2010. An overview of arthropod-associated fungi from Argentina and Brazil. Mycopathologia 170: 61–76.
- Speare AT. 1920. On certain entomogenous fungi. Mycologia 12: 62–76.
- Sung GH, Hywel-Jones NL, Sung JM, Luangsa-ard JJ, Shrestha B, Spatafora JW. 2007. Phylogenetic classification of *Cordyceps* and the clavicipitaceous fungi. Studies in Mycology 57: 5–59.
- Tankersley MS, Ledford DK. 2015. Stinging insect allergy: state of the art 2015. Journal of Allergy and Clinical Immunology: in Practice 3: 315–322.
- Yamane S, Wagner RE, Yamane S. 1980. A tentative revision of the subgenus *Paravespula* of eastern Asia (Hymenoptera: Vespidae). Insecta Matsumurana 19: 1–46.