

## **A Maximum Concentration Bioassay to Assess Insecticide Efficacy Against Hemipteran Pests of Tomato**

Authors: De Marchi, Bruno Rossitto, Hennessey, Megan, Turechek, William, and Smith, Hugh

Source: Florida Entomologist, 106(2) : 133-136

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.106.0211>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# A maximum concentration bioassay to assess insecticide efficacy against hemipteran pests of tomato

Bruno Rossitto De Marchi<sup>1</sup>, Megan Hennessey<sup>1</sup>, William Turechek<sup>2</sup>, and Hugh Smith<sup>1,\*</sup>

Florida is a primary producer of fresh market tomatoes (*Solanum lycopersicum* L.; Solanaceae) in the US, with over 9,700 ha (24,000 acres) harvested in 2020 at a value of USD \$463 million (USDA NASS 2021). Stink bugs (Hemiptera: Pentatomidae) and leaf-footed bugs (Hemiptera: Coreidae) are occasional pests of Florida tomato (UF IFAS 2006). Until recently, this pest complex was a problem primarily on tomato in Florida on certified organic farms where broad spectrum insecticides are not used, and management efforts focused on trap cropping (Gordon et al. 2017). Since 2019, for reasons that are not clear, stink bugs have resurged as a problem on conventional tomato grown in the spring season (Jan–May) in southern Florida. The most prevalent species attacking tomato in Florida are a brown stink bug (*Euschistus quadrator* Rolston) and the southern green stink bug (*Nezara viridula* [L.]) (both Hemiptera: Pentatomidae) (UF IFAS 2006). In addition, leaf-footed bugs (*Leptoglossus phyllopus* (L.) and *Phthia picta* (L.) (both Hemiptera: Coreidae) also attack tomato in Florida (UF IFAS 2006). Stink bug feeding damage can result in major economic losses to tomato. Stylet punctures and salivary enzymes from stink bug feeding cause deformed fruit (“cat-facing”), discoloration on the outer skin of tomatoes, and internal tissue damage reducing the economic value of fruit (Peiffer & Felton 2014; Rice et al. 2014, 2017).

To determine if the resurgence of stink bugs as a pest in tomato in Florida was related to lack of insecticide efficacy, 8 stink bug populations were collected from commercial tomato farms in the spring of 2021. In addition, we were able to collect 1 leaf-footed bug population for testing. The insecticide screening focused on neonicotinoid and pyrethroid insecticides, because these 2 insecticide groups have tended to show the greatest efficacy against stink bugs (Willrich et al. 2003; Snodgrass et al. 2005; Kuhar & Komminga 2017; Santos-Júnior et al. 2021). We hypothesized that insecticide tolerant populations would show limited response to maximum concentration treatments of com-

monly used insecticides, providing evidence that tolerance was present and that resistance monitoring efforts were warranted.

Stink bug and leaf-footed bug populations were collected from 6 commercial tomato field sites in Manatee County, 1 site in Hillsborough County, and 1 site in Hernando County (all in Florida) from Apr through Jun 2021. The species collected were *N. viridula* (the southern green stink bug), *E. quadrator* (a brown stink bug), and *L. phyllopus* (the leaf-footed bug). Populations were maintained on lima bean (*Phaseolus lunatus* L.; Fabaceae) plants in cages in a greenhouse at the Gulf Coast Research and Education Center in Wimauma, Florida, USA, and provided with tomatoes and sweet corn for additional food. The greenhouse was cooled with cooling pads and ventilated with fans, maintaining temperatures between 18 and 35 °C. Insect colonies experienced natural daylight. Relative humidity was not recorded. Egg clusters, nymphal stages and adults were maintained in separate cages to avoid cannibalism. Insect populations were given approximately 30 d to establish on lima bean plants in the greenhouse, at which point the F<sub>1</sub> generation was tested for susceptibility to key insecticides using third instar nymphs and a maximum concentration bioassay.

Each stink bug and leaf-footed bug population was tested separately for susceptibility to 3 pyrethroid insecticides, 3 neonicotinoid insecticides, and a deionized water control (Table 1). The active ingredients evaluated were bifenthrin (Brigade® 2 EC, FMC Corporation, Philadelphia, Pennsylvania, USA), zeta-cypermethrin (Mustang® Maxx, FMC Corporation, Philadelphia, Pennsylvania, USA), lambda-cyhalothrin (Warrior, Syngenta, Greensboro, North Carolina, USA), dinotefuran (Venom® 70 SG, Valent USA, Walnut Creek, California, USA), imidacloprid (Admire® Pro, Bayer Corporation, Whippany, New Jersey, USA), and thiamethoxam (Actara®, Syngenta, Greensboro, North Carolina, USA). The maximum labeled per ha application rate for tomato was used for each treatment and was prepared in beakers containing 500

**Table 1.** Trade name, active ingredient (ai), mode of action group (MoA), and application rate of 6 insecticides tested in a laboratory bioassay to control stink bug (Hemiptera: Pentatomidae) and leaf-footed bug (Hemiptera: Coreidae) populations collected from Florida tomato farms, spring 2021.

Trade name	Active ingredient	MoA	Per ha rate	mL or g of product per 500 mL	mL or g of ai per 500 mL
Brigade®	Bifenthrin	3A	380 mL	0.338 mL	0.085 mL
Mustang®	Zeta-cypermethrin	3A	314 mL	0.28 mL	0.026 mL
Warrior	Lambda-cyhalothrin	3A	140 mL	0.125 mL	0.029 mL
Admire® Pro	Imidacloprid	4A	160 mL	0.143 mL	0.061 mL
Actara®	Thiamethoxam	4A	385 g	0.325 g	0.081 g
Venom®	Dinotefuran	4A	280 g	0.25 g	0.175 g

<sup>1</sup>University of Florida, Department of Entomology, Wimauma, Florida 33598, USA; E-mail: bruno.rossittode@ufl.edu (B. R. D. M.); meganhen@ufl.edu (M. H.); hughasmith@ufl.edu (H. A. S.)

<sup>2</sup>USDA ARS, US Horticultural Research Laboratory, Fort Pierce, Florida 34945, USA; E-mail: william.turechek@usda.gov (W. T.)

\*Corresponding author; E-mail: hughasmith@ufl.edu

mL of deionized water at a rate equivalent to 561 water L per ha (60 gallons water per acre). This is the typical volume used for foliar treatment of tomatoes when plants are in the vegetative stage (the first 6–8 wk after transplanting). The per ha application rate, amount of insecticide, and active ingredient in 500 mL of water are listed in Table 1. Organic grape tomatoes were purchased from the supermarket, rinsed, and dipped in the treatments. After 2 h air drying, 1 treated tomato was placed in a clear Genpak 32 oz plastic container (Genpak LLC, Sebring, Florida, USA) with 5 nymphs. Each container with 5 insects was 1 experimental unit or replicate. Each treatment was replicated 3 times. After 72 h, the number of live, moribund, and dead nymphs was recorded. For analysis of mortality, moribund nymphs were grouped with dead nymphs.

A general linear mixed model analysis was used to characterize the effects of insecticide treatment on insect mortality across 9 different populations. The experimental design was considered a completely randomized factorial design with each level of insecticide treatment and insect population crossed in each block. The independent variables population, treatment, and their interaction were treated as fixed effects. Replication was treated as a random effect. Prior to analysis, the dependent variable mortality was converted to proportions and subjected to the arcsine square root transformation; data were back-transformed for presentation in tables and figures. The analysis was performed using the GLIMMIX procedure of SAS (SAS, Version 9.4, Cary, North Carolina, USA) with the default identity link function and error Gaussian distribution. Additionally, the 'NOBOUND' option was used to allow negative values of covariance and scale parameters, which guards against an inflated Type I error rate (Littell et al. 2006). Moreover, the DDFM=KR option was used, which invokes the Kenward-Roger estimation of the standard errors and degrees-of-freedom. The general linear mixed model fit the data well based on visual inspection for a non-random pattern in a scatter plot of the residuals, a histogram with normal density, a linear quantile-quantile (or Q-Q) plot, and boxplot of the residuals that showed no outliers. Pairwise treatment differences for the fixed effects were obtained using the LSMEANS statement and LINES option ( $P = 0.05$ ). Multiple comparisons for all pairwise comparisons were based on Bonferroni  $t$ -tests ( $P = 0.05$ ), which keeps the maximum experiment-wise error rate at the intended type I error rate for all comparisons. The effect of insecticide on mortality was compared on the pooled populations using the Bonferroni  $t$ -test.

Insecticide ( $F_{6,124} = 54.80$ ;  $P < 0.0001$ ), population ( $F_{8,124} = 8.95$ ;  $P < 0.0001$ ), and the interaction ( $F_{48,124} = 2.27$ ;  $P < 0.0002$ ) had a significant effect on nymphal mortality observed 72 h after treatment.

The mortality of nymphs in the untreated controls ranged from 0.00 to 33.33% (Fig. 1). Six populations had zero mortality in the untreated control, 2 populations produced 6.7% mortality, and 1 population, Brown1, experienced 33.3% mortality in the untreated control. Mortality was lower in the untreated control than all insecticide treatments. Mortality was higher in bifenthrin than all other treatments with the exception of dinotefuran. Among the products tested, bifenthrin was the most effective pyrethroid with mortality ranging from 73.3 to 100%. Dinotefuran was the most effective neonicotinoid with mortality ranging from 46.67 to 86.67%. In contrast, imidacloprid produced consistently poor results, with mortality ranging from 0 to 60.00% (Fig. 1). Mortality was not statistically different between dinotefuran and zeta-cypermethrin, both of which produced significantly higher mortality than lambda-cyhalothrin, thiamethoxam, and imidacloprid. Mortality was higher in the lambda-cyhalothrin treatment than imidacloprid, but not significantly different from thiamethoxam. Mortality was not significantly different between the thiamethoxam and imidacloprid treatments (Fig. 1).

Stink bugs and leaf-footed bugs are generalist pests affecting tomato and other crops (Mitchell 2000; UF IFAS 2006; Braman & West-erfield 2020). Historically this pest complex has been managed with broad spectrum insecticides including carbamate, organophosphate, and pyrethroid insecticides (Willrich et al. 2003; Snodgrass et al. 2005). Neonicotinoid insecticides also have shown efficacy against this group (Tillman & Mullinix 2004; Tillman 2006; Kamminga et al. 2012). In addition, trap cropping has been evaluated against stink bugs and leaf-footed bugs with limited success (Gordon et al. 2017; Braman & West-erfield 2020; Leppla et al. 2023). Current management methods focus on chemical control based on the application of neonicotinoid and pyrethroid insecticides (Kamminga et al. 2012; Kuhar & Kamminga 2017). Our results were consistent with studies cited above demonstrating efficacy of pyrethroid and neonicotinoid insecticides.


The maximum concentration test provides a rapid screening of insecticide materials and can be used to illustrate patterns of susceptibility among pest populations across production regions (Riley et al. 2020; De Marchi et al. 2021). We subjected 5 populations of *N. viridula*, 3 populations of *E. quadrator*, and 1 population of *L. phyllopus* to a maximum concentration test to screen for susceptibility to 3 pyrethroid and 3 neonicotinoid insecticides commonly used to manage stink bugs. Our objective was to determine if most populations were susceptible to insecticides or if there was evidence of insecticide tolerance that might help explain the resurgence in stink bugs as pests in Florida tomato fields. Not all insecticides were equally effective, and there was variability in the overall susceptibility of populations studied. However, each population was susceptible to insecticides, with bifenthrin and dinotefuran being the most effective.

Our objective was to test populations promptly so that susceptibility profiles would be representative of responses in the field. As a result, we had to focus on nymphal stages and used a low number of replicates to have sufficient individuals to screen all 6 insecticides. However, the test served the purpose of providing information quickly to growers that insecticide resistance did not seem to be the primary reason for pest resurgence. One possible explanation for the resurgence of stink bug populations on some farms may be the substitution of cyantraniliprole in place of dinotefuran at-planting and in early vegetative stages for management of *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae) and tomato yellow leaf curl virus, which is transmitted by *B. tabaci*. Cyantraniliprole is effective against *B. tabaci* and other tomato pests, including leafminers (*Liriomyza* spp. Mik; Diptera: Agromyzidae) and lepidopteran larvae, but it is not labelled for use against stink bugs. In summary, judicious use of pyrethroid and neonicotinoid insecticides when hemipteran pest populations are first detected may mitigate damage caused by this pest group.

This research was funded with support from the Florida Tomato Committee. Laurie Chambers, Sherline Estaing, Gideon Alake, and Craig Frey assisted with this research.

## Summary

Hemipterans have resurged as a pest of tomato in south Florida. To determine if the problem was related to insecticide efficacy, populations of southern green stink bug (*Nezara viridula* [L.]; Hemiptera: Pentatomidae), brown stink bug (*Euschistus quadrator* Rolston; Hemiptera: Pentatomidae), and a leaf-footed bug (*Leptoglossus phyllopus* [L.]; Hemiptera: Coreidae) were collected from 8 commercial tomato farms in South Florida, and third instar nymphs were subjected to a maximum concentration bioassay. Insects from the  $F_1$  generation were confined in plastic containers with grape tomatoes treated with the highest labeled application rate of 6 insecticides commonly used for



**Fig. 1.** Mortality of third instar nymphs from 5 populations of *Nezara viridula* ('green'), 3 populations of *Euschistus quadrator* ('brown'), and 1 population of *Leptoglossus phyllopus* ('leaf') stink bug subjected to a maximum concentration test of 3 pyrethroid and 3 neonicotinoid insecticides and an untreated control (UTC). Percentage mortalities (+ SEM) are based on the mean of 5 nymphs per replicate with 3 replicates. Insecticides designated with the same letter in the horizontal axis are not statistically different (by Bonferroni t-test,  $P < 0.05$ ).

hemipteran management in tomatoes: bifenthrin, lambda-cyhalothrin, zeta-cypermethrin (pyrethroids), dinotefuran, imidacloprid, and thiamethoxam (neonicotinoids). Mortality was observed after 72 h. Bifenthrin was the most effective pyrethroid with mortality ranging from 73 to 100%. Dinotefuran was the most effective neonicotinoid with mortality ranging from 47 to 87%. In contrast, imidacloprid produced consistently poor results, with mortality ranging from 0 to 60%. Populations varied in their susceptibility to insecticides tested, but tests did not reveal evidence of lack of insecticide efficacy.

**Key Words:** *Nezara viridula*; *Euschistus quadrator*; pyrethroid; neonicotinoid

## Sumario

Los hemipteranos han resurgido como plaga del tomate en el sur de Florida. Para determinar si el problema estaba relacionado con la eficacia de los insecticidas, se recolectaron poblaciones del chinche hediondo verde del sur (*Nezara viridula* [L.]; Hemiptera: Pentatomidae),

el chinche hediondo café (*Euschistus quadrator* Rolston; Hemiptera: Pentatomidae) y un chinche de patas laminadas (*Leptoglossus phyllopus* [L.]; Hemiptera: Coreidae) de 8 granjas de tomate comerciales en el sur de Florida, y las ninfas de tercer estadio se sometieron a un bioensayo de concentración máxima. Los insectos de la generación  $F_1$  se limitaron en recipientes de plástico con tomates de uva tratados con la tasa de aplicación más alta etiquetada de 6 insecticidas comúnmente utilizados para el manejo de hemipteranos en tomates: bifentrina, lambda-cyhalothrin, zeta-cypermethrin (piretroides), dinotafurán, imidacloprida y thiametoxamam (neonicotinoides). La mortalidad se observó después de 72 horas. La bifentrina fue el piretroide más efectivo con mortalidad que varió del 73 al 100%. Dinotefuran fue el neonicotinoide más efectivo con mortalidad que varió del 47 al 87%. En contraste, el imidacloprid produjo resultados consistentemente pobres, con mortalidad que varió de 0 a 60%. Las poblaciones variaron en su susceptibilidad a los insecticidas probados, pero las pruebas no revelaron evidencia de falta de eficacia de insecticidas.

**Palabras Clave:** *Nezara viridula*; *Euschistus quadrator*; piretroide; neonicotinoide

## References Cited

- Braman SK, Westerfield B. 2020. Influence of trap crops on tomato and squash insect pests. *Journal of Entomological Science* 55: 578–583.
- De Marchi BR, Smith H, Turechek W, Riley D. 2021. A maximum dose bioassay to assess efficacy of key insecticides against *Bemisia tabaci* MEAM1 (Hemiptera: Aleyrodidae). *Journal of Economic Entomology* 114: 914–921.
- Gordon TL, Haseeb M, Kanga LHB, Legaspi JC. 2017. Potential of three trap crops in managing *Nezara viridula* (Hemiptera: Pentatomidae) on tomatoes in Florida. *Journal of Economic Entomology* 110: 2478–2482.
- Kamminga KL, Koppel AL, Herbert DA, Kuhar TP. 2012. Biology and management of the green stink bug. *Journal of Integrated Pest Management* 3: C1–C8.
- Kuhar TP, Kamminga K. 2017. Review of the chemical control research on *Halyomorpha halys* in the USA. *Journal of Pest Science* 90: 1021–1031.
- Littell RC, Milliken GA, Stroup WW, Wolfinger RD, Schabenberger O. 2006. SAS for Mixed Models, Second Edition. SAS Institute Inc., Cary, North Carolina, USA.
- Leppla NC, Stacey KJ, Rooney LM, Lennon KM, Hodges AC. 2023. Stink bug (Hemiptera: Pentatomidae) occurrence, reproduction, and injury to fruit in an organic tomato crop bordered by sorghum. *Journal of Economic Entomology* 116: 144–152.
- Mitchell PL. 2000. Leaf-footed bugs (Coreoidae), pp. 337–404 *In* Schaefer CW, Panizzi AR [Eds.], *Heteroptera of Economic Importance*. CRC Press, New York, USA.
- Peiffer M, Felton G. 2014. Insights into the saliva of the brown marmorated stink bug *Halyomorpha halys* (Hemiptera: Pentatomidae). *PLoS One* 9: e88483. doi: 10.1371/journal.pone.0088483
- Rice KB, Bergh CJ, Bergmann EJ, Biddinger DJ, Dieckhoff C, Dively G, Fraser H, Garipey T, Hamilton G, Haye T, Herbert A, Hoelmer K, Hooks CR, Jones A, Krawczyk G, Kuhar T, Martinson H, Mitchell W, Nielsen AL, Pfeiffer DG, Raupp MJ, Rodriguez-Saona C, Shearer P, Shrewsbury P, Venugopal PD, Whalen J, Wiman NG, Leskey TC, Tooker JF. 2014. Biology, ecology, and management of brown marmorated stink bug (Hemiptera: Pentatomidae). *Journal of Integrated Pest Management* 5: A1–A13.
- Rice KB, Troyer RR, Watrous KM, Tooker JF, Fleischer SJ. 2017. Landscape factors influencing stink bug injury in Mid-Atlantic tomato fields. *Journal of Economic Entomology* 110: 94–100.
- Riley D, Smith H, Bennett J, Torrance P, Huffman E, Sparks Jr A, Gruver C, Dunn T, Champagne D. 2020. Regional survey of diamondback moth (Lepidoptera: Plutellidae) response to maximum dosages of insecticides in Georgia and Florida. *Journal of Economic Entomology* 113: 2458–2464.
- Santos-Júnior JLC, Raetano CG, de Carvalho FK, Ferreira-Filho PJ, Guerreiro JC, Ferrari S, Nascimento V, Tomaz RS, Argioli AS, Prado EP. 2021. Effects of application rate on spray deposition, stink bug control, and yield in soybean crops. *Phytoparasitica* 50: 191–200.
- Snodgrass GL, Adamczyk JJ, Gore J. 2005. Toxicity of insecticides in a glass-vial bioassay to adult brown, green, and southern green stink bugs (Heteroptera: Pentatomidae). *Journal of Economic Entomology* 98: 177–181.
- Tillman PG. 2006. Susceptibility of pest *Nezara viridula* (Heteroptera: Pentatomidae) and parasitoid *Trichopoda pennipes* (Diptera: Tachinidae) to selected insecticides. *Journal of Economic Entomology* 99: 648–657.
- Tillman PG, Mullinix Jr BG. 2004. Comparison of susceptibility of pest *Euschistus servus* and predator *Podisus maculiventris* (Heteroptera: Pentatomidae) to selected insecticides. *Journal of Economic Entomology* 97: 800–806.
- UF IFAS (University of Florida Institute of Food and Agricultural Sciences). 2006. Grower's IPM Guide for Florida Tomato and Pepper Production. Gillette JL [Ed.]. [https://ipm.ifas.ufl.edu/agricultural\\_ipm/Growers\\_IPM\\_Guide\\_for\\_Florida\\_Tomato\\_and\\_Pepper\\_Production.html](https://ipm.ifas.ufl.edu/agricultural_ipm/Growers_IPM_Guide_for_Florida_Tomato_and_Pepper_Production.html) (last accessed 17 Mar 2023).
- USDA NASS. 2021. Vegetables Annual Summary. USDA Economics, Statistics and Market Information System (ESMIS), pp. 72–73. <https://usda.library.cornell.edu/concern/publications/02870v86p> (last accessed 17 Mar 2023).
- Willrich MM, Leonard BR, Cook DR. 2003. Laboratory and field evaluations of insecticide toxicity to stink bugs (Heteroptera: Pentatomidae). *The Journal of Cotton Science* 7: 156–163.