

High-Elevation Savanna Landscapes in the Cordillera Central, Dominican Republic, Hispaniola

Authors: Michael Clark, G., Horn, Sally P., and Orvis, Kenneth H.

Source: Mountain Research and Development, 22(3): 288-295

Published By: International Mountain Society

URL: https://doi.org/10.1659/0276-4741(2002)022[0288:HESLIT]2.0.CO;2

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

G. Michael Clark, Sally P. Horn, and Kenneth H. Orvis High-Elevation Savanna Landscapes in the Cordillera Central, Dominican Republic, Hispaniola

Dambos in the Caribbean?



288

Headwater unincised savanna (sabana) landscapes in the Cordillera Central, Dominican Republic, have a vegetation cover dominated by the native tussock grass, Danthonia domingensis. Sites occur on gentle-relief upland

surfaces approximately between 2000 and 2400 m, mainly in José del Carmen Ramírez National Park (JCR) and Juan B. Pérez Rancier National Park (JBPR). Surface and subsurface data from 1 site constrain regolith age overlying granitoid saprolite to <12,570 ¹⁴C years BP. We conclude that the savannas studied are dambos that developed in response to environmental change during and after the Pleistocene-Holocene transition. Dambos elsewhere are widely regarded as reliable indicators of major Quaternary environmental change. Dambos have unique hydrological, ecological, scenic, and utilitarian attributes; their importance will increase with greater tourism, as will the concerns for their sustainable use. This first publication on dambos in the Caribbean signals the need for more research into their origin, evolution, and state of equilibrium.

Keywords: Dambo; savanna.

Peer reviewed: February 2002. Accepted: March 2002.

Introduction

Small, treeless savanna landscapes occur on low- to moderate-relief uplands between 2000 and 2400 m that occupy low-relief topographic hollows and sideslopes without incised drainage lines in the Cordillera Central, Dominican Republic, Hispaniola (Figure 1; Table 1). These features lack previous scientific study despite their present and projected potential for basic and applied research. Sites are within José del Carmen Ramírez National Park (JCR) and Juan B. Pérez Rancier National Park (JBPR). Another savanna, El Valle de Bao in Armando Bermúdez National Park (AB), which borders JCR on the north, differs in climatic and geomorphic history and will be treated elsewhere.

The JCR sites are the Sabana Macutico principal site and the savannas to the northwest and west along the mule trail between the refugios of Macutico and Alto de la Rosa (Table 1). Trails are the only land transportation link in this area, creating logistical limitations for research. The Macutico site and other savannas in JCR

and JBPR bear striking similarities with geomorphic features described in the literature as dambos. Dambo is a Bantu term meaning "meadow grazing" and is extensively used to describe streamless, low-relief hollows and other low-slope-order grassland landforms in the plateau-like African uplands, dominantly south of the Equator to about 25° S (Mäckel 1974, 1985; Acres et al 1985; Boast 1990). Ackermann (1936) introduced the term dambo into scientific literature to describe channelless grass-covered openings in dry forest or bush vegetation that connect down-gradient to incised drainage systems. He restricted the term to headwater areas, excluding terraces but including areas inundated only during the rainy season. Classic areas occur in Malawi and Zambia (Boast 1990) and in the bordering countries. Other terms for these features include vleisforms the senior author observed in The Union of South Africa. Dambos may also exist in southern India and in South America (Boast 1990).

With increasing tourism, Dominican savannas will assume increasing importance despite their small areal extents in these montane landscapes (<1%). Savannas provide siting for *refugios* and grazing areas for mules, and constitute a tourist attraction. Although the preponderant vegetation is Danthonia, additional plant species are found that are either absent in the forest or occur sparingly. Savanna preservation may be necessary for survival of rare plant species and possibly endan-



FIGURE 1 Locations of JCR on the drier southwestern side of the topographic crest of the Cordillera Central, containing high-elevation savanna landforms. AB is on the northeastern side of the divide. The former Valle Nuevo Scientific Reserve, now Juan B. Pérez Rancier National Park (JBPR), a plateau-like upland separated from the Cordillera Central by a lowland, also contains a highelevation savanna landscape assemblage. Elevation range shading: darkest shading = 0-1000 m; intermediate shading = 1000-2000 m; lightest shading (almost all within parks) = >2000 m.

TABLE 1 Locations and elevations for selected savanna localities interpreted to be *dambos* on the southern flank of the Cordillera Central in JCR, and in JBPR. UTM Northings and UTM Eastings are field GPS readings except for JBPR sites Sabana Quéliz and Cerro de Sabana de la Cruz, which are from map grid data.

Site name (see map, Figure 1)	UTM Northing	UTM Easting	Elevation (m asl, map)	
JCR: Sabana Macutico	2,105,503-2,105,640	281,459–281,534	1980-2000	
JCR: Sabana de los Boquerones	2,106,565-2,107,768	277,721-279,059	2200-2240	
JCR: Sabana del Pino	2,108,177-2,108,241	277,786–277,811	2320-2340	
JBPR: Valle Nuevo	2,080,555	322,754	2300–2360	
JBPR: Sabana de los Robles	2,080,479	323,645	2272-2360	
JBPR: Sabana Quéliz	2,075,400-2,075,600	264,000–269,000	2260-2300	
JBPR: Cerro de Sabana de la Cruz	2,074,500-2,074,700	281,000-286,000	2240-2260	
JBPR: Nizao	2,070,791	330,185	2300–2380	
JBPR: La Nevera	2,069,010-2,069,144	330,940-331,017	2260-2380	

gered animals such as the large rodent *hutia* (Bolay 1997). Thus, the present and projected future importance of highland savannas cannot be overemphasized.

In this article we assess tectonic and climatic environments of these savanna landforms, characteristics of their landforms and regolith, and what is known of their historic, geomorphic, and ecologic environments. We then address the problems of classification and genesis and provide evidence that these features constitute two major types of *dambos*.

Study area and description

Introduction

The savannas have an essentially continuous vegetation cover of tussock grass (*Danthonia domingensis* Hack. & Pilger) in predominantly well-drained areas, and sedges and shrubs such as *Rubus* in poorly drained sites. Upvalley areas and sides of the treeless savannas are abruptly bordered by pure to nearly pure stands of the native West Indian pine (*Pinus occidentalis* Swartz). Down valley, savannas terminate at major topographic breaks, such as intersections with local perennial streams or small rivers.

Savannas occur in an intermediate altitudinal landscape belt between 2 other distinctive major landscape assemblages. Below 1500 m is an altitudinal belt with deeply incised, low-sinuosity, feral relief landscapes, typical of Caribbean montane environments (Ahmad et al 1993; Orvis et al 1997, figure 8, p 327). Above the savanna landscape belt are the high-peak areas with steep slopes, bedrock exposures, and blocky regolith. The intermediate savanna-containing landscape belt is composed of composite fluvial and hillslope terrains that respond dynamically to major precipitation events. Typical hillslope regoliths beneath these terrains are ancient, deeply weathered, saprolite profiles undergoing exhumation, and complex surficial diamicton deposits. Stream and small-river floodplain and terrace landforms are underlain by poorly to well-sorted coarse alluvium.

Tectonic geology and lithology

In the Dominican Republic the Cordillera Central is within the Greater Antilles Deformed Belt (Case and Holcombe 1980) or Greater Antilles Orogenic Belt (Draper et al 1994), an active 150-km-wide strike-slip zone forming a major part of the northern Caribbean Plate boundary. Main Cordillera Central uplift phases may have occurred in the Plio-Pleistocene (Lewis et al 1991). But Lewis et al (1990, p 101) note subsurface stratigraphic evidence in Valle del Cibao (Cibao Basin), north of the Cordillera Central, that suggests rapid Cordillera Central uplift in Early Miocene time. The present combination of dynamic and erosional relief produced is extreme for the Caribbean.

The Macutico site and related savannas in JCR are within the Tireo Stratigraphic Tectonic Terrane, containing Cretaceous lavas and tuffs with intercolated mudstone, siltstone, chert, and limestone. Most of these rocks have been metamorphosed to varying grades (Draper and Lewis 1991). Several large, unfoliated granitoid stocks and batholiths, mainly hornblende tonalites (Lewis et al 1990), intruded the volcaniclastic rocks and lavas between Late Cretaceous and Late Eocene times. These plutons and spatially associated rocks underlie the known savanna areas we traversed in JCR (Draper and Lewis 1991, figure 1, p 30).

The other area with savanna landforms and vegetation is JBPR, about 60 km southeast of the Cordillera Central High Peaks area (Figure 1) and separated from it by a lowland. The central part of the JBPR is a plateau-like upland, where Tireo rocks composed of tuff, limestone, and rhyolite are unconformably overlain by apparent mid-Tertiary conglomerates that are intruded by andesites and overlain by Plio-Pleistocene tuffs and basalts (Lewis et al 1991). It is unknown whether areas occupied by the main Cordillera Central to the northwest had a covermass of sedimentary rocks or sediments during the early to mid-Tertiary.

Present-day climate

The Cordillera Central forms an orographic barrier to the northeast trade winds and to trailing midlatitude fronts in winter. Vegetation patterns on the northern, windward flank indicate that precipitation in that area increases with elevation below the trade wind inversion at 2000–2500 m, then decreases. Summits are relatively arid. On the southern, leeward flank the maximum moisture appears to narrowly coincide with the cloud belt near 2400 m and decreases rapidly downslope.

Precipitation means at foothill stations south of the Cordillera, despite a stronger tropical storm component, are about 45% of those to the north (Horst 1992). But periods of prolonged precipitation do occur, and precipitation during hurricanes can be extreme. Thus the overall geomorphic effectiveness of precipitation on the southern flank of the Cordillera Central may be very significant over time, despite low annual means. Surface air temperature decreases markedly with increasing elevation. Horst (1992) reported an annual mean of 25.5°C at Santo Domingo (sea level), 21.5°C at Jarabacoa (529 m), and 18°C at Constanza (1164 m). Air temperatures below freezing occur in all months at all known sites approximately above 2100 m (Pedersen 1953) and at least as low as 1800 m, where cold air drainage is trapped. We are unaware of any records of snow cover, although local guides report the frequent occurrence of hailstorms, and hail may temporarily cover the land surface. Arctic air mass outbreaks reach Hispaniola on rare occasions, producing mountain temperatures at least as low as -8°C under dramatically clear skies.

Methodology

The principal research site, Sabana Macutico JCR is an incisionless savanna in Macutico (Figure 1; 19°02' N, 71°05' W; 1980–2000 m), about 3–8 km west–southwest of and 1000 m lower than the High Peaks altitudinal zone (Pico Duarte, 3102 m; Loma la Pelona, 3097 m; and adjacent peaks, >2500 m). The site occupies a small, low-gradient streamless valley trending down-gradient about 200 m to the NE, then about 300 m to the

	Horizon	Depth (cm)	Color (moist) matrix	Color (moist) mottles	Field texture (moist)	
	0	0–2.5	2.5Y/2/0	NA	Clay	
	A1	2.5–15	7.5YR/2/0	NA	Silty clay	
	A ₂	15–33	10YR/3/4	NA	Sandy clay loam	
	A ₃ / B ₁	33–46	10YR/4/4	10YR/6/8	Silty clay loam	
	B ₂	46-61	10YR/5/8	NA	Silty clay loam	
	B ₃	61–86	7.5YR/5/6	NA	Sandy loam	
	B ₄	86–97	7.5YR/5/8	NA	Clay loam	
	B ₅	97–132	10YR/5/8	10YR/6/4	Clay loam	
	B₅/C	132–155	7.5YR/5/8	10YR/7/8	Loam	
	Cr₁ 155–173 –		-	NA	-	
il	Cr ₂	173-	-	NA	-	

TABLE 2 Soil pit field description for Macutico Sabana. Field soil profile descriptions follow established guidelines (Soil Survey Division Staff 1993; Schoeneberger et al 1998; Soil Survey Staff 1999).



FIGURE 2 Approximate lower half of length of Sabana Macutico, January 1999. View is down gradient toward termination of savanna at its truncation by the channel of upper Río Macutico (in trees, at base of near ridge, photographic center). Small fanlike features can be seen entering the savanna on the left and right sides of the view. Debris slide scars on ridges date from Hurricane Georges (22 September 1998–23 September 1998, Class IV). (Photo by G. M. Clark)

Structure	Consistency	Boundary	Clay films	Rock fragments	Other features
Loose	Loose	Very abrupt	None	None	Charcoal dominant
Fine weak subangular blocky	Loose	Diffuse	None	None	-
Fine subangular blocky	Friable	Diffuse	None	About 1%	-
Fine subangular blocky	Friable	Diffuse	On ped faces	About 25%, very angular	Very abundant rock fragments
Fine weak subangular blocky	Friable	Diffuse	On ped faces and sand grains	None	Abundant charcoal
Fine weak subangular blocky	Friable	Diffuse	On ped faces and sand grains	None	_
Fine weak subangular blocky	Friable	Abrupt	On ped faces and sand grains	About 6%	Charcoal
Fine weak subangular blocky	Friable	Abrupt	On ped faces and sand grains	None	Charcoal uncal. ¹⁴ C date 12480 +/- 85
Fine weak subangular blocky	Friable	Abrupt	On ped faces	Very stony, rounded fragments	Basal 8 cm very stony
-	Very firm	Abrupt	-	-	Soft granitoid saprolite with minor metabasaltic fragments
-	Extremely hard	Diffuse	-	-	Firm saprolite with minor metabasaltic fragments

In March 1996 we used a tubiform root auger (Horn et al 1994) and took two 10-cm-diameter soil cores in 5-cm-depth intervals to 50 cm. Because charcoal >2 mm was present in every interval, and a radiocarbon calibrated age range from one of the cores of 11,060-10,290 cal years BP (2o range) was obtained at the 45- to 50-cm-depth interval (Horn et al 2000), we investigated this savanna in further detail. In January 1999 a soil pit $0.75 \times 2.4 \times 1.75$ m deep was excavated with hand tools in the medial position of the savanna. We also took an additional tubiform root auger core to a depth of 95 cm and excavated deeper samples with a soil auger. Geomorphic field procedures followed techniques given by Goudie (1990). Field soil procedures followed established guidelines (Soil Survey Division Staff 1993; Schoeneberger et al 1998; Soil Survey Staff 1999).

Results

292

Sabana Macutico

Sabana Macutico has low-gradient, concave-up cross sections and even gentler longitudinal profiles. Incision is lacking, and smoothness dominates the microtopography. With the exception of several boulders, the savanna soil surface is free of large rocks.

The 1999 tubiform root auger core segments contained charcoal in all sample intervals but one (30–37 cm) to a depth of 93 cm (Horn et al 2000, table 1: core 3). Sampling continued to greater depth with the use of spoon augering, reaching an interval of deeply weathered granitoid parent soil material (10YR/5/8) at 142–154 cm. Refusal was reached at 154–164 cm at the base of the next change in regolith to granitoid saprolite (10YR/5/8, with mottling, 10YR/8/4).

The soil pit exposed a solum overlying weakly stratified soil parent material overlying saprolite-derived grus overlying saprolite (Table 2). Soil parent material is interpreted as locally transported surficial sediment, probably emplaced by sheetwash (see Mäckel 1974). Soil profile development is moderate, evidenced by clay translocation into B-horizons. Winter soil temperature (22 January 1999) at 1-m depth averaged 13.3°C. The site's estimated mean annual air surface temperature is about 14°C at 2000 m, derived from the mean annual lapse rate (Dominican Republic temperature records, Orvis et al 1997, p 325), indicating a mesic soil temperature regime. The moderate degree of soil development at the site is consistent with drainage, the radiocarbon ages obtained, and available soil temperature and precipitation data.

Effects of intense precipitation events on the sabana surface are minimal, to judge from the paucity

of effects of Hurricane Georges (22 September 1998–23 September 1998, Class IV). Modifications to Sabana Macutico occurred at its downslope terminus with Río Macutico, with channel entrenchment and overbank deposition along the reach crossing the terminus of the sabana. Older, coarse overbank alluvium, vegetation disturbance, and coarse, poorly sorted terrace gravels in the transition zone between the lower end of the sabana proper and the near river bank indicate a similar history.

Other savanna areas

Other savannas studied in JCR and JBPR from 1995 to 1999 (Table 1) have similar vegetation and ecotone contrast with bordering forest environments, but some landforms vary to some extent from those of Sabana Macutico. Two types of savanna landform variation occur. Valley head savannas are low-relief, gently concaving upward incisionless valleys that slope toward modern drainage lines (eg, up-gradient areas of La Nevera and Nizao, JBPR). More common are the topographically more diverse sideslope savanna landscapes consisting of interspersed nose, sideslope, and hollow landforms with somewhat steeper slope gradients (eg, Sabana de los Boquerones, Sabana del Pino, JCR). In Africa, dambo siting and orientation are linked to fracture patterns in the underlying bedrock (Giardino and Mäckel 1985; Whitlow 1985). Bedrock fracture effects on Dominican savanna location and orientation may exist, but the observed bedrock exposures were insufficient for evaluation. Most forms lack fluvial incision, and slope breaks and channel incisions occur only at or near down-valley ends of savannas at incised fluvial networks. Effects of Hurricane Georges that were visible in January 1999 were confined to such preexisting fluvial channels and their margins.

The additional savannas observed on the dry southern flank of the Cordillera Central in JCR are of the sideslope type (Orvis et al 1997, figure 7, p 327). Bedrock underlying these savannas is mapped as the granitoid Macutico Batholith complex and related rocks (Draper and Lewis 1991, figure 1, p 30). Where exposed or sampled by tubiform root auger coring, augering, and shallow soil pits, the underlying regolith is of local provenance, although not residual until massive subsoil and saprolite are reached. Typical sections of the modern soil have histic surface and near-surface soil horizons underlain by mineral soils. Where reached, the underlying parent material is weathered granular rock.

Both valley head and sideslope savannas occur in JBPR. Mineral soil underlies histic surface and near-surface horizons, and deeper soil parent material typically is cobbly to bouldery regolith ranging in weathering state from moderately weathered to totally decomposed.

Discussion

The following synopsis emphasizes African *dambo* properties as described in the literature and comparisons with and contrasts against savanna characteristics in JCR and JBPR.

Most of the reported climates in African regions with dambo morphology exhibit pronounced seasonality with average annual precipitation of 800-1500 mm and a high recurrence interval of intense precipitation events (Bond 1967; Mäckel 1974) with resultant geomorphically effective sheetfloods (Thorbecke 1973). Except in areas where dambos are increasing their areal extent (usually headward), abrupt demarcation lines exist between bordering arboreal vegetation and grassy near-level dambo floors. Although other factors are cited, sharp contacts between grassland and the surrounding woodland are most commonly attributed to poor soil drainage in the dambos (Meadows 1984; Acres et al 1985; Whitlow 1985), as evidenced by the surface wetness and widespread existence of organic-rich horizons (Mäckel 1974; Meadows 1984). Wide soil texture variations are reported in African soil descriptions and reflect underlying and bordering rock types. Sandy textures are characteristic of dambo soil profiles derived from quartzite, granitic, and gneissic parent rock materials derived from bedrock beneath many African dambos (Meadows 1984; Acres et al 1985; Whitlow 1985).

Acres et al (1985) gave a 4-fold geomorphic classification of *dambos* that emphasized landscape position. These categories are headwater *dambos* with extremely gentle gradients; slope *dambos* that extend up the valley sides with greater slope gradients; hanging or scarp *dambos* on plateau edges; and river *dambos*, with flanking terraces due to subsequent fluvial incision. Savanna features in JCR and JBPR best fit their models of headwater *dambos* and slope *dambos*. Sabana Macutico is an excellent fit with the headwater *dambo* type, whereas several other savannas such as the Sabana de los Boquerones and Sabana del Pino in JCR fit descriptions of the slope *dambo* category. We, therefore, interpret these savannas to be *dambos*.

Most *dambo* sites in both major areas of the Dominican Highlands were not waterlogged during our observation periods (February 1995; March 1996; February 1997; May 1997; July 1998; January 1999–February 1999). Exceptions include small blockages behind levee-like features (Valle Nuevo site, JBPR) and small fans (Macutico site, JCR). But good drainage for pine forest growth may not always have existed or may not exist the year round. Histic surface horizons are common and may partly be relicts from earlier wetter conditions, raising an unanswered question about whether or not some *dambo* features are relict.

Given its long history and apparent frequency

(Horn et al 2000), fire is likely a major influence on savanna vegetation development and its interactions with the surficial geomorphology. Several prehistoric charcoal dates and common historic occurrences of lightning strikes and droughts in the pine forest suggest that many fires may be natural. Macroscopic charcoal (>2-mm sieve catchment) is present in all but 2 depth intervals in the 3 Sabana Macutico soil cores, and one of these intervals is at the base of the deepest core (93-95 cm, Horn et al 2000). During recent widespread fires in JCR and JBPR, complete burning of Danthonia, but with little mortality, occurred in many sites-bogs and other wet areas excepted. Abundant charcoal was produced during burns, and the dambos have burned in the past, as evidenced by abundant subsurface charcoal in the Macutico dambo and other investigated sites (Horn et al 2000). Frost may also inhibit pine invasion into the dambos. In the Haitian savannas of Acani, Boujcan Pierre, Phillipe, and Pistache between 1524 and 1829 m, frost killed seedlings and injured saplings of P. occidentalis during the winter of 1950-1951 (Pedersen 1953). Severe effects occurred in open low-slope gradient areas in pine plantations with poor air drainage and were rare on land with intermediate to high slope gradients. A Forest Division meteorological station at 5500 feet (1676 m) reported temperatures of 24°F (-4°C) in January and 27°F (-2°C) in February (Pedersen 1953). Temperatures as low as -8°C are reported in the Dominican Highlands, and we observed morning frost on Danthonia in savannas on several occasions during 1999. Human effect on *dambo* development is probably minimal-highland activity comprised only cattle grazing and lumbering, and in JBPR, limited agriculture immediately before park establishment.

Conclusions

Dominican savannas that we interpret to be dambos have an abrupt contrast in vegetation-from bordering pine forest cover on hillslopes to grass-dominated vegetation-that is remarkably similar to the reported characteristics of African dambos. Topography upslope from the ecotone does not markedly differ from that at the boundary. Judging from surface and near-surface rock fragments, soil parent rock material is visually identical to that in the savannas. Other factors, for example, ponding of cold air drainage and subsurface hydrology, apparently have dictated formative savanna environments, and those processes must have differed markedly from conditions in the bordering forest. Hillslope expression is more variable. Gentle concave-upward transverse profiles and low longitudinal gradients characterize most savannas, but more complex shapes occur. In map view, straight trends and sharp angular bends suggest the influence of bedrock fracture patterns.

Dambos are widely regarded as strong indicators of major Quaternary environmental change, and their origins may be linked to such instability (Meadows 1985; Thomas 1994). A speculative chronology has the Late Pleistocene–Holocene transition open with rain-shadow montane aridity, enhancing the prospects for fire, slow vegetation recovery, and slopewash. Abundant evidence of prehistoric fires is preserved in Dominican dambo sediments, although definitive links to climatic variation are not yet established (Horn et al 2000). Recolonization of dambo floors by pine seedlings on open areas could be inhibited further by severe frost accompanying outbreaks of northern cold air masses.

The present and projected future importance of highland *dambos* cannot be overemphasized and must be brought forth in Dominican National Park plans for sustainable mountain development. Research will be necessary for further understanding of the origin, evolution, and present state of stability of *dambos* in the Cordillera Central. Subsurface information is especially needed to assess the 3-dimensional nature of the regolith-weathered bedrock surface and to study the effects of *dambo* formative process mechanisms that operate underground.

AUTHORS

G. Michael Clark

Department of Geological Sciences, 306 Geological Sciences Building, The University of Tennessee, Knoxville, TN 37996-1410, USA. clarkgmorph@utk.edu

Sally P. Horn

Department of Geography, The University of Tennessee, 304 Burchfiel Geography Building, Knoxville, TN 37996-0925, USA. shorn@utk.edu

Kenneth H. Orvis

Department of Geography, The University of Tennessee, 304 Burchfiel Geography Building, Knoxville, TN 37996-0925, USA. orvis@utk.edu

ACKNOWLEDGMENTS

This research was supported by the National Science Foundation (SBR-9809798 and SBR-9512484), the Committee on Research and Exploration and the Education Foundation of the National Geographic Society, the Tennessee Geographic Alliance, the Stewart K. McCroskey Memorial Fund, and The University of Tennessee. Additional support was provided by the Panasonic Corporation and the Fundación Moscoso Puello (FMP). We thank the National Parks Directorate for granting permission for this work and FMP Executive Director Andrés Ferrer for critical logistical and financial support, as well as field assistance. For field help we also thank R. Rodríguez, R. Briones, L. Kennedy, J. Speer, A. Walter, J. Rosado, M. Mejía, R. García, G. Drinnon, J. Clouse, C. Kelly, A. Levin, D. Ray, M. Snelson, M. Young, I. Pérez, B. Rámos, and our backcountry guides and muleteers. Constructive comments from two anonymous reviewers greatly aided in manuscript improvement.

REFERENCES

Ackermann E. 1936. Dambos in Nordrhodesien. Wissenschaftliche Veröffentlichungen des Deutschen Museums für Länderkunde, Leipzig. N.F. 4:147–157.

Acres BD, Blair-Rains A, King RB, Lawton RM, Mitchell AJB, Rackham LJ. 1985. African dambos: their distribution, characteristics and use. Zeitschrift für Geomorphologie. Supplementband 52:63–86.

Ahmad R, Scatena FN, Gupta A. 1993. Morphology and sedimentation in Caribbean montane streams. Examples from Jamaica and Puerto Rico. Sedimentary Geology 85(1–4):157–169.

Boast R. 1990. Dambos: a review. Progress in Physical Geography 14(2):153–177.

Bolay E. 1997. The Dominican Republic: A Country Between Rain Forest and Desert. Contributions to the Ecology of a Caribbean Island. Weikersheim, Germany: Margraf.

Bond G. 1967. River valley morphology, stratigraphy, and palaeoclimatology in southern Africa. *In:* Bishop WW, Clark JD, editors. *Background to Evolution in Africa*. Chicago: The University of Chicago Press, pp 303–312. **Case JE, Holcombe TL.** 1980. Geologic-Tectonic Map of the Caribbean Region. United States Geological Survey [USGS], Miscellaneous Investigations Series Map I-1100. Scale 1:2,500,000. Reston, VA: USGS.

Draper G, Jackson TA, Donovan SK. 1994. Geologic provinces of the Caribbean Region. *In:* Donovan SK, Jackson TA, editors. *Caribbean Geology: an Introduction.* Mona, Kingston, Jamaica: The University of the West Indies Publishers' Association, pp 3–12.

Draper G, Lewis JF. 1991. Metamorphic belts in central Hispaniola. Geological Society of America Special Paper 262:29–45.

Giardino JR, Mäckel R. 1985. Correlative development of dambos and dwalas: plateau regions of Zambia. Zeitschrift für Geomorphologie, Supplementband 52:187–200.

Goudie A. 1990. Geomorphological Techniques. London: Unwin Hyman.

Horn SP, Orvis KH, Kennedy LM, Clark GM. 2000. Prehistoric fires in the highlands of the Dominican Republic: evidence from charcoal in soils and sediments. *Caribbean Journal of Science* 36:10–18.

Horn SP, Wallin TR, Northrop LA. 1994. Nested sampling of soil pollen, charcoal, and other soil components using a root corer. *Palynology* 18: 87–89.

Horst OH. 1992. Climate and the "encounter" in the Dominican Republic. Journal of Geography 91(5):205–210.

Lewis JF, Amarante A, Bloise G, Jiménez GJG, Dominguez HD. 1991. Lithology and stratigraphy of upper Cretaceous volcanic and volcaniclastic rocks of the Tireo Group, Dominican Republic, and correlations with the Massif du Nord in Haiti. Geological Society of America Special Paper 262:143–163.

Lewis JF, Draper G, Bourdon C, Bowin C, Mattson P, Maurrasse F, Nagle F, Pardo G. 1990. Geology and tectonic evolution of the northern Caribbean margin. *In:* Dengo G, Case JE, editors. *The Caribbean Region.* The Geology of North America, Volume H. Boulder, CO: Geological Society of America, pp 77–140.

Mäckel R. 1974. Dambos: a study in morphometric activity on the plateau regions of Zambia. *Catena* 1:327–365.

Mackel R. 1985. Dambos and related landforms in Africa. An example for the ecological approach to tropical geomorphology. *Zeitschrift für Geomorphologie, Supplementband* 52:1–23.

Meadows ME. 1984. Past and present environments of the Nyika Plateau, Malawi. Palaeoecology of Africa and the Surrounding Islands 16:353–390. **Meadows ME.** 1985. Dambos and environmental change in Malawi, Central Africa. Zeitschrift für Geomorphologie, Supplementband 52:147–169.

Orvis KH, Clark GM, Horn SP, Kennedy LM. 1997. Geomorphic traces of Quaternary climates in the Cordillera Central, Dominican Republic. *Mountain Research and Development* 17:323–331. **Pedersen A.** 1953. Frost damage in the pine forest. *Caribbean Forester* 2(3–4):93–96.

Schoeneberger PJ, Wysocki DA, Benham EC, Broderson WD. 1998. Field Book for Describing and Sampling Soils. Lincoln, NE: Natural Resources Conservation Service, USDA [US Department of Agriculture], National Soil Survey Center.

Soil Survey Division Staff. 1993. Soil Survey Manual. US Department of Agriculture, Agricultural Handbook 18. Washington, DC: US Government Printing Office. Soil Survey Staff. 1999. Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys. 2nd ed. US Department of Agriculture, Natural Resources Conservation Service, Agricultural Handbook 436. Washington, DC: US Government Printing Office.

Washington, DC: US Government Printing Office. **Thomas MF.** 1994. Geomorphology in the Tropics. Chichester: Wiley. **Thorbecke F.** 1973. Landforms of the savanna zone with a short dry season. In: Derbyshire ED, editor. Climatic Geomorphology. New York: Barnes and Noble, pp 96–103.

Whitlow JR. 1985. Dambos in Zimbabwe: a review. Zeitschrift für Geomorphologie, Supplementband 52:115–146.