

Response to Comment by Craig Loehle (MRD Vol 30 No 3) on Bao et al (2010)

Authors: Bao, Kunshan, Yu, Xiaofei, Jia, Lin, and Wang, Guoping

Source: Mountain Research and Development, 30(4) : 411-412

Published By: International Mountain Society

URL: <https://doi.org/10.1659/MRD-JOURNAL-D-10-00100.1>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Response to Comment by Craig Loehle (MRD Vol 30 No 3) on Bao et al (2010)

Recent Carbon Accumulation in
Changbai Mountain Peatlands,
Northeast China (MRD Vol 30 No 1)

We appreciate Dr Loehle's comments (Loehle 2010) on our recent article (Bao et al 2010). We agree with his contention that the peat zones we analyzed using ^{210}Pb dating for short-term carbon (C) accumulation history are within the acrotelm (aerobic layer), and further analyses are necessary to evaluate the status of peatlands in the Changbai Mountains. However, we disagree with his view that our analysis needs to take account of loss of peat mass and fails to contribute to global C budgets.

As stated in our paper and shown by previous research (eg Turetsky et al 2004), it is important to focus research efforts on C cycling of recently accumulated peat, not only because global warming will most likely affect the mass balance near peat surfaces where peat temperature fluctuates with air temperature, but because estimating short-term rates of peat accumulation in the acrotelm is still poorly understood and studied relative to assessing the long-term rate of C accumulation. In addition, the ^{210}Pb dated interval of peat (the last 100–150 years) is generally assumed to be acrotelm only, even though we know that acrotelm may best be defined by water table depths (Yu et al 2001). This is why we quantified recent rates of C accumulation by collecting and examining short peat cores (< 45 cm), but did not provide additional data on onsite water-level monitoring and measurements to distinguish between the near-surface layer (acrotelm) and the underlying layer (catotelm).

Dr Loehle states that “unfortunately, their analysis assumed that

there is no loss of peat mass to the atmosphere.” In fact, it is possible to neglect peat decomposition when only estimating apparent rates of C accumulation from the thickness of peat deposits. We think Dr Loehle may not have taken account of the 3 rates of estimation of C accumulation, namely, the long-term (apparent) rate of C accumulation (LORCA), the recent (apparent) rate of C accumulation (RERCA), and the actual net rate of C accumulation (ARCA). LORCA and RERCA are analogous and can be estimated for a given peatland from peat columns of known (dry) bulk density, C content, and age. Differences exist in age determination. The former dates the basal peat and the latter is based on the given dated horizon in a surface core. ARCA can only be estimated by modeling approaches like Clymo-type peat accumulation (eg Clymo 1984) and is lower than the apparent rate of C accumulation because the slow decay that takes place in the anoxic deeper layers is ignored in this approach (Tolonen and Turunen 1996). We multiply bulk density (g/cm^3) and depth (cm) to calculate cumulative peat mass (g/cm^2), and then use ^{210}Pb ages and C contents to calculate recent C accumulation rates ($\text{g}/\text{cm}^2/\text{y}$) for each pair of dated intervals. The calculated rates are apparent rates—namely RERCA as observed at present from the peat cores—so we do not need to take into account peat loss due to decomposition.

Dr Loehle also thinks that proper estimation of peat accumulation rates in the top zone requires gas exchange measurements integrated over the year, and thus in his view our analysis is unsuitable for global C cycling. We do not agree with this, as we believe he overstates the role of gas exchange measurements in assessing C dynamics in peatlands. It is certainly possible to estimate directly both net primary production and decomposition rates in peatlands. Methodological limitations, however, combined with typi-

cally large spatial and temporal variability, lead to substantial errors in extrapolating from direct measurements to annual C budgets in near-surface peat (Turetsky et al 2004). Moreover, peatlands have some unique and complicated features that cannot be easily accommodated in standard ways, as in other ecosystems, because they are ecosystems located between upland and aquatic ecosystems and are commonly inaccessible on the ground (Wieder 2001). Changbai Mountain peatlands are locations where continuous flux measurements are either not available or not feasible. As a result, the method we use is an alternative approach to estimating the short-term C sink capacity of a peatland; it takes advantage of the long-term records stored in vertically accumulating peat and allows for the quantitative estimation of recent rates of organic matter accumulation. For these reasons, we continue to maintain that our results in Bao et al (2010) can contribute to the global C cycle and climate change research as a start in evaluating the Changbai Mountain peatlands.

REFERENCES

- Bao K, Yu X, Jia L, Wang G.** 2010. Recent carbon accumulation in Changbai Mountain peatlands, northeast China. *Mountain Research and Development* 30(1):33–41. doi: 10.1659/MRD-JOURNAL-D-09-00054.1.
- Clymo RS.** 1984. The limits to peat bog growth. *Philosophical Transactions of the Royal Society of London B* 303:605–654.
- Loehle C.** 2010. Is peat accumulation really accelerating in northeast China? Comment on Bao et al (2010): Recent carbon accumulation in Changbai Mountain peatlands, northeast China (MRD vol 30 no 1). *Mountain Research and Development* 30(3):316. doi: 10.1659/MRD-JOURNAL-D-10-00058.1.
- Tolonen K, Turunen J.** 1996. Accumulation rates of carbon in mires in Finland and implications for climate change. *The Holocene* 6(2):171–178.
- Turetsky MR, Manning SW, Wieder RK.** 2004. Dating recent peat deposits. *Wetlands* 24(2):324–356.
- Wieder RK.** 2001. Past, present and future peatland carbon balance—An empirical model based on ^{210}Pb -dated cores. *Ecological Applications* 7:321–336.
- Yu Z, Turetsky MR, Campbell ID, Vitt DH.** 2001. Modelling long-term peatland dynamics. Processes and rates as inferred from litter and peat-core data. *Ecological Modelling* 145:159–173.

AUTHORS

Kunshan Bao¹, Xiaofei Yu¹, Lin Jia², and Guoping Wang^{1*}

* Corresponding author:
wangguoping@neigae.ac.cn

¹ Key Laboratory of Wetland Ecology and

Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, 3195 Weishan Road, Gaoxin District, Changchun 130012, Jilin, China

² Department of Geographical Environment and Human Health, Institute of Geographic Sciences and Natural Resources Research,

Chinese Academy of Sciences, 11A Datun Road, Anwai District, Beijing 100101, China

Open access article: please credit the authors and the full source.