

# Expression and Function of Xmsx-2B in Dorso-Ventral Axis Formation in Gastrula Embryos

Authors: Onitsuka, Izumi, Takeda, Masatoshi, and Maéno, Mitsugu

Source: Zoological Science, 17(8): 1107-1113

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zsj.17.1107

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Expression and Function of *Xmsx-2B* in Dorso-Ventral Axis Formation in Gastrula Embryos

Izumi Onitsuka, Masatoshi Takeda and Mitsugu Maéno\*

Department of Biology, Niigata University, Niigata 950-2181, Japan

ABSTRACT—Msx is a homeodomain-containing transcriptional factor that plays an essential role in pattern formation in vertebrata and invertebrata embryos. In Xenopus laevis, two msx genes have been identified (Xmsx-1 and Xmsx-2). In the present study, we attempted to elucidate the expression and function of Xmsx-2B (formerly designated as Xhox7.1') in early embryogenesis. Whole mount in situ hybridization analyses showed that the expression pattern of Xmsx-2B at gastrula and neurula stages was very similar to that of Xmsx-1: the transcript of Xmsx-2B was observed in ventral and lateral sides of the embryo. At the tailbud stage, however, the expression pattern of Xmsx-2B in neural tissues was distinct from that of Xmsx-1. An RNA injection experiment revealed that, like BMP-4, Xmsx-2B has a strong ventralizing activity. In the Xmsx-2B-injected embryos, differentiation of axial structures such as the notochord, muscle, and neural tissue was completely suppressed, whereas  $\alpha$ -globin mRNA, a blood cell marker, was highly expressed. Simultaneous injection of Xmsx-1 and Xmsx-2B RNAs showed that they function in an additive manner. It was also shown that coinjection of Xmsx-2B with a dominant-negative BMP-4 receptor (tBR), which can induce formation of secondary axis when injected alone in ventral blastomeres, suppressed secondary axis formation. Furthermore, Xmsx-2B also suppressed secondary axis formation, which was induced by a dominant-negative form of Xmsx-1 (VP16/msx-1). Therefore, like Xmsx-1, Xmsx-2B is a downstream nuclear factor of the BMP-4derived ventralizing signal, and these two factors probably share the same target molecules. In conclusion, Xmsx-1 and Xmsx-2B function in dorso-ventral axis formation in early Xenopus laevis development.

#### INTRODUCTION

A number of studies have indicated that BMP-4 (bone morphogenetic protein-4), a member of the TGF-  $\beta$  superfamily, is a key regulator in body planning, especially in dorsoventral patterning of mesodermal tissues and in neural tissue differentiation, in the Xenopus laevis embryo (Dale et al., 1992; Jones et al., 1992; Fainsod et al., 1994; Graff et al., 1994; Maéno et al., 1994; Suzuki et al., 1994; Schmidt et al., 1995). Recently, homeodomain-containing transcriptional factors that mediate the BMP-4 signal have been isolated and analyzed: Xmsx-1 (Maeda et al., 1997; Suzuki et al., 1997), Xvent-1 (PV. 1) (Gawantka et al., 1995; Ault et al., 1996), Xvent-2 (Xbr-1, Vox, Xom) (Onichtchouk et al., 1996; Papalopulu and Kintner 1996; Ladher et al., 1996) and Xvex-1 (Shapira et al., 1999). We and other investigators have revealed that Xmsx-1 is expressed in the prospective ventral ectoderm and mesoderm at mid to late gastrula stages and that it has a ventralizing activity if its RNA is injected into the prospective dorsal region (Maeda et al., 1997; Suzuki et al., 1997). In ectodermal cells, Xmsx-1 functions as an inhibitor of neurogenesis, and it has

\* Corresponding author: Tel. +81-25-262-6183; FAX. +81-25-262-6116. E-mail maenobio@sc.niigata-u.ac.jp been shown that *Xmsx-1* is a direct target gene of the BMP-4 signal (Suzuki *et al.*, 1997). More recently, we have demonstrated that *Xmsx-1* is an upstream factor of Xwnt-8 in the ventralizing signal cascade and a possible direct repressor of organizer genes such as *goosecoid* (Takeda *et al.*, 2000).

In comparison to the numerous studies on Xmsx-1, quite a few studies have been carried out on the function of Xmsx-2 (Su et al., 1991; Gong and Kiba, 1999). In other species, msx-2 has been studied extensively in various developing tissues such as bone, tooth, and limb bud (Monachan et al., 1991; Coelho et al., 1991; Mackenzie et al., 1992; Jowett et al., 1993). Most likely, the function of msx-2 overlaps with that of msx-1 (Semenza et al., 1995), but in some cases msx-2 may have distinct expression patterns and functions. For instance, expression domains of msx-1 and -2 overlap in some regions but are discrete along with anterior-posterior identity in chicken and mouse limb buds (Davidson et al., 1991; Coelho et al., 1991; Muneoka and Sassoon, 1992; Nohno et al., 1992). The BMP-4/msx-2 signal is involved in elimination of cranial neural crest cells in rhombomeres 3 and 5 (Graham et al., 1993; Graham et al., 1994; Takahashi et al., 1998; Farlie et al., 1999) and the interdigit regions of developing limbs (Zou and Niswander 1996; Ganan et al., 1996; Ferrari et al., 1998) by mediating apoptosis. Furthermore, the mechanisms of expressional control in mouse mammary development are different between *msx-1* and *-2* (Friedmann *et al.*, 1996; Phippard *et al.*, 1996).

These observations suggest that msx-2 could have a distinct expression pattern and function in the early development of Xenopus laevis. In spite of the importance of the issue, expressional and functional natures of msx-2 in Xenopus embryogenesis are poorly understood. Xenopus msx-2 (Xhox7.1') was first isolated by Su et al. (1991). In their report, however, major attention was paid to the expression of Xmsx-1. Afterward, no report has described the expression and function of Xmsx-2 except for a very recent one in which a related but distinct gene from Xhox7.1' was described (Gong and Kiba, 1999). In this study, we attempted to elucidate the expression and the function of Xhox7.1'. We propose designating Xhox7.1' as Xmsx-2B in order to distinguish it from Xmsx-2, which was isolated by Gong and Kiba (1999). Our results showed that there is a strong correlation between the expression patterns of Xmsx-1 and Xmsx-2B at gastrula to neurula stages and that those two genes play a role in the establishment of the dorso-ventral axis in developing Xenopus embryos.

#### MATERIALS AND METHODS

#### **Cloning and cDNA construction**

Xmsx-2B cDNA fragments of two different sizes were amplified by polymerase chain reaction (PCR) from Xenopus stage10 cDNA. A short fragment including 402 N-terminal base pairs was used as a probe for in situ hybridization, and another fragment including the whole protein-coding region (815 bp) was used for an RNA injection assay. The primer sequences for the PCR were as follows: 5'-TC T-GCG-CCC-CTT-TAC-TCA-CCG-CA-3' and 5'-TCA-AAG-TGC-AAG-AGG-ATG-GGC-TCA-3' for the 402-bp fragment; and 5'-TCT-GCG-CCC-CTT-TAC-TCA-CCG-CA-3' and 5'- CTA-TGT-CAT-GTC-ACC-CTC-CTC-AGA-TA-3' for the 815-bp fragment (Su et al., 1991). Each product was ligated into pCRTM II (In vitro gene) for subcloning, and the DNA sequence of the inserts was determined using an Auto Read Sequencing kit (ALF DNA Sequencer, Pharmacia Biotech). The obtained sequence was completely identical to that of Xhox-7.1', which was previously reported (Su et al., 1991). For injection of sense RNA of Xmsx-2, an 815-bp fragment including an open reading frame was excised with Not I, blunt ended, and ligated into a blunt-ended pCS2+ expression vector.

#### Embryos, mRNA micro-injection, and DAI judgement

Xenopus laevis embryos were obtained by artificial insemination after induction of females with 250 i.u. of human chorionic gonadotropin. Fertilized embryos were dejellied with 2.5% thioglycolic acid (pH8.3) and washed several times in 5% MMR. After the embryos were transferred into 3% Ficoll/Steinberg's solution, equatorial regions of either two ventral or dorsal blastomeres at the 4-cell stage were injected with mRNAs using a Nanoject microinjector (3.7-6 nl/embryo). pCS2+ plasmids containing Xmsx-1 whole cDNA (1.8 kb) or the 815bp Xmsx-2B fragment were linearized with Xho I, and capped mRNA was synthesized using SP6 polymerase (Ambion). VP16/msx-1, a dominant-negative form of Xmsx-1, has been described previously (Takeda et al., 2000). Injected embryos were incubated in 50% Steinberg's solution at 23°C until the sibling embryos reached stage 33/34 for determining DAI (dorso-anterior index) as described by Kao and Elinson (1988). Developmental stages were designated according to Nieuwkoop and Faber (1967).

#### Whole-mount in situ hybridization

Whole-mount in situ hybridization was performed as described previously (Shain et al., 1996). Digoxygenin-labeled antisense mRNA probes were synthesized from the following plasmids: Xmsx-1, an Nterminal fragment (EcoRI/HindIII, 440 bp) in pSP72; and Xmsx-2B, an N-terminal fragment (402 bp) in pCRTMII. The former plasmid was linearized with EcoRI, and RNA was synthesized by SP6 polymerase. The latter plasmid was linearized with BamHI, and RNA was synthesized by T7 polymerase. The positive signals were visualized using BM purple (Moors et al., 1995). For neural marker probes, fragments of slug and krox-20 were amplified by the PCR method from the cDNAs of stage18 whole embryo and of stage 27 head tissue, respectively. The primer sequences for the PCR were as follows: 5'-GAA-TCT-GGT-TGC-TGT-GTA-G-3' and 5'-CGT-TGC-TGG-ATT-GTC-TAG-GC-3' for slug; and 5'-CCA-ACC-GCC-CCA-GTA-AGA-CC-3' and 5'-GTG-TCA-GCC-TGT-CCT-GTT-AG-3' for krox-20. Both fragments were subcloned into a pCR2.1 vector (In vitro gene), and their DNA sequences were determined. For making an antisense RNA probe, the plasmids were cut with BamHI, and RNAs were synthesized by T7 polymerase.

#### RESULTS

#### Expression pattern of Xmsx-2B in the early embryo

To investigate the expression pattern of Xmsx-2B during embryogenesis and to compare it with that of Xmsx-1, wholemount in situ hybridization analysis was performed using various stages of albino embryos. As shown in Fig.1, during gastrula and early neurula stages, the Xmsx-2B transcript was observed in the ventral and lateral marginal zones and animal pole area (Fig.1 B, D and F), as was also found in the case of Xmsx-1 (Maeda et al., 1997; Suzuki et al., 1997; Fig.1A, C and E). At the neurula stage (stages 16–17), two parallel lines of transcript were observed at the dorsal side along the border between the prospective neural tissue and epidermis (arrowheads in Fig. 1G and H), and also strong spot signals were detected on the medial and lateral sides of the prospective neural crest (arrows in Fig. 1 G and H), with reference to the expression of slug, a neural crest marker (Mayor et al., 1995; Fig. 1 J). We also compared the site of expression with that of krox-20, a marker of rhombomeres 3 and 5 (Bradley et al., 1992; Fig.1 I and L), and we found that strong spot signals were located at the level of the hind brain (rhombomeres 3-5) (Fig. 1 H and K). At the early tailbud stage (stage 23), expression patterns became different between Xmsx-1 and Xmsx-2B in the neural tissue (Fig.1 M, N, P and Q). Xmsx-1 was highly expressed in the broad areas including fore-, mid- and hind-brains, according to the expression of krox-20 (Fig.1 O and R), and spinal cord, while Xmsx-2B was expressed in the medial region of neural tissues. These observations suggest that Xmsx-1 and Xmsx-2B have the same or similar functions in gastrula stages but have different functions after the neurula stage.

#### *Xmsx-2B* has a ventralizing activity

It has been shown in previous studies that *Xmsx-1* has ventralizing and anti-neuralizing activities in early embryogenesis (Maeda *et al.*, 1997; Suzuki *et al.*, 1997; Ishimura *et al.*, 2000; Takeda *et al.*, 2000). To examine whether *Xmsx-2B* 



**Fig. 1.** Whole-mount *in situ* hybridization showing the expressions of *Xmsx-1* (A, C, E, G, M, P), *Xmsx-2B* (B, D, F, H, K, N, Q), *slug* (a neural crest marker) (J), and *krox-20* (a hindbrain marker) (I, L, O, R). At the mid gastrula stage (st.12), *Xmsx-1* and *Xmsx-2B* mRNAs are expressed in the ventral and lateral marginal zone (A–D) (A and B, vegetal view and dorsal side is toward the top; C and D, lateral view and dorsal side is to the right). At the late gastrula stage (st.13), transcripts are accumulated in the lateral regions adjacent to the presumptive neural fold (E, F). (E and F, dorsal view and anterior is toward the top). At the neurula stage (st.16–17), transcripts are detected along the border lines of prospective neural and non-neural tissues (arrowheads in G and H), and high levels of expressions are observed on the medial and lateral sides of the prospective neural crest cells (arrows in G and H), as shown by *slug* expression (J). These regions correspond to the level of the prospective hindbrain (rhombomeres 3-5), with reference to the expression of *krox-20* (I and L) (G–J, dorsal view and anterior is toward the top). At the tailbud stage (st.23), the expression patterns of *Xmsx-1* and *Xmsx-2B* in neural tissue become clearly distinct (M, N, P and Q). *Xmsx-1* is expressed in broad areas, including the fore-, mid- and hindbrains and the spinal cord, while *Xmsx-2B* is expressed in a narrower region of neural tissues. (M-O, dorsal view and anterior is toward the top; P-R, anterior view and dorsal side is toward the top). Bars in A, M and P indicate 0.5 mm.

#### I. Onitsuka et al.









has the same or similar activity, we performed mRNA injection experiments. Two dorsal blastomeres of 4-cell stage embryos were injected with various concentrations of mRNAs encoding for the full-length Xmsx-2B protein, and the injected embryos were incubated for 2 days until the sibling embryos reached the tailbud stage. The embryos injected with Xmsx-2B mRNA showed ventralized phenotypes and the average DAI was reduced in dose-dependent manner (Fig. 2 D, E, G, J and M). This activity of Xmsx-2B seemed very strong among the known ventralizing agents, and as little as 10 pg of mRNA was enough to exert a ventralizing effect (average DAI=4.3, n=154; data not shown). Injection of 1ng RNA led to a hyperventralized phenotype completely lacking the dorso-ventral axis (average DAI=0.3, n=98; Fig. 2 J, M). The phenotype induced by Xmsx-2B overexpression was very similar to that obtained by the injection of BMP-4 mRNA (average DAI=0.1, n=92; Fig. 2 I, L). In these embryos, ectodermal, mesodermal and endodermal layers were clearly distinguished, but differentiation of dorsal tissues such as the notochord, neural tube or muscle was not observed (Fig.2 O and P). Furthermore, in situ hybridization analysis indicated that in these hyperventralized embryos, α-globin mRNA was strongly expressed in the anterior region (Fig. 2 I, J) but  $\alpha$ -actin was not (Fig. 2 L, M).

Since the phenotypes of the embryos injected with *Xmsx-2B* and *BMP-4* mRNA were similar, we examined the blood cell-inducing ability of *Xmsx-2B* in the dorsal marginal zone (DMZ) explants. After injection of several concentrations of *Xmsx-2B* mRNAs (200–800 pg/embryo) into the marginal zone

Fig. 2. Ventralizing activities of Xmsx-1 and Xmsx-2B. A-F. Typical ventralizing phenotypes obtained in the mRNA injection experiments. Two dorsal blastomeres of 4-cell stage embryos were injected with designated amounts of RNAs, and the embryos were allowed to develop until stage 33/34 to determine the DAI (dorso-anterior index; Kao and Elinson, 1988). Injection of 400 pg of Xmsx-1 (B) and 25 pg of Xmsx-2B (E) led to a moderate ventralizing phenotype (average DAIs being 3.1 and 3.5, respectively). Injection of 800 pg of Xmsx-1 (C) and 50 pg of Xmsx-2B (E), which are twice the amounts of B and E, resulted in a severely ventralized phenotype (average DAIs being 2.2 and 2.3, respectively). The embryos coinjected with 400 pg of Xmsx-1 and 25 pg of Xmsx-2B (F) showed a similar degree of ventralization to that of C and E. G. Average DAI and SD of RNAinjected embryos are represented by a bar graph. The numbers of embryos injected in each groups were as follows: n=157 (uninjected), n=185 (Xmsx-1 400 pg), n=77(Xmsx-1 800 pg), n=122 (Xmsx-2B 25 pg), n=152 (Xmsx-2B 50 pg), and n=140 (Xmsx-1 400 pg+ Xmsx-2B 25 pg). Note that both Xmsx-1 and Xmsx-2B had ventralizing activities and that their effects were additive. H-P. Hyperventralized phenotype (DAI=0.3, n=98) induced by a high dose of Xmsx-2B RNA. Embryos were injected with 1ng of Xmsx-2B RNA (J, M) or 1ng of BMP-4 RNA as a control (I, L), and they were allowed to develop until the sibling embryos reached st.33/34 (H, K). Whole-mount in situ hybridization was performed to show the expressions of  $\alpha$ -globin (H, I and J) and  $\alpha$ -actin (K, L and M). The expression patterns of these markers in the embryos, which had been injected with BMP-4 and Xmsx-2B, were guite similar. N, O, and P represent histological views of embryos shown in H, I and J. Note that no axial structure was observed in BMP-4 or Xmsx-2B mRNA-injected embryos, and, instead, *α-globin*-positive cells were dramatically increased in comparison with those in uninjected embryos. Bars in A, H and N indicate 1 mm, 1 mm, and 0.5 mm respectively.

Talble 1. Incidence of secondary axis formation in injected embryos

mRNA	Number of injected embryos	Secondary axis (%)
tBR	127	56
tBR+Xmsx-1	112	0
tBR+ Xmsx-2B	117	0
VP16/msx-1	95	65
VP16/msx-1+Xmsx-1	81	0
VP16/msx-1+ Xmsx-2B	118	0

The experimental protocol is described in the legend of **Fig. 3.** Note that *Xmsx-1* and *Xmsx-2B* completely suppressed formation of the secondary axis induced by *tBR* (a dominant-negative form of the BMP-4 receptor) and *VP16/msx-1* (a dominant-negative form of Xmsx-1).

A uninjected Uni

**Fig. 3.** A-D. Injection of *Xmsx-1* or *Xmsx-2B* mRNA suppressed the secondary axis formation induced by *tBR*, a dominant-negative form of the BMP-4 receptor. Two ventral blastomeres of 4-cell stage embryos were injected with *tBR* (500 pg) RNA (B), *tBR* (500 pg) with *Xmsx-1* (1000 pg) RNA (C), and *tBR* (500 pg) with *Xmsx-2B* (200 pg) RNA (D). The embryos were allowed to develop until st.33/34, and the existence of secondary axis was determined morphologically. E-H. Injection of *Xmsx-1* or *Xmsx-2B* mRNA suppressed the secondary axis formation induced by *VP16/msx-1*, a dominant-negative form of Xmsx-1. The embryos were injected with *VP16/msx-1* (600 pg) RNA (F), *VP16/msx-1* (600 pg) with *Xmsx-2B* (200 pg) (G), or *VP16/msx-1* (600 pg) with *Xmsx-2B* (200 pg) (H). Bar in A indicates 1mm.

of dorsal blastomeres, DMZs were excised at stage10.5 and cultured until stage 39. Northern blot analysis showed that  $\alpha$ -globin mRNA was not induced in the explants even though the highest amount of mRNA was injected (data not shown). Thus, we concluded that among the target genes of the BMP-4 signal, *Xmsx-1* and *Xmsx-2B* are not sufficient for blood cell differentiation.

To investigate whether Xmsx-1 and Xmsx-2B act coop-

eratively, these RNAs were injected alone or simultaneously into two dorsal blastomeres (Fig.2 B-G). The average DAI obtained by 25 pg of *Xmsx-2B* mRNA was  $3.1\pm1.4$  (n=122), whereas that of embryos that received 400 pg of *Xmsx-1* mRNA was  $3.5\pm1.1$  (n=185). When these mRNAs were coinjected, the DAI fell to  $2.3\pm1.0$  (n=140). Since similar values of DAI were obtained in the embryos injected with 50 pg *Xmsx-2B* (DAI= $2.3\pm1.4$ ; n=152) or 800 pg *Xmsx-1* (DAI= $2.2\pm1.2$ ; n=77), we concluded that *Xmsx-1* and *Xmsx-2B* function in ventralization in an additive manner.

#### Xmsx-2B mediates BMP-4 signaling

It has been shown that the expression of *Xmsx-1* is regulated by the BMP-4 signal (Maeda *et al.*, 1997; Suzuki *et al.*, 1997). Taken together with these expression patterns of *Xmsx-2B* in the early embryo (Fig. 1), it seems that *Xmsx-2B*, like *Xmsx-1*, is regulated by the BMP-4 signal. To confirm this hypothesis, we examined whether *Xmsx-2B* could disturb the formation of the secondary axis, which was induced by *tBR* mRNA injection. Two ventral blastomeres of 4-cell stage embryos were injected with 500 pg of *tBR* mRNA alone or coinjected with 1ng of *Xmsx-1* or 200 pg of *Xmsx-2B*. As shown in Table 1 and Fig.3 B, 56 % (n=127) of *tBR*-injected embryos formed the secondary axis. In contrast, the formation of the secondary axis was completely suppressed if either *Xmsx-1* (n=112) or *Xmsx-2B* (n=117) was coinjected with *tBR* mRNA.

We also examined whether the ventralizing activity of *Xmsx-1* can be replaced by that of *Xmsx-2B*. For this purpose, we used a fusion construct of *Xmsx-1* and VP16, a dominant-negative form of *Xmsx-1*, which was shown to induce formation of the secondary axis (Takeda *et al.*, 2000). The embryos injected ventrally with *VP16/msx-1* mRNA formed an incomplete secondary axis without a head structure (65%; n=95, Fig.3 F). Simultaneous injection either of *Xmsx-1* (Fig.3 G; n=81) or *Xmsx-2B* (Fig.3 H; n=118) completely suppressed the formation of the secondary axis. These results provide us with evidence that the activity of *Xmsx-1* can be replaced by that of *Xmsx-2B* and, therefore, these two transcriptional factors share the same target molecules in the ventralizing signal.

#### DISCUSSION

### The expression of *Xmsx-2B* overlaps that of *Xmsx-1* in gastrula and neurula stages

Since Su *et al.* (1991) first reported the isolation of an *Xhox7.1*' (*Xmsx-2B*) cDNA clone in *Xenopus laevis*, there have been no reports describing the expression and function of *Xenopus msx-2*, despite the general view of importance in various vertebrate embryogenesis. Very recently, however, Gong and Kiba (1999) isolated a *msx-2* cDNA clone and emphasized its function in antero-posterior patterning of meso-derm formation. From the viewpoints of sequence similarity, expression pattern, and biological activity, *Xmsx-2B* and the cDNA isolated by Gong and Kiba (1999) are subtypes of *Xenopus msx-2*.

As expected, the expression pattern of *Xmsx-2B* was very similar to that of *Xmsx-1* (Maeda *et al.*, 1997; Suzuki *et al.*, 1997) during gastrula and neurula stages. Concomitant expressions of *msx-1* and *-2* have been reported in various tissues in mammalian species (Mackenzie *et al.*, 1992; Jowett *et al.*, 1993; Wang and Sassoon, 1995; Friedmann *et al.*, 1996; Phippard *et al.*, 1996) and in birds (Coelho *et al.*, 1991; Ganan *et al.*, 1996; Ganan *et al.*, 1998). At the gastrulation stage, the transcript was detected in the lateral and ventral marginal zone (Fig.1), suggesting that *Xmsx-2B* and *Xmsx-1* have the same or similar functions at this stage. At the tailbud stage, however, the expression pattern in neural tissues was clearly distinct. We suggest that the mechanism of expressional control in the two msx genes is conserved at least in early embryogenesis.

#### Xmsx-2B acts as a ventralizing factor

As suggested by the expression pattern of Xmsx-2B in gastrulating embryos, the forced expression of Xmsx-2B mRNA in dorsal blastomeres induced a strong ventralized phenotype (Fig. 2 J, M), and *a-globin* mRNA was strongly expressed in the embryos (Fig. 2 J). By analyzing the ventralizing activity of Xmsx-1, we noticed that, unlike BMP-4, Xmsx-1 alone lacks activity to upregulate the expression of ventral genes (Maeda et al., 1997). On the other hand, Xmsx-2B -injected embryos completely lost the body axis, while the same dose of Xmsx-1 had a weaker ventralizing effect. We predicted, therefore, that Xmsx-2B may be a cofactor of Xmsx-1 in blood cell-inducing activity. However, Northern blot analysis revealed that Xmsx-2B could not induce  $\alpha$ -globin expression in the dorsal marginal zone explants, while BMP-4 could. Although there was no difference in the translational activities of Xmsx-1 and Xmsx-2B RNAs in the reticulocyte lysate (data not shown), the stabilities of proteins in the cells were not determined. As the Xmsx-2B cDNA used in this study lacks the 3' UTR, there is a possibility that this 3' UTR might regulate the stability of Xmsx-2B mRNA. From these observations, we concluded that even though Xmsx-2B has a strong activity, Xmsx-1 and Xmsx-2B ventralize the embryo by suppressing the expression of organizer genes (Takeda et al., 2000), and in the prospective ventral cells in which the BMP-4 signal is activated, another unknown factor is required to trigger the blood cell differentiation.

#### Xmsx-1 and Xmsx-2B function in the dorso-ventral patterning of mesoderm

In the present study, we found that *Xmsx-1* and *Xmsx-2B* function in the ventralizing cascade in an additive manner (Fig. 2 A-G). In addition, the ability of *Xmsx-2B* to suppress the secondary axis formation induced by *VP16/msx-1* RNA (a dominant negative form of msx-1 protein) supports our conclusion that the ventralizing activity of msx-1 can be replaced by that of *Xmsx-2B* and that their target gene(s) might be common. So far, the molecular nature of how these msx proteins interact with each other is unclear, but previous reports have suggested that msx and dlx proteins can form homo- and

heterodimers in various combinations (Zhang et al., 1997). Since msx proteins are transcriptional repressors (Catron et al., 1995) and dlx proteins are transcriptional activators (Zhang et al., 1997), the biological activities in transcriptional regulation are mutually exclusive (Zhang et al., 1997). Other candidates of msx's partner are Xvent-1 (PV. 1) (Gawantka et al., 1995; Ault et al., 1996) and Xvent-2 (Xbr-1, Vox, Xom) (Onichtchouk et al., 1996; Papalopulu and Kintner, 1996; Ladher et al., 1996). These factors are also homeodomaincontaining proteins and possessing ventralizing activity. The similarity of expression patterns and functions between msx and vent suggests that these proteins might interact with each other, directly or indirectly, in prospective ventral cells. To elucidate the molecular mechanisms of msx's function in the ventralizing activity and the following specification of tissue distribution, isolation and analysis of the interacting molecules will be required in the future.

#### ACKNOWLEDGEMENTS

This work was partly supported by grants from The Saneyoshi Foundation, The Sumitomo Foundation and The Ministry of Education, Science and Culture of Japan (No. 09680721).

#### REFERENCES

- Ault KT, Dirksen ML, Jamrich M (1996) A novel homeobox gene *PV.1* mediates induction of ventral mesoderm in *Xenopus* embryos. Proc Natl Acad Sci USA 93: 6415–6420
- Bradley LC, Snape A, Bhatt S, Wilkinson DG (1992) The structure and expression of the *Xenopus Krox-20* gene: conserved and divergent of expression in rhombomeres and neural crest. Mech Dev 40: 73–84
- Catron KM, Zhang H, Marshall SC, Inostroza JA, Wison JM, Abate C (1995) Transcriptional repression by Msx-1 does not require homeodomain DNA-binding sites. Mol Cell Biol 15: 861–871
- Coelho CND, Sumoy L Rodgers BJ, Davidson DR, Hill RE, Upholt WB, Kosher RA (1991) Expression of the chicken homeoboxcontaining gene *Ghox-8* during embryonic chick limb development. Mech Dev 34: 143–154
- Dale L, Howes G, Price BMJ, Smith JC (1992) Bone morphogenetic protein 4: a ventralizing factor in *Xenopus* development. Development 115: 573–585
- Davidson DR, Crawley A, Hill RE, Tickle C (1991) Position-dependent expression of two related homeobox genes in developing vertebrate limbs. Nature 352: 429–431
- Fainsod A, Steinbeisser H, De Robertis EM (1994) On the function of BMP-4 in patterning the marginal zone of the *Xenopus* embryo. EMBO J 13: 5015–5025
- Farlie PG, Kerr R, Thomas P, Symes T, Minichiello J, Hearn CJ, Newgreen D (1999) A paraxial excusion zone creates patterned cranial neural crest cell outgrowth adjacent to rhombomeres 3 and 5. Development 213: 70–84
- Ferrari D, Lichtler AC, Pan ZZ, Dealy CN, Upholt WB, Kosher RA (1998) Ectopic expression of *Xhox-7.1* in posterior limb bud mesoderm impairs limb morphogenesis while inducing *BMP-4* expression, inhibiting cell proliferation, and promoting apoptosis. Dev Biol 197: 12–24
- Friedmann Y and Daniel CW (1996) Regulated expression of homeobox gene *Msx-1* and *Msx-2* in mouse mammary gland development suggests a role in hormone action and epithelialstromal interactions. Dev Biol 177: 347–355
- Ganan Y, Macias D, Duterque-Coquillaud M, Ros MA, Hurle JM (1996)

Role of TGF  $\beta$  s and BMPs as signals controlling the position of the digits and the areas of interdigital cell death in the developing chick limb autopod. Development 122: 2349–2357

- Ganan Y, Macias D, Basco RD, Merino R, Hurle J (1998) Morphological diversity of the avian foot is related with the pattern of *msx* gene expression in the developing autopod. Dev Biol 196: 33–41
- Gong SG, Kiba A (1999) The role of Xmsx-2 in the anterior-posterior patterning of the mesoderm in *Xenopus laevis*. Differentiation 65: 131–140
- Gawantka V, Delius H, Hirschfeld K, Blumenstock C, Niehrs C (1995) Antagonizing the Spemann organizer: role of the homeobox gene *Xvent-1*. EMBO J 14: 6268–6279
- Graff JM, Thies SR, Song JJ, Celeste AJ, Melton DA (1994) Studies with a *Xenopus* BMP receptor suggest that ventral mesoderminducing signals override dorsal signals in vivo. Cell 7 9: 169– 179
- Graham A, Heyman I, Lumsden A (1993) Even-numbered rhombomeres control the apoptotic elimination of neural crest cells from odd-numbered rhombomeres in the chick hindbrain. Development 119: 233–245
- Graham A, Francis-West P, Brickell P, Lumsden A (1994) The signalling molecule BMP4 mediates apoptosis in the rhombencephalic neural crest. Nature 372: 684–686
- Ishimura A, Maeda M, Takeda M, Kikkawa M, Daar IO, Maéno M (2000) Involvement of BMP-4/msx-1 and FGF pathways in neural induction in *Xenopus* embryo. Develop. Growth & Differ. 42: 307–316
- Jones CM, Lyons KM, Lapan PM, Wright CVE, Hogan BLM (1992) DVR-4 (bone morphogenetic protein-4) as a posterior-ventralizing factor in *Xenopus*. Development 115: 639–647
- Jowett AK, Vainio S, Ferguson MW, Sharpe PT, Thesleff I (1993) Epithelial-mesenchymal interactions are required for *msx1* and *msx2* gene expression in the developing murine molar tooth. Development 117: 461–470
- Kao KR and Elinson RP (1988) The entire mesodermal mantle behaves as Spemann's organizer in dorsoanterior enhanced *Xenopus laevis* embryos. Dev Biol 127: 64–77
- Ladher R, Mohun J, Smith JC, Snape A (1996) *Xom*: a *Xenopus* homeobox gene that mediates the early effects of BMP-4. Development 122: 2385–2394
- Mackenzie A, Ferguson MWJ, Sharpe PT (1992) Expression patterns of the homeobox gene, Hox-8, in the mouse embryo suggest a role in specifying tooth initiation and shape. Development 115: 403–420
- Maeda R, Kobayashi A, Sekine R, Lin JJ, Kung HF, Maéno M (1997) Xmsx-1 modifies mesodermal tissue pattern along dorsoventral axis in Xenopus laevis embryo. Development 124: 2553–2560
- Maéno M, Ong RC, Suzuki A, Ueno N, Kung HF (1994) A truncated bone morphogenetic protein-4 receptor alters the fate of ventral mesoderm to dorsal mesoderm: Roles of animal pole tissue in the development of ventral mesoderm. Proc Natl Acad Sci USA 91: 10260–10264
- Mayor R, Morgan R, Sargent MG (1995) Induction of the prospective neural crest of *Xenopus*. Development 121: 767–777
- Monachan AP, Davidson DD, Sime C, Graham E, Baldock R, Bhattacharya SS, Hill RE (1991) The *Msx*-like homeobox genes define domains in the developing vertebrate eye. Development 112: 1053–1061
- Moors JM, Wang S, Krinks M (1995) Anti-dorsalizing morphogenetic protein is a novel TGF- $\beta$  homolog expressed in the Spemann organizer. Development 121: 4293–4301
- Muneoka K and Sassoon D (1992) Molecular aspects of regeneration in developing vertebrate limbs. Dev Bio 152: 37–49

- Nieuwkoop PD and Faber J (1967) Normal Table of Xenopus laevis (Daudine). Amsterdam: North-Holland.
- Nohno T, Noji S, Koyama E, Nishikawa K, Myokai F, Saito T, Taniguchi S (1992) Differential expression of two *msh*-related homeobox genes *Chox-7* and *Chox-8* during chick limb development. Biochem Biophys Res Commun 182: 121–128
- Onichtchouk D, Gawantka V, Dosch R, Delius H, Hirschfeld K, Blumenstock C, Niehrs C (1996) The *Xvent-2* homeobox gene is part of BMP-4 signaling pathway controling dorsoventral patterning of *Xenopus* mesoderm. Development 122: 3045–3053
- Papalopulu N and Kintner C (1996) A *Xenopus* gene, *Xbr-1*, defines a novel class of homeobox genes and is expressed in the dorsal ciliary margin of the eye. Dev Biol 174: 104–114
- Phippard DJ, Weber-Hall SJ, Sharpe PT, Naylor MS, Jayatalake H, Maas R, Hill RE, Dale TC (1996) Regulation of *Msx-1, Msx-2, Bmp-2* and *Bmp-4* during foetal and postnatal mammary gland development. Development 122: 2729–2723
- Schmidt JE, Suzuki A, Ueno N, Kimelman D (1995) Localized BMP-4 mediates dorsal/ventral patterning in early *Xenopus* embryo. Dev Biol 169: 37–50
- Semenza GL, Wang GL, Kundu R (1995) DNA binding and transcriptional properties of wild-type and mutant formes of the homeodomain protein msx2. Biochem. Biophys Res Commun 209: 257–262
- Shaine DH and Zuber MX (1996) Sodium dodecyl sulfate (SDS)-based whole-mount in situ hybridization of *Xenopus laevis* embryos. Biochem Biophys Meth 31: 185–188
- Shapira E, Marom K, Yelin R, Levy A, Fainsod A (1999) A role for the homeobox gene *Xvex-1* as part of the BMP-4 ventral signaling pathway. Mech Dev 86: 99–111
- Su MW, Suzuki HR, Solursh M, Ramirez F (1991) Progressively restricted expression of a new homeobox-containing gene during *Xenopus laevis* embryogenesis. Development 111: 1179–1187
- Suzuki A, Thies RS, Yamaji N, Song JJ, Wozney JM, Murakami K, Ueno N (1994) A truncated bone morphogenetic protein receptor affects dorsal-ventral patterning in the early *Xenopus* embryo. Proc Natl Acad Sci USA 91: 10255–10259
- Suzuki A, Ueno N, Hemmati-Brivanlou A (1997) *Xenopus msx-1* mediates epidermal induction and neural inhibition by BMP4. Development 124: 3037–3044
- Takahashi K, Nuckolls GH, Tanaka O, Semba I, Takahashi I, Dashner R, Shum L, Slavkin HC (1998) Adenovirus-mediated ectopic expression of msx2 in even-numbered rhombomeres induces apoptotic elimination of cranial neural crest cells in ovo. Development 125: 1627–1635
- Takeda M, Saito Y, Sekine R, Onitsuka I, Maeda R, Maéno M (2000) *Xenopus msx-1* regulates the dorsoventral axis formation by supressing the expression of organizer genes. Comp Biochem Physiol 126: 157–168
- Zhang H, Hu G, Wang H, Sciavolino P, Iler N, Shen MM, Abate-Shen C (1997) Heterodimerization of Msx and Dlx homeoproteins results in functional antagonism. Mol Cell Biol 17: 2920–2932
- Zou H and Niswander L (1996) Requirement for BMP signaling in interdigital apoptosis and scale formation. Science 272: 738-741
- Wang Y and Sassoon D (1995) Ectoderm-mesenchyme and mesenchyme-mesenchyme interactions regulate Msx-1 expression and cellular differentiation in the murine limb bud. Dev Biol 168: 374– 382

(Received March 17, 2000 / Accepted May 20, 2000)