

# Effects of Olfactory Tract Section on the Immunohistochemical Distribution of Brain GnRH Fibers in the Female Goldfish, Carassius auratus

Authors: Kim, Myung-Hee, Kobayashi, Makito, Oka, Yoshitaka, Amano, Masafumi, Kawashima, Seiichiro, et al.

Source: Zoological Science, 18(2): 241-248

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zsj.18.241

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Effects of Olfactory Tract Section on the Immunohistochemical Distribution of Brain GnRH Fibers in the Female Goldfish, *Carassius auratus*

Myung-Hee Kim<sup>1</sup>, Makito Kobayashi<sup>1\*</sup>, Yoshitaka Oka<sup>2</sup>, Masafumi Amano<sup>1†</sup>, Seiichiro Kawashima<sup>3</sup> and Katsumi Aida<sup>1</sup>

 <sup>1</sup>Department of Aquatic Bioscience, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo-ku, Tokyo 113-8657,
<sup>2</sup>Misaki Marine Biological Station, Graduate School of Science, The University of Tokyo, Misaki, Miura, Kanagawa 238-0225,
<sup>3</sup>Research Laboratory, Zenyaku Kogyo Co., Ltd., Nerima-ku, Tokyo 178-0062, Japan

**ABSTRACT**—The organization of the gonadotropin-releasing hormone (GnRH) system originating from the terminal nerve (TN) neuron in the olfactory bulb was examined immunohistochemically by tracing the changes in the distribution of salmon type (sGnRH) and chicken-II type (cGnRH-II) GnRH fibers after olfactory tract section (OTX) in the female goldfish. Following OTX, which blocks the axonal transport of GnRH from the TN to other brain regions, the density of sGnRH-ir fibers in various brain regions decreased from 7 days to 28 days, with the exception of rather restricted areas surrounding cell bodies in the ventral telencephalon, preoptic area, and hypothalamus. The density of cGnRH-II-ir fibers in several brain areas, neither type of GnRH fibers showed marked changes in the pituitary gland during the experiment. These results indicate that the TN GnRH system project fibers to wide brain areas and that most of sGnRH and some of the cGnRH-II-ir fibers in the brain of goldfish are TN origin. Furthermore, the GnRH system that project fibers to the pituitary does not primarily originate from the TN-GnRH system.

### INTRODUCTION

Gonadotropin-releasing hormone (GnRH) is a decapeptide which stimulates the release of gonadotropins (GTH) from the pituitary gland. To date, twelve types of GnRH molecules have been isolated, and their primary structures have been determined (See reviews by Okuzawa and Kobayashi, 1999; Parhar, 1999). Recent studies have shown that in teleosts, more than one type of GnRH molecule exists even within the same species, and that GnRH neurons are found not only in the hypothalamic region but also in the extrahypothalamic regions of the brain (Amano *et al.*, 1997; Okuzawa and Kobayashi, 1999; Parhar, 1999; Yamamoto *et al.*, 1995). These results led us to consider whether all of the brain GnRHs function in the regulation of GTH release, and if not, which population and which type of GnRH regulates GTH release in teleosts.

In the goldfish, Carassius auratus, two types of GnRH,

\* Corresponding author: Tel. +81-3-5841-5286; FAX. +81-3-5841-5288.

E-mail: amakito@mail.ecc.u-tokyo.ac.jp <sup>†</sup> Present address: School of Fisheries Sciences, Kitasato University, Sanriku, Iwate 022-0101, Japan salmon (sGnRH) and chicken-II (cGnRH-II) have been identified in the brain and pituitary gland at peptide and mRNA levels by chromatography combined with radioimmunoassays (RIA) and molecular cloning, respectively (Yu et al., 1988; Kobayashi et al., 1992; Lin and Peter, 1996, 1997; Suetake et al., 2000). Our previous immunohistochemical study in the goldfish brain revealed that sGnRH-immunoreactive (ir) cell bodies are localized in the terminal nerve (TN) ganglion (TN GnRH system), the ventral telencephalon, the preoptic area (POA), and the hypothalamus (POA GnRH system) (Kim et al., 1995a). cGnRH-II-ir cell bodies occur in the same areas as those of sGnRH and also in the midbrain tegmentum (midbrain GnRH system). sGnRH and cGnRH-II fibers are widely distributed in the brain and the pituitary. The wide distribution of GnRH-ir fibers in the brain suggested that, in the goldfish, sGnRH and cGnRH-II regulate not only gonadotropin release from the pituitary gland, but also function as neuromodulators in the brain (Oka, 1997). In fact, changes in brain GnRH content does not always reflect levels of gonadal maturation in goldfish (Yu et al., 1987). Since the olfactory bulbs of the goldfish are of the pedunculated type with elongated olfactory tracts and the TN cell bodies are localized in the rostral part of the olfactory bulb, the axonal transport of GnRH from the TN to

other brain regions can be easily blocked by sectioning the olfactory tracts. Our studies using RIA showed that olfactory tract section (OTX) causes remarkable decreases in brain sGnRH content without inhibiting gonadal maturation (Kim *et al.*, 1995b,c; Kobayashi *et al.*, 1992,1994,1997), suggesting that GTH secretion in the goldfish is primarily regulated by GnRH systems other than the TN-GnRH system. However, it is difficult to confirm by RIA whether the origin of the decreased sGnRH is solely the TN, or alternatively, whether the TN influenced GnRH production in other GnRH neurons.

In the present study, we examined the chronological changes in the distribution of sGnRH- and cGnRH-II-immunoreactive fibers after OTX in order to elucidate the anatomical organization of the TN GnRH system in the goldfish.

#### MATERIALS AND METHODS

#### Experiment 1

Sexually regressed female goldfish weighing 40 to 70 g (gonadosomatic index 0.5-1.5) were obtained from a commercial source in August and kept in stock tanks at  $20^{\circ}$ C.

Fish were olfactory tract-sectioned (N=35) or sham-operated (N=35) as described in the previous reports (Kim *et al.*, 1995b; Kobayashi *et al.*, 1992, 1994). For the olfactory tract section (OTX), a four-sided flap was cut in the skull cap using a disc saw. The first cut was made aligned with the center of the iris. The second cut, parallel to the first, was made just behind the nares. The third and fourth cuts, parallel to the longitudinal axis of the fish, was made to connect the first and second cuts. After removal of a square flap detached from the skull cap, fat and cranial fluid were gently removed. The olfactory tracts were bilaterally sectioned at two places with Wecker's scissors, and the resultant sections were removed to prevent possible regeneration (OTX fish). The cavity resulting from the operation was filled with gelatin sponge (Spongel, Yamauchi Pharmaceutical Co., Tokyo, Japan). Sham operations were performed in the same manner without cutting the olfactory tracts (sham-operated fish).

After the operation, fish were kept at 20°C and fed commercial trout pellets once a day. Fish were sampled at 3, 7, 14, 21, and 28 days after OTX for immunohistochemical analysis. At each sampling time, two or three fish were sampled. Intact fish were also sampled as initial controls.

#### Experiment 2

Sexually immature female goldfish weighing 8.0 to 22 g (gonadosomatic index 1.5-4.2) were obtained from a commercial source in September and kept at 20°C. Fish were olfactory tract-sectioned (N=20) or sham-operated (N=20). After the operation, fish were kept at 20°C and fed commercial trout pellets once a day. Four fish were sampled before and 14 and 28 days after the operation to trace changes in the density of GnRH-ir fibers. In this experiment, changes in the density of GnRH fibers were quantified after the operation. Representative parasagittal sections containing the whole brain regions except the olfactory bulb and the tract were selected from each of the sGnRH and cGnRH-II staining. The density of GnRH-ir fibers were calculated as total length of fiber (mm) / brain areas (mm<sup>2</sup>) in representative sections (48 µm thickness) using KIMA image analyzing system (Ecosolution Co., Seoul, Korea). The difference of the means among groups was analyzed by ANOVA and Duncan's test with raw or log-transformed data.

#### **Experiment 3**

Sexually maturing or mature female goldfish weighing 20 to 30 g (gonadosomatic index 5.0-8.5) were obtained from a commercial source in May and kept at  $18^{\circ}$ C, which is favorable for maintianing

the maturity of the fish. Fish were olfactory tract-sectioned (N=23) or sham-operated (N=23). After the operation, fish were kept at 18°C and fed commercial trout pellets once a day. Fish were sampled for immunohistochemical analysis 28 days after the operation.

#### Immunohistochemistry

Fish were deeply anesthetized by immersion in 0.02% tricane methanesulfonate. They were then perfused through the heart with 0.75% NaCl, followed by a fixative that had been freshly prepared by mixing 40 ml of 10% paraformaldehyde, 50 ml of 0.2 M phosphate buffer (pH 7.4), 7 ml of saturated picric acid, and 3 ml of distilled water. The head was post-fixed in the same fixative for 24 hr at 4°C. Next, the brain and the pituitary were dissected out of the head and fixed again for 24 hr at 4°C. Tissues were embedded in 15% gelatin dissolved in a 0.1 M phosphate buffer (pH 7.4) and were quickly frozen by immersing them in powdered dry ice. Forty-eight micrometer sagittal and frontal sections were cut on a cryostat, and the subsequent procedures were done on floating sections. Alternate sections were used for sGnRH and cGnRH-II staining. We used antisera raised against sGnRH (lot No. 2) and cGnRH-II (lot R-A) (Okuzawa et al., 1990; Hasegawa et al., 1986). These antisera were diluted 10,000and 20,000-fold respectively, with 0.1 M phosphate buffer (pH 7.4) containing 0.75% NaCl and 0.3% Triton X-100 (PBST). For immunohistochemical reactions, we used the streptavidin method with a commercial kit (Histofine; Nichirei Co., Tokyo, Japan). Immunoreactivity of each GnRH neuron was judged to be positive when the labeling was denser than that of the background. Meanwhile, there was no difference of immunoreactivity in each individual. Representative parasagittal sections that are considered to be the best areas to show the localization of GnRH neurons (Kim et al., 1995a), were selected for the camera lucida drawings (Figs. 1, 2). Localization of cell bodies shown in Figs. 1, 2 were determined from the observation of several sections. The specificity of anti-sGnRH and cGnRH-II have been demonstrated by RIA (Kobayashi et al., 1992; Okuzawa et al., 1990) and by immunocytochemistry (Amano et al., 1991; Kim et al., 1995a). To test the specificity of the immunoreactions, the diluted sGnRH and cGnRH-II antisera were each incubated overnight at 4°C with an excess amount of synthetic sGnRH and cGnRH-II (1 µg GnRH in 1 ml of diluted antiserum) respectively, or conversely, sGnRH and cGnRH-II antisera with synthetic cGnRH-II and sGnRH, respectively. The subsequent histological procedures were performed in the same manner as in the experimental sections. No immunoreactive fibers were observed when the antisera were preincubated with respective GnRHs. On the contrary, immunoreactive fibers were observed when the antisera were preincubated with different GnRHs. When the sGnRH and cGnRH-II antisera were incubated with seabream GnRH (1 µg GnRH in 1ml of diluted antiserum) (Powell et al., 1994), they were not absorbed, and immunoreactive fibers were observed with both the sGnRH and cGnRH-II antisera. Synthetic seabream GnRH was obtained from Fujiya Bioscience Co. (Hadano, Kanagawa, Japan). Moreover, we tested the specificity of the antisera by immunoreaction on nitrocellulose membranes. Both synthetic sGnRH and cGnRH-II of different amount, 1 ng, 100 ng, 1 µg, 10 µg, 20 µg, and 30  $\mu g$  in 100  $\mu$  l of 0.1 M PBST were spotted on nitrocellulose membrane. The membrane was treated in the same fixative as for the brain tissues. The subsequent immunoreaction procedures were done in the same manner as in the experimental sections. When the antisera were incubated with heterologous antigens of 10  $\mu$ g or more, only faint staining reactions were observed, while strong reactions were observed between the antisera and the homologous antigens at these doses.

We used the terminology of brain structures or subdivisions according to Northcutt and Davis (1983) for the telencephalon, Braford and Northcutt (1983) for the diencephalon and pretectum, and Nieuwenhuys and Pouwels (1983) for the brainstem.

## RESULTS

#### **Experiment 1**

The distribution of sGnRH-ir fibers 3 days after OTX showed no changes in various brain regions and in the pituitary gland in comparison with those of the initial controls (Fig. 1). At Day 7, sGnRH-ir fibers in various brain areas started to diminish. In particular, the decrease in the density of ir fibers in the telencephalon, the optic tectum, and the thalamus was more remarkable than in the hypothalamus, cerebellum, and medulla oblongata. At Day 14, further decrease of sGnRH-ir fibers was observed in the brain areas except in the ventral telencephalon, the preoptic area, and the hypothalamus where ir cell bodies were located. Thereafter, there were no further changes in the distribution of ir fibers in all brain areas.

cGnRH-II-ir fibers decreased mainly in the telencephalon from Day 7 to Day 14 in comparison with that of the initial control (Fig. 2). Thereafter, there were no further changes in the distribution of cGnRH-II-ir fibers in all brain areas.

On the other hand, both types of GnRH fibers showed no

marked changes in the pituitary gland during the experiment (Figs. 1, 2). The ovaries of the OTX and sham-operated fish remained regressed.

#### **Experiment 2**

Distribution of GnRH fibers in initial control fish, OTX fish, and sham-operated fish was similar to that in equivalent fish in Experiment 1 (data not shown). When the density of GnRH fibers was calculated by summing the length of all GnRH-ir fibers observed in the representative parasagittal brain section (olfactory bulb and tract are not included) and normalized by the brain area, the density of sGnRH fibers showed a significant decrease after OTX compared to initial control fish (day 14, p<0.01; day 28, p<0.01) and sham-operated fish of the same sampling days (day 14, p<0.01; day 28, p<0.01) (Fig. 3). There were no significant differences in the density of sGnRH-ir fibers between initial control fish and sham-operated fish.

The density of cGnRH-II-ir fibers in the brain showed a significant decrease after OTX compared to initial control fish



**Fig. 1.** Semidiagramatic illustrations of salmon gonadotropin-releasing hormone-immunoreactive (sGnRH-ir) cell bodies (solid circles) and fibers (lines) in a parasagittal section of the immature goldfish brain (Exp. 1). Shaded areas (dense) and lines (scarce) in the pituitary indicate sGnRH-ir fibers. D0, initial; D3, 3 days after olfactory tract section (OTX); D7, 7 days after OTX; D14, 14 days after OTX; D21, 21 days after OTX; D28, 28 days after OTX. Arrows indicate location of OTX. C, cerebellum; M, medulla oblongata; OB, olfactory bulb; OLF, olfactory nerve; ON, optic nerve; OT, optic tectum; PIT, pituitary; T, telencephalon. Scale bar, 1 mm.



**Fig. 2.** Semidiagramatic illustrations of chicken gonadotropin-releasing hormone-II-immunoreactive (cGnRH-II-ir) cell bodies (solid circles) and fibers (lines) in a parasagittal section of the immature goldfish brain (Exp. 1). Shaded areas (dense) and lines (scarce) in the pituitary indicate cGnRH-II-ir fibers. D0, initial; D3, 3 days after olfactory tract section (OTX); D7, 7 days after OTX; D14, 14 days after OTX; D21, 21 days after OTX; D28, 28 days after OTX. Arrows indicate location of OTX. Abbreviations for brain areas are as in Fig. 1. Scale bar, 1 mm.



Fig. 3. Changes in the density of sGnRH-ir fibers in the brain after olfactory tract section in goldfish. The density was calculated by sum-

(day 14, p<0.001; day 28, p<0.001) and sham-operated fish of the same sampling days (day 14, p<0.05; day 28, p<0.001) (Fig.4). Sham-operated fish showed a decrease in cGnRH-II-ir fibers at day 14 after operation compared to that in initial control fish (p<0.01).

## **Experiment 3**

Distribution of GnRH fibers in OTX fish, and sham-operated fish was similar to that in equivalent fish in Experiments 1 and 2. Most of sGnRH-ir fibers and some cGnRH-II-ir fibers disappeared in various brain areas with the exception of those in the pituitary gland, the ventral telencephalon, the preoptic area, and the hypothalamus where immunoreactive cell bod-

ming the length of sGnRH-ir fibers observed in the representative parasagittal brain section (the olfactory bulb and tract are not included) and normalized by brain area. IC, Initial control; Sham, sham-operated fish; OTX, olfactory tract-sectioned fish. D14, 14 days after operation; D28, 28 days after operation. Each column represents the mean and SEM (N=4). , Significant difference compared to initial control fish (p<0.01). \*\*, Significant difference compared to sham-operated fish (p<0.01).



**Fig. 4.** Changes in the density of cGnRH-II-ir fibers in the brain after olfactory tract section in goldfish. The density was calculated by summing the length of cGnRH-II-ir fibers observed in the representative parasagittal brain section (the olfactory bulb and tract are not included) and normalized by brain area. IC, Initial control; Sham, sham-operated fish; OTX, olfactory tract-sectioned fish. D14, 14 days after operation; D28, 28 days after operation. Each column represents the mean and SEM (N=4). , Significant difference compared to initial control fish. \*, Significant difference compared to sham-operated fish. Levels of significance: one symbol, p<0.05; two symbols, p<0.01; three symbols, p<0.001.

ies were located (Figs. 5–16). Despite the disappearance of most of the sGnRH-ir fibers and some of the cGnRH-II-ir fibers, the fish maintained mature ovaries exhibiting vitellogenic oocytes and showed no difference from those of sham-oper-ated fish.

## DISCUSSION

The localization of two types of GnRH cell bodies and fibers in initial controls and sham-operated goldfish showed a pattern similar to that in goldfish reported in our previous study (Kim et al., 1995a). Both sGnRH- and cGnRH-II-ir cell bodies were localized in the TN (TN GnRH system), the ventral telencephalon, the preoptic area, and the anterior hypothalamus (POA GnRH system). In addition, cGnRH-II-ir but sGnRHnegative cell bodies were also observed in the midbrain tegmentum (midbrain GnRH system). Both sGnRH- and cGnRH-II-ir fibers were detected in diverse brain areas and in the pituitary. In OTX fish, there was a marked decrease in the GnRH-ir fibers in the brain. The density of sGnRH- and cGnRH-II-ir fibers started to decrease 7 days after OTX, and then Day 14, the density of sGnRH-ir fibers continued to decrease in various brain regions except for the rather restricted areas surrounding cell bodies in the ventral telencephalon, the preoptic area, and the hypothalamus. At 21 and 28 days, the density of fibers immunoreactive for both molecular types showed no further decrease. The results of present immunohistochemical study support those of our RIA study which showed that brain sGnRH contents starts to decrease at 7 days after OTX and reach stable levels at 14 days (Kim *et al.*, 1995b) as well as immunohistochemical studies which showed the disappearance of TN GnRH-derived fibers in the dwarf gourami, *Colisa Ialia* two weeks after TN lesion (Yamamoto *et al.*, 1995;1997).

Chicken GnRH-II-ir fibers decreased in the telencephalon after OTX. However, in the RIA study (Kim *et al.*, 1995b), no decrease in the content of cGnRH-II was observed in all brain areas after OTX. This difference between RIA and immunocytochemistry studies is most likely due to the difference in sensitivities between the two different methods. Another possibility is that the GnRH content was measured by RIA using the whole telencephalon, whereas the number of GnRH-ir fibers was counted only in the restricted areas including mainly parasagittal sections.

The present immunohistochemical study indicates that the TN-GnRH neuronal system projects fibers to wide brain areas and that most of sGnRH and some of cGnRH-II in the brain originate from the TN GnRH system. Therefore, the decrease in the brain GnRH content after OTX as shown by RIA studies (Kim *et al.*, 1995b,c; Kobayashi *et al.*, 1992, 1994) is mostly attributable to the blockade of axonal transport of GnRH from the TN GnRH system rather than to the decrease in GnRH production in other GnRH systems. The projection of GnRH fibers to wide brain areas has also been reported in the dwarf gourami by TN lesion study (Yamamoto *et al.*, 1995, 1997).

Despite the decrease in GnRH-ir fibers after OTX in the various brain regions, GnRH-ir fibers and cell bodies of both molecular types in the ventral telencephalon, the preoptic area, the hypothalamus, and the midbrain tegmentum remained almost unchanged. Therefore, it is considered that remaining GnRH-ir fibers originate from the GnRH-ir cell bodies in these brain regions. Brain GnRH in OTX fish measured by RIA is considered to be GnRH produced in these remaining GnRH neurons (Kim et al., 1995a,b,c; Kobayashi et al., 1992, 1994). GnRH-ir fibers of both types in the pituitary showed no marked changes in immunohistochemical sections, although we did not conduct quantitative analysis because of the difficulty in tracing individual fibers in the pituitary. Meanwhile, RIA studies showed no marked changes in the pituitary GnRH contents after OTX, suggesting that GnRH fibers in the pituitary do not primarily originate from the TN system (Kim et al., 1995b; Kobayashi et al., 1992, 1994).

Our previous studies showed that OTX fish underwent normal gonadal development as did sham-operated fish (Kim *et al.*, 1995c; Kobayashi *et al.*, 1992, 1994). Taken together, the results of the present study strongly indicates that GnRH of TN origin is not a prerequisite for gonadal development in the goldfish. Recent studies in teleost GnRH neuronal systems showed that GnRH fibers in the pituitary gland mainly originate from the POA GnRH system and suggested that these GnRH neurons play hypophysiotropic function (Kah *et* 



**Fig. 5.** Sagittal section through the telencephalon which shows sGnRH-ir cell bodies and fibers in sham-operated goldfish 28 days after olfactory tract section. Arrowhead indicates a sGnRH-ir cell body of POA GnRH system. Scale bar, 400 μm.

Fig. 6. Sagittal section through the telencephalon similar to Fig. 5 in olfactory tract-sectioned goldfish 28 days after olfactory tract section. Arrowhead indicates a sGnRH-ir cell body of POA GnRH system. Scale bar, 400  $\mu$ m.

**Fig. 7.** Sagittal section through the preoptic area which shows sGnRH-ir cell bodies and fibers in sham-operated goldfish 28 days after olfactory tract section. Arrowhead indicates a sGnRH-ir cell body of POA GnRH system. Scale bar, 200 μm.

**Fig. 8.** Sagittal section through the preoptic area similar to Fig. 7 in olfactory tract-sectioned goldfish 28 days after olfactory tract section. Arrowhead indicates a sGnRH-ir cell body of POA GnRH system. Scale bar, 200 μm.

**Fig. 9.** Sagittal section through the hypothalamus which shows sGnRH-ir cell bodies and fibers in sham-operated goldfish 28 days after olfactory tract section. Arrowhead indicates a sGnRH-ir cell body of POA GnRH system. Scale bar, 200 μm.

**Fig. 10.** Sagittal section through the hypothalamus similar to Fig. 9 in olfactory tract-sectioned goldfish 28 days after olfactory tract section. Arrowhead indicates a sGnRH-ir cell body of POA GnRH system. Scale bar, 200 μm.

*al.*, 1986; Yamamoto *et al.*, 1995, 1998; Okuzawa and Kobayashi, 1999; Parhar, 1999). sGnRH and cGnRH-II neurons comprising the POA GnRH system is presumed to be the major GnRH population regulating gonadotropin release in the pituitary of the goldfish.

Physiological functions of the TN-GnRH and the midbrain

GnRH systems in the goldfish remain unknown. Neuromodulatory function is proposed as one of the most probable physiological functions of TN GnRH in teleosts (Oka, 1997), and the involvement of midbrain GnRH in sexual behavior is suggested in the musk shrew, *Suncus murinus* (Dellovade *et al.*, 1995) and in the regulation of the caudal neurosecretory



**Fig. 11.** Sagittal section through the pituitary (proximal pars distalis) in sham-operated goldfish showing sGnRH-ir fibers 28 days after olfactory tract section. Scale bar, 100 μm.

Fig. 12. Sagittal section through the pituitary similar to Fig.11 in olfactory tract-sectioned goldfish showing sGnRH-ir fibers 28 days after olfactory tract section. Scale bar, 100 μm.

**Fig. 13.** Sagittal section through the preoptic area which shows cGnRH-II cell bodies and fibers in sham-operated goldfish 28 days after olfactory tract section. Arrowhead indicates a cGnRH-II-ir cell body of POA GnRH system. Scale bar: 200 μm.

**Fig. 14.** Sagittal section through the preoptic area similar to Fig. 13 in olfactory tract-sectioned goldfish 28 days after olfactory tract section. Arrowhead indicates a cGnRH-II-ir cell body of POA GnRH system. Scale bar, 200 µm.

Fig. 15. Sagittal section through the pituitary (proximal pars distalis) in sham-operated goldfish showing cGnRH-II-ir fibers 28 days after olfactory tract section. Scale bar, 100 μm.

Fig. 16. Sagittal section through the pituitary similar to Fig. 15 in olfactory tract-sectioned goldfish showing cGnRH-II-ir fibers 28 days after olfactory tract section. Scale bar, 100 μm.

neurons in poecilid fishes (Miller and Kriebel, 1986). Recently, a stimulatory effect of GnRH on prostaglandin-induced female spawning behavior was shown in the goldfish by GnRH administration (Volkoff and Peter, 1999). GnRH from the TN GnRH and the midbrain GnRH neuronal systems may be involved in the modulation of spawning behavior in the goldfish (Stacey and Kyle, 1983).

In summary, the present immunohistochemical study together with our previous studies (Kim *et al.*, 1995a, b, c; Kobayashi *et al.*, 1992, 1994) indicate that in the goldfish the TN GnRH system projects fibers to wide brain areas and that most of sGnRH and a part of the cGnRH-II in the brain are TN origin. However, the TN GnRH system is not a prerequisite for gonadal maturation in the goldfish, and the GnRH system that projects fibers to the pituitary does not primarily originate from the TN GnRH system.

#### ACKNOWLEDGMENTS

We thank Dr. Matthew S. Grober of Arizona State University and Dr. Ishwar S. Parhar of Nippon Medical School for critical reading of the manuscript. This study was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture of Japan.

## REFERENCES

- Amano M, Oka Y, Aida K, Okumoto N, Kawashima S, Hasegawa Y (1991) Immunocytochemical demonstration of salmon GnRH and chicken GnRH-II in the brain of masu salmon, *Oncorhynchus masou*. J Comp Neurol 314: 587–597
- Amano M, Urano, A, Aida K (1997) Distribution and function of gonadotropin-releasing hormone (GnRH) in the teleost brain. Zool Sci 14: 1–11
- Braford MR, Northcutt RG (1983) Organization of the diencephalon and pretectum of the ray-finned fishes. In "Fish Neurobiology, Vol II" Ed by RE Davis and RG Northcutt, Univ. of Michigan Press, Ann Arbor, pp 117–164
- Dellovade TL, Ottinger MA, Rissman EF (1995) Mating alters gonadotropin-releasing hormone cell number and content. Endocrinology 136: 1648–1657
- Hasegawa Y, Miyamoto K, Igarashi M (1986) Regional dissociation of chicken LHRH and GnRH-II in chicken and Japanese quail brains. In "Proceedings of the 10th Annual Meeting of the Japan Soc for Comp Endocrinol" Ed by H Kobayashi and M Wada, Japan Soc Comp Endocrinol, Tokyo, p 7
- Kah O, Breton B, Dulka JG, Nunez-Rodriguez J, Peter RE, Corrigan A, Rivier JE, Vale WW (1986) A reinvestigation of the Gn-RH (gonadotropin-releasing hormone systems in the goldfish brain using antibodies to salmon Gn-RH. Cell Tissue Res 221: 1–13
- Kim MH, Oka Y, Amano M, Kobayashi M, Okuzawa K, Hasegawa Y, Kawashima S, Suzuki Y, Aida K (1995a) Immunocytochemical localization of sGnRH and cGnRH-II in the brain of goldfish, *Carassius auratus*. J Comp Neurol 355: 2–12
- Kim MH, Kobayashi M, Aida K (1995b) Changes in brain GnRH after olfactory tract section in female goldfish. Fisheries Sci 61: 614– 617
- Kim MH, Kobayashi M, Aida K (1995c) Seasonal changes in levels of sGnRH and cGnRH-II in discrete areas of the brain of the female goldfish, *Carassius auratus*. Proc Japan Soc for Comp Endocrinol 10: 53
- Kobayashi M, Amano M, Hasegawa Y, Aida K (1992) Effects of olfactory tract section on brain GnRH distribution, plasma gonadotropin levels, and gonadal stage in goldfish. Zool Sci 9: 765–773
- Kobayashi M, Amano M, Kim MH, Furukawa K, Hasegawa Y, Aida K (1994) Gonadotropin-releasing hormones of terminal nerve origin are not essential to ovarian development and ovulation in goldfish. Gen Comp Endocrinol 95: 192–200
- Kobayashi M, Amano M, Kim MH, Yoshiura Y, Sohn YC, Suetake H, Aida K (1997) Gonadotropin-releasing hormone and gonadotropin in goldfish and masu salmon. Fish Physiol Biochem 17: 1–8
- Lin XW, Peter RE (1996) Expression of salmon gonadotropin-releasing hormone (GnRH) and chicken GnRH-II precursor messenger ribonucleic acids in the brain and ovary of goldfish. Gen Comp Endocrinol 101: 282–296

- Lin XW, Peter RE (1997) Cloning and expression pattern of a second [His<sup>5</sup>Trp<sup>7</sup>Tyr<sup>8</sup>] gonadotropin-releasing hormone (chicken GnRH-II) mRNA in the goldfish: Evidence for two distinct genes. Gen Comp Endocrinol 107: 262–272
- Miller KE, Kriebel RM (1986) Peptidergic innervation of caudal neurosecretory neurons. Gen Comp Endocrinol 64: 396–400
- Nieuwenhuys R, Pouwels E (1983) The brain stem of actinopterygian fishes. In "Fish Neurobiology, Vol I" Ed by RE Davis and RG Northcutt, Univ. of Michigan Press, Ann Arbor, pp 25–87
- Northcutt RG and Davis RE (1983) Telencephalic organization in rayfinned fishes. In "Fish Neurobiology, Vol. II" Ed by RE Davis and RG Northcutt, Univ. of Michigan Press, Ann Arbor, pp 203–235
- Oka Y (1997) GnRH neuronal system of fish brain as a model system for the study of peptidergic neuromodulation. In "GnRH Neurons, Gene to Behavior" Ed by IS Parhar and Y Sakuma, Brain Shuppan Publishers, Tokyo, pp 245–276
- Okuzawa K, Amano M, Kobayashi M, Aida K, Hanyu I, Hasegawa Y, Miyamoto K (1990) Differences in salmon GnRH and chicken GnRH-II contents in discrete brain areas of male and female rainbow trout according to age and stage of maturity. Gen Comp Endocrinol 80: 116–126
- Okuzawa K, Kobayashi M (1999) Gonadotropin-releasing hormone neuronal systems in the teleostean brain and functional significance. In "Neuronal regulation in the vertebrate endocrine system" Ed by PD Prasada Rao and RE Peter, Kluwer Academic/ Plenum Publishers, New York, pp 85–100
- Parhar IS (1999) Multiple gonadotropin-releasing hormone neuronal systems in vertebrates. Korean J Biol 3: 1–7
- Powell JFF, Zohar Y, Elizur A, Park M, Fischer WH, Craig AG, Rivier JE, Lovejoy DA, Sherwood NM (1994) Three forms of gonadotropin-releasing hormone characterized from brain of one species. Proc Natl Acad Sci USA 91: 12081–12085
- Stacey NE, Kyle AL (1983) Effects of olfactory tracts lesions on sexual and feeding behavior in the goldfish. Physiol Behav 30: 621–628
- Suetake H, Yoshiura Y, Kikuchi K, Gen K, Ashihara M, Kobayashi M, Aida K (2000) Two salmon gonadotropin-releasing hormone genes and their differential expressions in the goldfish *Carassius auratus*. Fisheries Sci 66: 49–57
- Volkoff H, Peter RE (1999) Actions of two forms of gonadotropin releasing hormone and a GnRH antagonist on spawning behavior of the goldfish *Carassius auratus*. Gen Comp Endocrinol 116: 347–355
- Yamamoto N, Oka Y, Amano M, Aida K, Hasegawa Y, Kawashima S (1995) Multiple GnRH neuronal system in the dwarf gourami: Immunohistochemistry and radioimmunoassay. J Comp Neurol 355: 354–368
- Yamamoto N, Oka Y, Kawashima S (1997) Lesions of gonadotropinreleasing hormone-immunoreactive terminal nerve cells: effects on the reproductive behavior of male dwarf gourami. Neuroendocrinol 65: 403–412
- Yamamoto N, Parhar IS, Sawai N, Oka Y, Ito H (1998) Preoptic gonadotropin-releasing hormone (GnRH) neurons innervate the pituitary in teleosts. Neurosci Res 31: 31–38
- Yu KL, Nahorniak CS, Peter RE, Corrigan A, Rivier JE, Vale WW (1987) Brain distribution of radioimmunoassayable gonadotropin-releasing hormone in female goldfish: Seasonal variation and periovulatory changes. Gen Comp Endocrinol 67: 234–246
- Yu KL, Sherwood NM, Peter RE (1988) Differential distribution of two molecular forms of gonadotropin-releasing hormone in discrete brain areas of goldfish (*Carassius auratus*). Peptide 9: 625–630

(Received June 16, 2000 / Accepted November 6, 2000)