



## **Development of SSR Markers by 454 Sequencing in the Endemic Species *Gentianella praecox* subsp. *bohemica* (Gentianaceae)**

Authors: Šurinová, Mária, Brabec, Jiří, and Münzbergová, Zuzana

Source: Applications in Plant Sciences, 5(1)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1600114>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# DEVELOPMENT OF SSR MARKERS BY 454 SEQUENCING IN THE ENDEMIC SPECIES *GENTIANELLA PRAECOX* SUBSP. *BOHEMICA* (GENTIANACEAE)<sup>1</sup>

MÁRIA ŠURINOVÁ<sup>2,3,5</sup>, JIŘÍ BRABEC<sup>4</sup>, AND ZUZANA MÜNZBERGOVÁ<sup>2,3</sup>

<sup>2</sup>Institute of Botany of the Czech Academy of Sciences, Zámek 1, 252 43 Průhonice, Czech Republic; <sup>3</sup>Department of Botany, Faculty of Science, Charles University in Prague, Benátská 2, 128 01 Prague, Czech Republic; and <sup>4</sup>Muzeum Cheb, Krále Jiřího z Poděbrad 493/4, 350 11 Cheb, Czech Republic

- **Premise of the study:** Polymorphic microsatellite loci were developed and used to genotype individuals of *Gentianella praecox* subsp. *bohemica* (Gentianaceae), a highly protected taxon in Europe, to study the genetic structure of the remaining populations.
- **Methods and Results:** Thirty-eight primer pairs were successfully amplified; of these, 12 polymorphic microsatellite loci were developed using a 454 sequencing approach and used to genotype 180 individuals of *G. praecox* subsp. *bohemica* from six populations. Allelic richness ranged between one and nine alleles per locus. We detected a high frequency of polyploid individuals (77.8%). The highest average percentage of heterozygous genotypes was identified for samples from the Hroby population (75.5%). All loci can also be amplified in the congeneric species *G. praecox* subsp. *praecox*, *G. amarella* subsp. *amarella*, and *G. obtusifolia* subsp. *sturmiana*.
- **Conclusions:** These markers will provide knowledge on patterns of gene flow and population genetic structure, which is necessary for current protection actions and for effective conservation of this species in the future.

**Key words:** genotyping; Gentianaceae; *Gentianella praecox* subsp. *bohemica*; microsatellites; polyploidy.

*Gentianella praecox* (A. Kern. & Jos. Kern.) Dostál ex E. Mayer subsp. *bohemica* (Skalický) Holub (IUCN: e.T161825A5500524) is a strictly biennial herb endemic to the Bohemian Massif, with most populations occurring in the Czech Republic but extending to Bavaria (Germany), Upper and Lower Austria, and Poland. *Gentianella* Moench (Gentianaceae) is a highly diverse and taxonomically complicated genus due to seasonal dimorphism, introgression, and hybridization between closely related species (Winfield et al., 2003; Greimler and Jang, 2007; Plenk et al., 2016). It is expected that *G. praecox* subsp. *bohemica* is tetraploid (Oberdorfer, 1983), but cytotype distribution is unknown. It occurs in seminatural, nutrient-poor grasslands. Strong reduction of population size was recorded during the 20th century, probably due to land-use intensification or abandonment of traditional land use, which led to the disintegration of large habitats and fragmentation of original populations. *Gentianella praecox* subsp. *bohemica* is highly protected in Europe (Annexes II and IV of the Habitats Directive; Council of

the European Community, 1992). By using amplified fragment length polymorphism, Königer et al. (2012) studied the genetic structure of 11 *G. praecox* subsp. *bohemica* populations, but this taxon is known from 99 localities (Brabec, 2010). For effective protection of this subspecies, it is necessary to identify populations with high genetic diversity so these populations can be prioritized for protection. Moreover, knowledge about the genetic structure of all remaining populations will reveal patterns of gene flow among populations and the potential for inbreeding depression.

## METHODS AND RESULTS

**Microsatellite development**—Total genomic DNA of 14 individuals (two individuals per population collected across the whole distribution range) of *G. praecox* subsp. *bohemica* was extracted from dehydrated leaves using the cetyltrimethylammonium bromide (CTAB) method of Lodhi et al. (1994), with all amounts downscaled 10×. The sequencing facility GenoScreen (Lille, France) was used to prepare libraries and design primers. Extracted DNA was pooled for microsatellite library preparation. The fragmented DNA was hybridized with eight probes (TG, TC, AAC, AAG, AGG, ACG, ACAT, and ACTC) to enrich the DNA library. Sequencing was performed using a GS FLX sequencer (Roche, 454 Life Sciences, Branford, Connecticut, USA). A total of 19,152 reads were obtained. Raw sequencing data were submitted to the National Center for Biotechnology Information (NCBI) Sequence Read Archive (accession no. SRR5113067). QDD software (Megléczy et al., 2009) with default settings was used to identify microsatellite loci and for design of the microsatellite primers. A total of 3017 reads contained microsatellite motifs, and 373 candidate microsatellite loci were identified (Appendix S1), with an average sequence length of 325 bp. Markers belonged to di-, tri-, tetra-, penta-, and hexanucleotide

<sup>1</sup>Manuscript received 13 September 2016; revision accepted 27 November 2016.

This work was supported by grants from Iceland, Liechtenstein, and Norway (EHP funds 2009–2014) and by the Ministry of the Environment of the Czech Republic. The authors thank Kristina Plenk and Matthias Kropf for providing *Gentianella praecox* subsp. *praecox* samples and Jana Kadlecová and Eva Ničová for technical assistance.

<sup>5</sup>Author for correspondence: maria.surinova@ibot.cas.cz

doi:10.3732/apps.1600114

*Applications in Plant Sciences* 2017 5(1): 1600114; <http://www.bioone.org/loi/apps> © 2017 Šurinová et al. Published by the Botanical Society of America. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC-BY-NC-SA 4.0), which permits unrestricted noncommercial use and redistribution provided that the original author and source are credited and the new work is distributed under the same license as the original.

repeats (40.2%, 52.8%, 5.4%, 0.8%, and 0.8%, respectively). Across all candidate loci, 3378 primer pairs (3–15 primer pairs per locus) were designed using Primer3, as implemented within QDD (Malausa et al., 2011) with amplicon lengths between 90 and 319 bp. For each microsatellite candidate locus, one primer pair was selected for further analysis. Of these, we selected 50 primer pairs (Appendix S1) recommended by GenoScreen to identify polymorphic markers. Primers were synthesized (Sigma-Aldrich, St. Louis, Missouri, USA) with M13 tails preceding the 5' end of the forward primer sequences (Schuelke, 2000). Six individuals from six populations of *G. praecox* subsp. *bohemica* (Appendix 1) were used to test amplification efficiency and polymorphism. DNA amplification was performed in 10-μL reactions consisting of 5 μL of QIAGEN Multiplex PCR Master Mix (QIAGEN, Hilden, Germany), 0.25 μL of each M13-labeled forward, reverse, and fluorolabeled (5'-FAM) M13 primer (10 μM each in initial volume), 20 ng of DNA dissolved in 1 μL TE buffer, and 3.25 μL of H<sub>2</sub>O.

The following PCR protocol was performed using an Eppendorf Mastercycler pro S Thermal Cycler (Eppendorf, Hamburg, Germany): an initial denaturation step at 95°C for 15 min; followed by 25 cycles of denaturation (95°C for 20 s), annealing (59°C for 30 s), and extension (72°C for 20 s); followed by 10 cycles of denaturation (95°C for 30 s), annealing (53°C for 45 s), and extension (72°C for 45 s); and a final extension at 72°C for 10 min. Thirty-eight primer pairs (76%) were successfully amplified. Due to allele dosage uncertainty in polyploid individuals, preliminary statistics included determination of polymorphic information content (PIC) for each locus by PICcalc (Nagy et al., 2012). Based on PIC, 20 (53%) of the 38 primer pairs were selected for detailed variability screening on 36 individuals of *G. praecox* subsp. *bohemica* (two individuals from each population). Based on the multiplex PCR performance and variability screening, 12 polymorphic primer pairs were identified.

To confirm primer specificity for these 12 loci, we ran PCRs for each primer pair separately under the same conditions described in the next paragraph. PCR products were purified using the QIAquick PCR Purification Kit (QIAGEN) and cloned using pGEM-T Vector Systems II (Promega Corporation, Madison, Wisconsin, USA) in accordance with the manufacturer's instructions, but down-scaled to half reactions. Approximately 10 colonies per sample were transferred into 20 μL of ddH<sub>2</sub>O and denatured at 95°C for 10 min. These served as templates for subsequent PCR amplifications for sequencing. Sequencing was performed by the commercial company SEQme (Dobříš, Czech Republic), and the resulting sequences were aligned using MAFFT 7.017 (Kato et al., 2002) as implemented in Geneious 8.1.6 (Kearse et al., 2012). Repeat motifs with variation in number of repeats were confirmed in the obtained sequences. GenBank

accession numbers of identified sequences for 12 loci of *G. praecox* subsp. *bohemica* are provided in Table 1.

**Genotyping**—Total DNA was extracted from 180 *G. praecox* subsp. *bohemica* individuals from six populations and from 114 individuals from eight populations of three closely related taxa (Appendix 1) for initial primer screening. DNA amplification was carried out in three multiplex reactions consisting of 2.5 μL of QIAGEN Multiplex PCR Master Mix and 10 ng of DNA dissolved in 0.5 μL of TE buffer. For multiplex mix I (MM I), the PCR contained 1.1 μL of primer mix (10 μM each in initial volume) and 0.9 μL of H<sub>2</sub>O, for MM II the PCR consisted of 1.1 μL of primer mix (10 μM each in initial volume) and 0.9 μL of H<sub>2</sub>O, and for MM III the PCR contained 0.7 μL of primer mix (10 μM each in initial volume) and 1.3 μL of H<sub>2</sub>O. The sequence, labeling, motif information, final volumes, and PCR product size range are given in Table 1. The following PCR protocol was performed using an Eppendorf Mastercycler pro S Thermal Cycler: an initial denaturation step at 95°C for 15 min; followed by 35 cycles of denaturation (95°C for 20 s), annealing (59°C for 30 s), and extension (72°C for 20 s); and a final extension at 72°C for 10 min. PCR products were diluted with ddH<sub>2</sub>O in these ratios: 1:2 (PCR product of MM I and MM II PCRs: ddH<sub>2</sub>O), 1:9 (PCR product of MM III PCR: ddH<sub>2</sub>O). Each PCR product (1 μL) was mixed with 11 μL of a 120:1 solution of formamide: size standard (GeneScan 500 LIZ; Thermo Fisher Scientific, Waltham, Massachusetts, USA). Fragment lengths were determined by capillary gel electrophoresis with an ABI 3130 Genetic Analyzer using GeneMapper 4.0 (Thermo Fisher Scientific). Using SPAGeDi (Hardy and Vekemans, 2002), we calculated the number of alleles per locus, which ranged between one and nine (Table 2). All markers were polymorphic in all *G. praecox* subsp. *bohemica* populations, except marker GbM48, which was monomorphic in the Zidkovi population. The highest average percentage of heterozygous genotypes was identified for individuals from the Hroby population (75.5%) and the lowest percentage for individuals from the Zidkovi population (50.5%). We detected a high frequency of polyploid individuals (77.8%). The observed heterozygote excess is likely caused by the fact that the species is tetraploid.

We also tested cross-amplification of these loci in three other *Gentianella* taxa: *G. praecox* subsp. *praecox*, *G. amarella* (L.) Börner subsp. *amarella*, and *G. obtusifolia* (F. W. Schmidt) Holub subsp. *sturmiana* (A. Kern. & Jos. Kern.) Holub. We tested 114 individuals from eight populations (Appendix 1). DNA amplification was carried out in three multiplex reactions as described above. Tests for cross-amplification in the three congeneric taxa resulted in successful amplification of up to seven of the 12 polymorphic loci (Table 2). These results (Table 3) demonstrate that these primer pairs may be of broad utility throughout *Gentianella*.

TABLE 1. Characteristics of 12 polymorphic loci designed for genotyping of *Gentianella praecox* subsp. *bohemica*.

Locus <sup>a</sup>	Primer sequences (5'–3')	Repeat motif	PCR product size range (bp)	Fluorescent label	Volume of forward primer (μL)	Multiplex	GenBank accession no.
GbM46	F: CAACCACAAGAAGCTTCCAA R: GCATTGCCAACAGATGCAG	(CTT) <sub>4</sub>	81–129	PET	0.1	I	KX420610
GbM11	F: TGGTTTGATTTCAGACCCCTTG R: CAGGTTGCCCTACCAAGATG	(TTG) <sub>16</sub>	138–180	PET	0.25	I	KX420608
GbM34	F: GAAGCGTCCGTTTCAGTTTC R: GCTTAGAGCCCAAGATACCTAGA	(TGT) <sub>5</sub>	119–152	NED	0.075	I	KX420611
GbM3	F: AGTTGAGAATTGGCCTGGAG R: GATGCATTGGAAGCAGGATT	(GTAT) <sub>5</sub>	134–174	VIC	0.125	I	KX420606
GbM12	F: ATCAGGCATTGCCATTAAGC R: GAGATTTCATAGTTGGCGAGG	(AC) <sub>5</sub>	96–108	VIC	0.15	II	KX420604
GbM19	F: GGAATTCTCTTGTAAGCCAG R: TTGCTGCTTCTTTTCCATGA	(GAG) <sub>8</sub>	136–202	VIC	0.225	II	KX420609
GbM38	F: TTCAAGGTTGCTTTTGGCT R: GCCTTGTTGTTAAATTAGTTGCAG	(AGA) <sub>6</sub>	129–162	NED	0.075	II	KX420612
GbM5	F: CTCCTTCCCTTTTCCCAAC R: GCTTATGTCGCAGTGCAGAA	(AG) <sub>8</sub>	158–180	PET	0.1	II	KX420615
GbM2	F: GGGAGAGCGAGTTCAAAG R: AAGCTGCTAAACTTCAATACTTCG	(GGA) <sub>13</sub>	147–180	VIC	0.075	III	KX420607
GbM48	F: ACCGAAGCGAGTTTCAACAC R: CCAACAACTTAGCTACCTTAGCA	(GGA) <sub>3</sub>	84–93	NED	0.175	III	KX420613
GbM39	F: AACAGAGCAAAAACAAAACAGG R: CAAGAAAGCAATGAATCCCC	(AGA) <sub>8</sub>	79–94	VIC	0.05	III	KX420614
GbM43	F: AATCATGTCCAGCTCAGCCT R: GCCGACGTAGAATGTTTGGT	(CCT) <sub>4</sub>	158–185	NED	0.05	III	KX420605

<sup>a</sup>Optimal annealing temperature was 59°C for all loci.

TABLE 2. Results of initial primer screening of 12 microsatellite loci developed in *Gentianaella praecox* subsp. *bohemica* and congeners.

Species	Population code <sup>a</sup>	Primer/Index <sup>b</sup>	GbM46	GbM11	GbM34	GbM3	GbM12	GbM19	GbM38	GbM5	GbM2	GbM48	GbM39	GbM43
<i>G. praecox</i> subsp. <i>bohemica</i>	HROBY	A	3	6	4	7	5	9	4	4	4	4	5	6
		A <sub>ind</sub>	2.3	2.3	1.43	1.73	2.7	2.8	1.63	2.53	1.43	1.77	1.5	2.6
	PODVORI	% Het	96.7	86.7	40	73.3	100	96.7	60	96.7	43.3	63.3	50	100
		A	6	4	4	3	5	5	4	4	5	2	3	5
	POLNA	A <sub>ind</sub>	2.3	1.9	1.43	1.47	1.7	1.9	1.27	2.57	1.33	1.2	1.3	2.57
		% Het	86.7	86.7	36.7	46.7	56.7	70	26.7	100	33.3	20	30	100
<i>G. amarella</i> subsp. <i>amarella</i>	VYSNY	A	4	5	4	4	4	5	4	4	6	3	2	5
		A <sub>ind</sub>	1.43	2.07	1.83	1.5	2.17	2	1.43	2.47	1.5	1.3	1.13	2.43
	ZIDKOVI	% Het	40	80	73.3	50	83.3	80	43.3	100	50	30	13.3	100
		A	7	3	2	3	5	6	3	5	3	3	3	4
	VANIC	A <sub>ind</sub>	1.73	1.73	1.5	1.13	2.13	2.63	1.53	2.6	1.5	1.27	1.5	2.23
		% Het	73.3	73.3	50	13.3	83.3	100	53.3	100	50	26.7	50	100
<i>G. obtusifolia</i> subsp. <i>sturmiana</i>	ČER.S.	A	5	2	4	6	2	4	2	3	3	1	2	3
		A <sub>ind</sub>	2.7	1.77	1.1	1.37	1.53	2.3	1.03	2.03	1.13	1	1.17	2.1
	PP PILA	% Het	100	76.7	10	36.7	53.3	100	3.3	100	13.3	0	16.7	96.7
		A	9	4	3	4	4	5	4	4	3	3	3	4
	KOCOL	A <sub>ind</sub>	2.8	2.27	1.03	1.53	1.63	2.77	1.67	2.47	1.43	1.53	1.57	2.53
		% Het	96.7	100	6.7	53.3	60	96.7	66.7	100	43.3	46.7	56.7	100
<i>G. praecox</i> subsp. <i>praecox</i>	RANK	A	4	3	2	3	4	4	4	3	4	2	3	7
		A <sub>ind</sub>	1.63	1.67	1.17	1.33	2.17	2.17	1.17	2.17	83.3	83.3	16.7	2.83
	BUBE	% Het	60	50	16.7	33.3	100	100	16.7	100	83.3	83.3	16.7	100
		A	4	1	1	1	2	2	1	2	2	2	1	3
	GIEE	A <sub>ind</sub>	2.53	1	1	1	2	2	1	2	2	2	1	3
		% Het	96.7	0	0	0	100	100	0	100	100	100	0	100
<i>G. praecox</i> subsp. <i>praecox</i>	LEOE	A	4	5	4	6	2	7	4	2	2	3	3	4
		A <sub>ind</sub>	1.43	2.3	1.6	1.5	1.7	2.6	1.6	1.2	1.1	1.4	1.4	2.6
	RANK	% Het	43.3	90	60	50	70	90	60	20	10	40	40	100
		A	4	4	3	3	4	7	4	3	1	1	2	4
	BUBE	A <sub>ind</sub>	1.77	2.5	1.7	1.6	2.3	2.6	1.8	1.8	1	1	1.1	2.8
		% Het	63.3	100	70	60	100	100	80	70	0	0	10	100
<i>G. praecox</i> subsp. <i>praecox</i>	LEOE	A	5	3	1	2	3	2	1	2	1	1	1	2
		A <sub>ind</sub>	1.5	1.1	1	1.3	2.3	2	1	2	1	1	1	2
	GIEE	% Het	50	10	0	30	100	100	0	100	0	0	0	100
		A	6	2	3	4	4	3	2	2	4	5	3	2
	LEOE	A <sub>ind</sub>	2.6	1.71	1.21	1.29	1.54	1.71	1.21	2.29	1.25	1.21	1.17	2.46
		% Het	100	62.5	20.8	29.2	45.8	66.7	20.8	100	25	20.8	16.7	100
<i>G. praecox</i> subsp. <i>praecox</i>	LEOE	A	1	1	2	2	2	1	1	2	3	3	3	2
		A <sub>ind</sub>	1.8	1.25	1	1	1.05	1	1.4	2	1	1.25	1.15	2.3
	LEOE	% Het	75	25	0	0	5	0	40	20	0	25	15	20
		A	2	2	2	1	2	1	1	2	3	4	2	2
	LEOE	A <sub>ind</sub>	1.79	1	1	1.13	1.83	1	1.04	2	1.13	1.04	1.08	2.13
		% Het	79.2	0	0	12.5	83.3	0	4.2	100	12.5	4.2	8.3	100

Note: A = number of alleles; A<sub>ind</sub> = mean number of alleles per individual; % Het = percentage of heterozygous genotypes.

<sup>a</sup>Population code refers to collecting locality. Detailed information is provided in Appendix 1.

<sup>b</sup>Due to allele ambiguity in polyploids, the H<sub>o</sub>:H<sub>e</sub> ratio is replaced by the percentage of heterozygous genotypes.

TABLE 3. Allele size ranges obtained during cross-amplification trials of microsatellite loci isolated from *Gentianella praecox* subsp. *bohemica* and tested in three additional taxa.

Locus	<i>G. amarella</i> subsp. <i>amarella</i>	<i>G. praecox</i> subsp. <i>praecox</i>	<i>G. obtusifolia</i> subsp. <i>sturmiana</i>
GbM46	111–126	114–117	114–126
GbM11	159–168	150–165	138–168
GbM34	131–134	128–131	119–131
GbM3	142–170	134–158	134–170
GbM12	100–102	100–106	100–108
GbM19	142–154	142–157	142–178
GbM38	137–143	137–146	134–153
GbM5	168–176	168–176	168–178
GbM2	150–177	150–156	147–153
GbM48	84–87	87–90	84–90
GbM39	85–91	82–91	79–85
GbM43	167–179	164–176	167–176

CONCLUSIONS

We developed and successfully multiplexed 12 polymorphic markers in several taxa of *Gentianella*. These polymorphic loci will be valuable for the future management of the extremely rare *G. praecox* subsp. *bohemica*.

LITERATURE CITED

BRABEC, J. 2010. Monitoring hořečku mnohotvarého českého (*Gentianella praecox* subsp. *bohemica*) v ČR (sezóna 2010). [Monitoring of *Gentianella praecox* subsp. *bohemica* in the Czech Republic (vegetation season 2007)]. Deposit in: Central Office of Agentura Ochrany Přírody a Krajiny České Republiky, Prague, Czech Republic.

COUNCIL OF THE EUROPEAN COMMUNITY. 1992. Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Official Journal of the European Communities Series L* 206: 7–49.

GREIMLER, J., AND C. G. JANG. 2007. *Gentianella stiriaca*, a case of reticulate evolution in the northeastern and eastern Central Alps. *Taxon* 56: 857–870.

HARDY, O. J., AND X. VEKEMANS. 2002. SPAGeDi: A versatile computer program to analyse spatial genetic structure at the individual or population levels. *Molecular Ecology Notes* 2: 618–620.

KATOH, K., K. MISAWA, K. KUMA, AND T. MIYATA. 2002. MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30: 3059–3066.

KEARSE, M., R. MOIR, A. WILSON, S. STONES-HAVAS, M. CHEUNG, S. STURROCK, S. BUXTON, ET AL. 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics (Oxford, England)* 28: 1647–1649.

KÖNIGER, J., C. A. REBERNIG, J. BRABEC, K. KIEHL, AND J. GREIMLER. 2012. Spatial and temporal determinants of genetic structure in *Gentianella bohemica*. *Ecology and Evolution* 2: 636–648.

LODHI, M. A., G. N. YE, N. F. WEEDEN, AND B. I. REISCH. 1994. A simple and efficient method for DNA extraction from grapevine cultivars and *Vitis* species. *Plant Molecular Biology Reporter* 12: 6–13.

MALAUSSA, T., A. GILLES, E. MEGLÉCZ, H. BLANQUART, S. DUTHOY, C. COSTEDOAT, V. DUBUT, ET AL. 2011. High-throughput microsatellite isolation through 454 GS-FLX Titanium pyrosequencing of enriched DNA libraries. *Molecular Ecology Resources* 11: 638–644.

MEGLÉCZ, E., C. COSTEDOAT, V. DUBUT, A. GILLES, T. MALAUSSA, N. PECH, AND J. F. MARTIN. 2009. QDD: A user friendly program to choose microsatellite markers and design primers from large sequencing projects. *Bioinformatics (Oxford, England)* 26: 403–404.

NAGY, S., P. POZAI, I. CERNÁK, A. M. GORJI, G. HEGEDUS, AND J. TALLER. 2012. PICcalc: An online program to calculate polymorphic information content for molecular genetic studies. *Biochemical Genetics* 50: 670–672.

OBERDORFER, E. 1983. Pflanzensoziologische Exkursionsflora. Eugen Ulmer Verlag, Stuttgart, Germany.

PLENK, K., F. GÖD, M. KRIECHBAUM, AND M. KROPP. 2016. Genetic and reproductive characterisation of seasonal flowering morphs of *Gentianella bohemica* revealed strong reproductive isolation and possible single origin. *Plant Biology* 18: 111–123.

SCHUELKE, M. 2000. An economic method for the fluorescent labeling of PCR fragments. *Nature Biotechnology* 18: 233–234.

WINFIELD, M. O., P. J. WILSON, M. LABRA, AND J. S. PARKER. 2003. A brief evolutionary excursion comes to an end: The genetic relationship of British species of *Gentianella* sect. *Gentianella* (Gentianaceae). *Plant Systematics and Evolution* 237: 137–151.

APPENDIX 1. Accession information for *Gentianella* species used in this study.<sup>a</sup>

Species name	Population code	Collection locality	Country	n	Latitude	Longitude
<i>G. praecox</i> (A. Kern. & Jos. Kern.) Dostál ex E. Mayer subsp. <i>bohemica</i> (Skalický) Holub	HROBY	Hroby	CZ	30	49.3932222	14.85622
<i>G. praecox</i> subsp. <i>bohemica</i>	PODVORI	Podvoří	CZ	30	48.8356111	14.20819
<i>G. praecox</i> subsp. <i>bohemica</i>	POLNA	Polná in the Šumava Mountains	CZ	30	48.7928056	14.14997
<i>G. praecox</i> subsp. <i>bohemica</i>	VYSNY	Vyšný	CZ	30	48.8266944	14.30211
<i>G. praecox</i> subsp. <i>bohemica</i>	ZIDKOVI	Olešnice in the Orlické Mountains	CZ	30	50.3619444	16.28389
<i>G. praecox</i> subsp. <i>bohemica</i>	VANIC	Nature Reserve Opolenec	CZ	30	49.0866667	13.79706
<i>G. amarella</i> (L.) Börner subsp. <i>amarella</i>	VANIC	Nature Reserve Opolenec	CZ	6	49.0866667	13.79706
<i>G. amarella</i> subsp. <i>amarella</i>	ČER.S.	Kouty nad Desnou	CZ	10	50.1225	17.16111
<i>G. obtusifolia</i> (F. W. Schmidt) Holub subsp. <i>sturmiana</i> (A. Kern. & Jos. Kern.) Holub	PP PILA	Pila u Karlových Varů	CZ	10	50.1747222	12.92694
<i>G. obtusifolia</i> subsp. <i>sturmiana</i>	KOCEL	Kocelovice	CZ	10	49.475	13.82444
<i>G. obtusifolia</i> subsp. <i>sturmiana</i>	RANK	Rankovice	CZ	10	50.0067778	12.84208
<i>G. praecox</i> subsp. <i>praecox</i>	BUBE	Buchberg, Lower Austria	AU	24	48.376944	15.39722
<i>G. praecox</i> subsp. <i>praecox</i>	GIEE	Gießhübl, Lower Austria	AU	20	48.320833	15.36306
<i>G. praecox</i> subsp. <i>praecox</i>	LEOE	Leopolds, Lower Austria	AU	24	48.427778	15.28611

Note: AU = Austria; CZ = Czech Republic; n = number of individuals.

<sup>a</sup>Because all investigated species are rare and highly protected, it was not possible to sample whole plants for herbarium vouchers. Leaf samples were collected in the field for up to five individuals per population and were dried in silica gel before performing DNA extraction. The leaf samples and DNA extracts were deposited at the Institute of Botany of the Czech Academy of Sciences, Průhonice, Czech Republic.