

## **Fossil Vesicomyid Bivalves from Miocene Hydrocarbon Seep Sites, North Island, New Zealand**

Authors: Amano, Kazutaka, Saether, Kristian P., Little, Crispin T.S., and Campbell, Kathleen A.

Source: Acta Palaeontologica Polonica, 59(2) : 421-428

Published By: Institute of Paleobiology, Polish Academy of Sciences

URL: <https://doi.org/10.4202/app.2012.0070>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Fossil vesicomimid bivalves from Miocene hydrocarbon seep sites, North Island, New Zealand

KAZUTAKA AMANO, KRISTIAN P. SAETHER, CRISPIN T.S. LITTLE,  
and KATHLEEN A. CAMPBELL



Amano, K., Saether, K.P., Little, C.T.S., and Campbell, K.A. 2014. Fossil vesicomimid bivalves from Miocene hydrocarbon seep sites, North Island, New Zealand. *Acta Palaeontologica Polonica* 59 (2): 421–428.

Two fossil species of vesicomimids are described from Lower to Middle Miocene hydrocarbon seep carbonates in eastern North Island, New Zealand. One elongate species is proposed as a new genus and species: *Notocalyptogena neozelandica*. The other species probably belongs to the genus *Pliocardia*, but due to poor preservation is not identified further. The composition of this Miocene vesicomimid seep fauna differs from that found in modern New Zealand seeps located on the offshore Hikurangi convergent margin, which contain the genera *Calyptogena*, *Archivesica*, and *Isorropodon*. The fossil fauna went extinct locally after the Middle Miocene and has been since replaced by the modern vesicomimid taxa.

Key words: Mollusca, Bivalvia, Vesicomiyidae, hydrocarbon seep, Miocene, New Zealand.

Kazutaka Amano [amano@juen.ac.jp], Department of Geoscience, Joetsu University of Education, 1 Yamayashiki, Joetsu 943-8512, Japan;

Kristian P. Saether [kris.saether@gmail.com] and Kathleen A. Campbell [ka.campbell@auckland.ac.nz], School of Environment, Faculty of Science, University of Auckland, Private Bag 92019, Auckland Mail Centre, Auckland 1142, New Zealand;

Crispin T.S. Little [earctsl@leeds.ac.uk], School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK.

Received 18 May 2012, accepted 24 August 2012, available online 7 September 2012.

Copyright © 2014 K. Amano et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Introduction

The bivalve family Vesicomiyidae is a characteristic and species-rich molluscan taxon among the modern chemosynthesis-based fauna from hydrocarbon seeps (e.g., Smith and Baco 2003; Levin 2005; Taylor and Glover 2010; Kiel 2010). For some time the taxonomy of the family has been in a state of flux with, for example, some authors placing all large vesicomimid species in the genus *Calyptogena* (e.g., Boss and Turner 1980; Okutani et al. 2000). However, recent detailed work on shell and soft part morphology, combined with molecular data, has resolved a number of well characterized genera (Krylova and Sahling 2006, 2010; Amano and Kiel 2007; Krylova and Cosel 2011). Moreover, Krylova and Sahling (2010) subdivided Vesicomiyidae into two subfamilies: Vesicomiyinae and Pliocardiinae. The subfamily Vesicomiyinae comprises the small-sized genus *Vesicomya* only; the subfamily Pliocardiinae contains medium to large sized genera with reduced guts and thiotrophic bacterial symbionts in their gills (e.g., Dubilier et al. 2008). The majority of occurrences of modern and fossil vesicomimid species are from the northern hemisphere, with few data points from

high southern latitudes, which is likely an artifact of research effort rather than true distributional bias.

One southern high latitude area for which there are records of fossil and modern vesicomimids is New Zealand. Lewis and Marshall (1996) first described and illustrated modern vesicomimids from hydrocarbon seeps of the Hikurangi margin, offshore eastern North Island. They recorded *Calyptogena* spp. A, B, C, and *Vesicomya* sp. While it is evident that *Calyptogena* sp. A should be included in *Calyptogena* sensu stricto, a review of this genus by Krylova and Sahling (2006) did not consider the New Zealand records. Campbell et al. (2010) also illustrated *Calyptogena* sp. and another vesicomimid from the Hikurangi margin deposits. Of these, their *Calyptogena* sp. resembles *Calyptogena* sp. A noted by Lewis and Marshall (1996).

Many inferred chemosynthesis-based fossils also are known from Miocene hydrocarbon seep deposits in eastern North Island. McKay (1877a, b) was the first to describe these geographically isolated carbonates within voluminous mudstones, and noted that they contained coquinas of fossil mussels. The carbonates were recognized in early geological mapping of the region (Adams 1910; Henderson and Ong-




Fig. 1. Geological setting of the Miocene seep sites and fossil localities. **A.** Boundary between the Australian and Pacific Plates, position of the Hikurangi subduction zone (HSZ) and arc volcanoes of the Taupo Volcanic Zone (TVZ); M, Marlborough. **B.** Overview of the geology of the East Coast centered around Hawke's Bay, North Island, New Zealand, showing locations of known Miocene hydrocarbon seep sites (numbered circles). 1–11 = northern sites: 1, Waipapu; 2, Waipiro; 3, Karikarihuata; 4, Bexhaven; 5, Tauwharepare; 6, Puketawa; 7, Totaranui; 8, Moonlight North; 9, Rocky Knob; 10, Waikairo; 11, Turihaua. 12–16 = southern sites: 12, Wanstead; 13, Ugly Hill; 14, Haunui; 15, Ngawaka; 16, Wilder. The fossil vesicomysids described in this study are from fossil seep locations 4, 6, 8, 9, 12, 13, and 14. **C.** Cross-section (modified from Barnes 2010) showing transpressive subduction of the Pacific Plate beneath the Australian Plate, and relationships of tectonic elements of the northern New Zealand plate margin (see Campbell et al. 2008 and Barnes et al. 2010 for further details on geologic context).

ley 1920), and termed “*Modiolus* limestone” by Ongley and MacPherson (1928). The first stratigraphic grouping of these deep-water carbonates was made by Kamp and Nelson (1988) in a compendium of limestone occurrences from the Neogene plate margin of New Zealand, in which they were informally dubbed the “Moonlight limestone” after a nearby sheep station. Beu and Maxwell (1990) noted that the Moonlight deposits occur widely in the Miocene East Coast Basin (cf. Field et al. 1997) of eastern North Island, as isolated pods from 10–100 m across. They recorded a mytilid resembling *Idasola*, *Lucinoma* aff. *taylori*, trochid and lepetellid gastropods. However, they did not comment on any other taxa, including vesicomysids, from these sites. Mazengarb et al. (1991) for-

mally elevated the scattered Moonlight/*Modiolus* limestone occurrences to formation status, describing them collectively as the Bexhaven Limestone. These limestone lenses occur in deep-water massive mudstone deposits of the Tolaga Group (Early to Late Miocene; Mazengarb and Speden 2000) north of Gisborne. Similar limestones also have been reported from the southern Hawke's Bay area, which were mapped as the Ihungia Limestone by Lillie (1953).

Campbell et al. (2008) established that the Bexhaven and Ihungia limestone deposits, from north and south of Hawke's Bay, respectively, are hydrocarbon seep-related. The 14 reported fossil seep carbonate deposits developed during the Early to Late Miocene, when the modern convergent plate

boundary was initiated offshore eastern North Island, and are preserved today in deformed and uplifted deep-marine forearc strata for 300 km along the continental margin, from East Cape to Dannevirke (Fig. 1). Subsequent study of the Miocene seep limestone localities from New Zealand has recognized 16 sites, and described the petrographic, stable isotopic and paleontological character of several of the deposits (Ewen 2009; Troup 2010; Saether et al. 2010a, b, 2012; Saether 2011).

Campbell et al. (2008) and Saether (2011) illustrated fossil vesicomys from seven locations in the Hawke’s Bay area (Fig. 1), and briefly compared them with the modern vesicomys offshore. However, these fossil vesicomys have never been formally studied. In this paper, we describe the vesicomys from Lower to Middle Miocene seep sites in the Hawke’s Bay area. We also discuss the biogeographic significance of these species by comparing them with the undescribed modern seep fauna offshore.

*Institutional abbreviations.*—AU, Paleontology Collection, School of Environment, UOA; UOA L, bivalve specimen number, Paleontology Collection, School of Environment, UOA; GNS, GNS Science, Lower Hutt, New Zealand; NMNZ, Museum of New Zealand Te Papa Tongarewa, Wellington, New Zealand; UOA, University of Auckland, Auckland, New Zealand.

*Other abbreviations.*—AL, anterior length; H, height; L, length; W, width; Y and U, map codes in New Zealand.

## Material

The vesicomys specimens described herein were collected from the geographically isolated Lower to Middle Miocene deep-water seep limestone deposits to the north of Gisborne and east of Dannevirke, East Coast Basin, North Island, New Zealand (Fig. 1, Table 1). In the east of Dannevirke area, vesicomys were collected from the Lower Miocene Ihungia Limestone at the Ugly Hill, Wanstead, and Haunui seep fossil localities; north of Gisborne, vesicomys specimens were collected from the Lower to Middle Miocene Bexhaven Limestone seep sites at Bexhaven, Puketawa, Moonlight North, and Rocky Knob localities. Foraminiferal data (e.g., Hayward 1986) and lithological comparison to dredgings from modern NZ seeps (Saether et al. 2010b) suggest that these vesicomys likely occupied seep environments at roughly 500–2000 m depth in the Miocene. The lithological, stable isotopic and faunal character of these fossil seep deposits have been described in detail elsewhere (Campbell et al. 2008; Saether et al. 2010a, b, 2012; Saether 2011).

All fossil localities of the illustrated Miocene specimens (e.g., Y16/f1059, U23/f267) are registered in the New Zealand Fossil Record File database (<http://www.fred.org.nz>) jointly administered by GNS Science and the New Zealand Geoscience Society. Details of comparative modern vesicomys specimens from New Zealand are shown in Table 1.

Table 1. Collection and storage details of material discussed herein.

Specimens	Collection and storage institution
Fossils	by the authors in February 2012, UOA
Fossils	by the geology program, School of Environment, UOA
Fossils	by UOA paleontologists between 1997–2010; UOA
Fossils	by GNS Science
Modern	by 2007 joint German-New Zealand NEW VENTS Hikurangi margin cruise with the R/V SONNE; UOA
Modern	NMNZ

## Systematic paleontology

Class Bivalvia Linnaeus, 1758

Subclass Heterodonta Neumayr, 1884

Family Vesicomysidae Dall and Simpson, 1901

Subfamily Pliocardiinae Woodring, 1925

?Genus *Pliocardia* Woodring, 1925

*Type species:* *Anomalocardia bowdeniana* Dall, 1903; Bowden Formation, Late Pliocene, Bowden, Jamaica.

*Pliocardia?* sp.

Fig. 2A–C.

*Material.*—Four specimens from Moonlight North (locality Y16/f1059, collection AU19982, specimen numbers UOA L4587, UOA L4588, UOA L4589, UOA L4590); Lower to Middle Miocene, Bexhaven Limestone.

*Description.*—Shell medium size (L up to 36.2 mm), quadrate-ovate (H/L = 0.71–0.81), moderately inflated (W/L = 0.50–0.59), shell material partly to wholly absent in all specimens; surface sculpture unknown except for faint growth lines. Blunt ridge running from beak to postero-ventral corner. Beak prominent, prosogyrate, situated anteriorly at




Fig. 2. Vesicomysid bivalve *Pliocardia?* sp. from Moonlight North (Y16/f1059), Early to Middle Miocene. A. UOA L4587. Dorsal view of articulated internal mould (A<sub>1</sub>), arrow shows lunular incision. Internal mould of right valve (A<sub>2</sub>), showing anterior adductor scar and posterior adductor scar. B. UOA L4588. Right valve of hinge plate (B<sub>1</sub>), anterior tooth is not preserved. Left valve (B<sub>2</sub>), internal mould, arrow shows pallial sinus.

24–27% along shell length from anterior margin. Antero-dorsal margin short and broadly arcuate; postero-dorsal margin straight to gently convex, continuing to subtruncated posterior margin; ventral margin usually moderately convex. Lunular incision weakly visible in dorsal view; large, deep grooves flanking central ligamental area along postero-dorsal margin, suggesting ligament external. Hinge plate of right valve narrow, with very thin ventral tooth (1); rather strong posterior tooth (3b); anterior cardinal tooth (3a?) obscure. Anterior adductor muscle scar ovate; posterior muscle scar subquadrate. Pallial sinus indistinct and shallow.

*Dimensions.*—See Table 2.

Table 2. Measurements of *Pliocardia?* sp. Abbreviations: AL, anterior length (\* distance of umbo from anterior margin); H, height; L, length; W, width (\*\* width of articulated shell).

UOA specimen	L (mm)	H (mm)	AL (mm)*	W (mm)**	H/L	L/H	AL/L	Valve
L4587	29.2	21.8	6.9	14.6	0.75	1.34	0.24	both
L4588	28.8	21.7	7.8	—	0.74	1.33	0.27	left
L4589	36.2	29.4	8.7	—	0.81	1.23	0.24	right
L4590	11.2	9.0	3.0	6.2	0.80	1.24	0.27	both

*Discussion.*—From its size, blunt ridge, lunular incision, right valve hinge dentition and pallial sinus, this species probably represents a new species in the genus *Pliocardia* Woodring, 1925, which was redefined by Krylova and Janssen (2006). However, until more details are known about the external shell and the left valve dentition we refrain from naming a new species.

Our species resembles *Pliocardia kawadai* (Aoki, 1954) from the Lower to Middle Miocene in Japan (see Amano and Kiel 2012) in having a similar size and outline. However, *P. kawadai* has more inflated valves, a radial depression, and a V-shaped pallial sinus.

### Genus *Notocalyptogena* nov.

*Etymology:* A combination of Greek *notos*, south and the genus *Calypptogena*.

*Type species:* *Notocalyptogena neozelandica* sp. nov.; see below.

*Species included:* Only type species.

*Diagnosis.*—Shell medium size, moderately inflated, elongate; sculpture smooth except for growth lines; blunt external ridge running from beak to posterior corner. Radial internal ridge distinct, running from beak to postero-ventral corner and in contact with posterior adductor scar. Hinge plate narrow. Subumbonal pit absent. Hinge of right valve with three cardinal teeth; anterior tooth (3a) along postero-dorsal margin; middle tooth (1) strong, oblique anteriorly; posterior tooth (3b) vertical or slightly anteriorly inclined; deeply depressed or flat area behind posterior cardinal tooth. Hinge of left valve with three cardinal teeth; anterior tooth (2a) long, connecting with middle stout tooth (2b); posterior tooth (4b) also rather stout. Pallial line entire.

*Discussion.*—*Notocalyptogena* gen. et sp. nov. closely resembles *Calypptogena* Dall, 1891 in its size, elongate outline, lack of subumbonal pit, and integripalliate condition. However, this new genus differs from *Calypptogena* by having a narrow hinge plate without a U-shaped tooth overhanging a ventral tooth, and by having a depressed or flat area behind the 3b tooth in the right valve. Moreover, the strong internal ridge from the beak to posterior ventral corner of this new genus is not seen in *Calypptogena*. *Notocalyptogena* gen. nov. shares a narrow hinge plate, elongate shell shape, lack of subumbonal pit and an entire pallial line with *Christineconcha* Krylova and Cosel, 2011. However, the right valve hinge of *Christineconcha* has a much shorter 3a tooth and teeth that radiate less around the umbo. *Elenaconcha* Cosel and Olu, 2009 can be easily distinguished from *Notocalyptogena* gen. nov. by having a lunule, a subumbonal pit and multiple “posterior nymphal ridges”. Although *Hubertschenckia* Takeda, 1953 has a similar arrangement of cardinal teeth to *Notocalyptogena* gen. nov., the former genus also has a subumbonal pit and pallial sinus, which are never seen in the new genus. *Adulomya* Kuroda, 1931 shares an elongate shell and a lack of a subumbonal pit with *Notocalyptogena* gen. nov. However, *Adulomya* differs from the new genus by having two radiating cardinal teeth in the right valve.

*Stratigraphic and geographic range.*—Lower Miocene Ihungia Limestone and Lower to Middle Miocene, Bexhaven Limestone, North Island, New Zealand.




Fig. 3. Hinge of vesicomyid bivalve *Notocalyptogena neozelandica* gen. et sp. nov. from Ugly Hill (U23/f267 for A, B, D; U23/f266 for C), Early Miocene. Left (A, C) and right (B, D) valve hinge. A. UOA L4606. B. UOA L4605. C. UOA L4596, paratype. D. UOA L4593, paratype.

*Notocalyptogena neozelandica* sp. nov.

Figs. 3A–D, 4A–G.


*Etymology*: Named for the country of origin.*Type material*: Holotype UOA L4591 (Fig. 4D). Paratypes: UOA L4592 (Fig. 4G), UOA L4593 (Fig. 3D), UOA L4594 (Fig. 4F), UOA L4595 (Fig. 4C) from locality U23/f267, collection AU19983; UOA L4596 (Fig. 3C) from locality U23/f266, collection AU19664.*Type horizon*: Ihungia Limestone, Lower Miocene.*Type locality*: Ugly Hill, North Island, New Zealand.*Material*.—11 poorly to well preserved specimens from Ugly Hill: U23/f267, AU19983. Among them, the specimens catalogued as UOA L4606 (Fig. 3A), UOA L4605 (Fig. 3B), UOA L4607 (Fig. 4A), UOA L4600 (Fig. 4E) show the internal structure of shell. The specimen as UOA L4603 (Fig. 4B) shows the surface of left valve.*Dimensions*.—See Table 3.*Diagnosis*.—As for the genus.

Fig. 4. Vesicomyid bivalve *Notocalyptogena neozelandica* gen. et sp. nov. from Ugly Hill (U23/f267), Early Miocene. **A**. UOA L4607; right valve showing internal ridge from beak to posterior corner (arrow). **B**. UOA L4603; external surface of left valve. **C**. UOA L4595, paratype; dorsal view showing strong ligament. **D**. UOA L4591, holotype; dorsal view (**D**<sub>1</sub>). Note blunt external ridge from beak to posterior ventral corner (arrow). External right valve view (**D**<sub>2</sub>). **E**. UOA L4599; right valve showing anterior adductor scar. **F**. UOA L4594, paratype; left valve internal mould showing pallial line without sinus (arrow) and posterior adductor scar. **G**. UOA L4592, paratype; left valve.




Fig. 5. Modern vesicomids from the Hikurangi Margin. **A, D.** *Calyptogena* sp. **A.** UOA L4610; right valve external ( $A_1$ ) and internal ( $A_2$ ) views. **D.** UOA L4611; external views of right ( $D_1$ ) and left ( $D_2$ ) valves. **B, C.** *Archivesica* sp. **B.** UOA L4608; left valve hinge. **C.** UOA L4609; left valve external ( $C_1$ ) and internal ( $C_2$ ) views.

*Description.*—Shell medium size (L up to 51.7 mm), rather thick, moderately inflated, elongate-ovate ( $H/L = 0.43\text{--}0.59$ ), equivalve and inequilateral. Antero-dorsal margin broadly curved into narrowly rounded anterior margin; postero-dorsal margin nearly straight, steeply sloping into oblique posterior margin at obtuse angle; posterior end acutely rounded; ventral margin slightly convex. Blunt external ridge running from beak to posterior corner. Internal radial ridge also prominent from beak to just behind posterior adductor muscle scar, making deep groove on internal moulds. Beak prominent, swollen, prosogyrate and located at anterior one-fifth to one-third of shell length (at 20–37% of shell length from anterior margin). Nymph long, occupying about half of shell length. Lunule and lunular incision absent. Ligament strong, occupying about 60% of postero-dorsal length. Shell surface with numerous growth lines. Hinge of right valve with three cardinal teeth; anterior tooth (3a) thin, disposed along antero-dorsal margin; middle tooth (1) strong, oblique anteriorly; posterior tooth (3b) thin, vertical or slightly anteriorly inclined; deeply depressed or flat area behind posterior cardinal tooth. Hinge of left valve with three cardinal teeth; anterior tooth (2a) long and thin, connecting with middle stout tooth (2b); posterior

tooth (4b) also rather stout. Pallial line entire. Anterior adductor scar ovate; posterior adductor scar pyriform.

*Discussion.*—The modern vesicomid, *Calyptogena* sp. A by Lewis and Marshall (1996) and *Calyptogena* sp. by Campbell et al. (2010), collected from the Hikurangi margin, can be distinguished from the new species by having a broad posterior tooth (3b) connecting with anterior tooth (3a), no depressed area behind 3b tooth, and no distinct inner ridge running from beak to postero-ventral corner.

*Stratigraphic and geographic range.*—Lower Miocene, Ihungia Limestone, Ugly Hill, Wanstead and Haunui; Lower to Middle Miocene, Bexhaven Limestone, Bexhaven, Moonlight North, and Rocky Knob.

## Modern New Zealand vesicomids and palaeobiogeographic implications

Among the modern shell collections from seeps sites on the Hikurangi margin of New Zealand (Table 1) referred to in

Table 3. Measurements of *Notocalyptogena neozelandica* gen. et sp. nov. Abbreviations: AL, anterior length (\* distance of umbo from anterior margin); H, height; L, length; W, width (\*\* width of each valve).

UOA specimen	Type	L (mm)	H (mm)	AL (mm)*	W (mm)**	H/L	L/H	AL/L	Valve
L4591	holotype	37.6	20.0	9.3	7.4	0.53	1.88	0.25	right
L4592	paratype	42.7	21.2	10.0	8.9	0.50	2.01	0.23	left
L4597		34.9	18.5	6.9	—	0.53	1.89	0.20	right
L4598		29.4	14.5	8.1	—	0.49	2.03	0.28	right
L4599		32.5	18.5	9.3	—	0.57	1.76	0.29	right
L4600		31.3	16.9	8.1	—	0.54	1.85	0.26	right
L4601		33.0	17.9	7.3	—	0.54	1.84	0.22	right
L4602		33.6	17.0	6.9	—	0.51	1.98	0.21	left
L4603		30.8	15.3	7.0	—	0.50	2.01	0.23	left
L4604		38.3	17.2	8.2	—	0.45	2.23	0.21	left

Campbell et al. (2010) are three modern vesicomid species: *Calyptogena* sp., *Archivesica* sp., and *Isorropodon* sp.

*Calyptogena* sp. (Fig. 5A, D) was described as *C.* sp. A and B by Lewis and Marshall (1996: figs. E–H). This species has an elongate shell and a low posterior cardinal tooth in the right valve. It is similar in appearance to *C. tuerkayi* Krylova and Janssen, 2006 from the Edison Seamount and *C. makranensis* Krylova and Sahling, 2006 from the Makran margin, off Pakistan. *Archivesica* sp. (Fig. 4B, C) was illustrated in Campbell et al. (2010: fig. 6D, E) as a vesicomid bivalve. A more detailed examination of this specimen shows it has a subumbonal pit, pallial sinus and three cardinal teeth in the right valve. Its outline is very close to that of *Archivesica nanshaensis* (Xu and Shen, 1991) from the South China Sea (see also Lutaenko and Xu 2008). *Isorropodon* sp. was described as *Vesicomya* sp. A, B by Lewis and Marshall (1996: figs. I–L). Judging from its size, the left valve dentition and ill-defined lunular incision, this species belongs to *Isorropodon*. Of these modern vesicomids from the Hikurangi margin, *Calyptogena* sp. is the dominant species.

A similar dominance of one vesicomid species, in this case *Notocalyptogena neozelandica*, is seen in the Miocene New Zealand vesicomid fauna, with *Pliocardia?* sp. being a relatively uncommon element. However, the modern New Zealand vesicomids have no phylogenetic relationship at genus level with the fossil seep vesicomid fauna. This indicates that at some point since the Miocene the New Zealand vesicomid seep fauna suffered a local extinction and has been replaced by genera with probable South China Sea or South Pacific origins.

While *Notocalyptogena* was present in seeps in New Zealand in the Early and Middle Miocene, contemporary Pacific seep sites in Japan and Alaska were dominated by the genera *Adulomya* and *Archivesica* (Kanno 1971; Kiel and Amano 2010; Amano and Kiel 2011), and *Calyptogena* was absent.

## Acknowledgements

We thank Bruce Marshall (Te Papa Museum, Wellington, New Zealand) for showing us modern *Calyptogena* specimens from seeps of the Hikurangi margin, Neville Hudson (University of Auckland, New Zealand)

for his help with fossil curating and access to material stored in the University of Auckland paleontological collections, and Steffen Kiel (University of Göttingen, Germany) for his comments on some vesicomids collected by Kris Saether. Iain McInnes (Ernslaw One Ltd., Gisborne, New Zealand) kindly allowed access to the Moonlight North site. James and Sue Hewitt (Wanstead, New Zealand) granted access to the Haunui site, and Tim and Maggie Simcox, and Paul and Kate Dearden (Wanstead, New Zealand) gave us permission to work at Ugly Hill. Tim Simcox (Wanstead, New Zealand) in particular provided key technical support that allowed extraction of adequate fossil hinges, making this study possible. Campbell Nelson (University of Waikato, Hamilton, New Zealand) and David Francis (Geological Research Ltd., Lower Hutt, New Zealand) supplied us with field logistical assistance. We also thank reviewers Rudo von Cosel (Muséum national d'Histoire naturelle, Paris, France) and Elena Krylova (Institute of Oceanology RAS, Moscow, Russia) for their useful comments resulting in the improvement of the manuscript. This research was partly supported by a Grant-in-aid for Scientific Research from the Japan Society for the Promotion of Science (C, 23540546, 2011–2013) (to KA). Field work also was supported by the Royal Society of New Zealand's Marsden Fund (06-UOA-082) (to KAC) and a travel fund from the Royal Society (to CTSL).

## References

- Adams, J.H. 1910. The geology of the Whatatutu subdivision, Raukumara division, Poverty Bay. *New Zealand Geological Survey Branch Bulletin, New Series* 9: 1–48.
- Amano, K. and Kiel, S. 2007. Fossil vesicomid bivalves from the North Pacific region. *The Veliger* 49: 270–293.
- Amano, K. and Kiel, S. 2011. Fossil *Adulomya* (Vesicomidae, Bivalvia) from Japan. *The Veliger* 51: 76–90.
- Amano, K. and Kiel, S. 2012. Two Neogene vesicomid species (Bivalvia) from Japan and their biogeographic implications. *The Nautilus* 126:79–85.
- Aoki, S. 1954. Mollusca from the Miocene Kabeya Formation, Joban coalfield, Fukushima Prefecture, Japan. *Science Reports of the Tokyo Kyōiku Daigaku, Section C* 3: 23–41.
- Beu, A.G. and Maxwell, P.A. 1990. Cenozoic Mollusca of New Zealand. *New Zealand Geological Survey Paleontological Bulletin* 58: 1–518.
- Boss, K.J. and Turner, R.D. 1980. The giant white clam from the Galapagos Rift, *Calyptogena magnifica* species novum. *Malacologia* 20: 161–194.
- Campbell, K.A., Francis, D.A., Collins, M., Gregory, M.R., Nelson, C.S., Greinert, J., and Aharon, P. 2008. Hydrocarbon seep-carbonates of a Miocene forearc (East Coast Basin), North Island, New Zealand. *Sedimentary Geology* 204: 83–105.
- Campbell, K.A., Nelson, C.S., Alfaro, A.C., Boyd, S., Greinert, J., Nyman, S., Grosjean, E., Logan, G.A., Gregory, M. R., Cooke S., Linke, P., Mil-

- loy, S. and Wallis, I., 2010. Geological imprint of methane seepage on the seabed and biota of the convergent Hikurangi Margin, New Zealand: Box core and grab carbonate results. *Marine Geology* 272: 285–306.
- Cosel, R. von and Olu, K. 2009. Large Vesicomidae (Mollusca: Bivalvia) from cold seeps in the Gulf of Guinea off the coasts of Gabon, Congo and northern Angola. *Deep-Sea Research II* 56: 2350–2370.
- Dall, W.H. 1891. Scientific results of explorations by the U.S. Fish Commission Steamer Albatross. no. XX. On some new or interesting West American shells obtained from dredgings of the U.S. fish commission steamer Albatross in 1888. *Proceedings of the U.S. National Museum* 14: 174–191.
- Dall, W.H. 1903. Contributions of the Tertiary fauna of Florida with especial reference to the Silex Beds of Tampa and the Pliocene beds of Caloosahatchie River, including in many cases a complete revision of the generic groups treated of and their American Tertiary species. Part VI. Concluding the work. *Transactions of the Wagner Free Institute of Science of Philadelphia* 3: 1219–1654.
- Dall, W.H. and Simpson, C.T. 1901. The Mollusca of Porto Rico. *Bulletin of the United States, Fish and Fisheries Commission* 20: 351–524.
- Dubilier, N., Bergin, C., and Lott, C. 2008. Symbiotic diversity in marine animals: the art of harnessing chemosynthesis. *Nature Reviews Microbiology* 6: 725–740.
- Ewen, S.M. 2009. *Diagenetic Evolution of Some Modern and Ancient Cold Seep Carbonates from East Coast Basin, New Zealand*. 231 pp. Unpublished MSc Thesis, The University of Waikato, Hamilton.
- Field, B.D., Uruski, C.I., Beu, A., Browne, G., Crampton, J., Funnell, R., Killips, S., Laird, M., Mazengarb, C., Morgans, H., Rait, G., Smale, D., and Strong, P. 1997. Cretaceous–Cenozoic geology and petroleum systems of the East Coast Region, New Zealand. *Institute of Geological and Nuclear Sciences Monograph* 19: 1–301.
- Hayward, B.W. 1986. A guide to paleoenvironmental assessment using New Zealand Cenozoic foraminiferal (sic) faunas. *New Zealand Geological Survey Report. Paleontology Group* 109: 1–73.
- Henderson, J. and Ongley, M. 1920. The Geology of the Gisborne and Whatatutu Subdivisions, Raukumara Division. *New Zealand Geological Survey Bulletin* 21: 1–88.
- Kamp, P.J.J. and Nelson, C.S. 1988. Nature and occurrence of modern and Neogene active margin limestones in New Zealand. *New Zealand Journal of Geology and Geophysics* 31: 1–20.
- Kanno, S. 1971. Tertiary molluscan fauna from the Yakataga district and adjacent areas of southern Alaska. *Palaeontological Society of Japan, Special Papers* 10: 1–154.
- Kiel, S. 2010. The fossil record of vent and seep mollusks. In: S. Kiel (ed.), *The Vent and Seep Biota. Topics in Geobiology* 33: 255–277.
- Kiel, S. and Amano, K. 2010. Oligocene and Miocene vesicomid bivalves from the Katalla district in southern Alaska, USA. *The Veliger* 51: 76–84.
- Krylova, E.M. and Cosel, R. von 2011. A new genus of large Vesicomidae (Mollusca, Bivalvia, Vesicomidae, Pliocardiinae) from the Congo margin, with the first record of the subfamily Pliocardiinae in the Bay of Biscay (northeastern Atlantic). *Zoosystema* 33: 83–99.
- Krylova, E.M. and Janssen, R. 2006. Vesicomidae from Edison Seamount (South Western Pacific: Papua New Guinea: New Ireland fore-arc basin) (Bivalvia: Glossoidea). *Archiv für Molluskenkunde* 135: 233–263.
- Krylova, E.M. and Sahling, H. 2006. Recent bivalve molluscs of the genus *Calyptogena* (Vesicomidae). *Journal of Molluscan Studies* 72: 359–395.
- Krylova, E.M. and Sahling, H. 2010. Vesicomidae (Bivalvia): Current taxonomy and distribution. *PLoS ONE* 5: 1–9.
- Kuroda, T. 1931. Fossil Mollusca [in Japanese]. In: F. Honma (ed.), *Geology of the Central Part of Shinano, Part 4*. 90 pp. Kokon Shoin, Tokyo.
- Levin, L.A. 2005. Ecology of cold seep sediments: Interactions of fauna with flow, chemistry and microbes. *Oceanography and Marine Biology: An Annual Review* 43: 1–46.
- Lewis, K.B. and Marshall, B.A. 1996. Seep faunas and other indicators of methane rich dewatering on New Zealand convergent margins. *New Zealand Journal of Geology and Geophysics* 39: 181–200.
- Lillie, A. 1953. The geology of the Dannevirke Subdivision. *New Zealand Geological Survey Bulletin* 46: 1–156.
- Lutaenko, K.A. and Xu, F. 2008. A catalogue of types of bivalve mollusks in the Marine Biological Museum, Chinese Academy of Sciences (Qingdao). *The Bulletin of the Russian Far East Malacological Society* 12: 42–70.
- Mazengarb, C., Francis, D.A., and Moore, P.R. 1991. *Geological Map of New Zealand, 1:50,000, Sheet Y16, Geology of the Tauwhareparae area*. New Zealand Geological Survey, Wellington.
- Mazengarb, C. and Speden, I.G. 2000. *Geology of the Rakumara Area. 1:250,000 geological map 6*. 60 pp. Institute of Geological & Nuclear Sciences, Wellington.
- McKay, A. 1877a. Reports made relative to collections of fossils made in the East Cape District, North Island, New Zealand. *New Zealand Geological Survey, Reports of Geological Explorations during 1873–1874* 8: 116–164.
- McKay, A. 1877b. On the geology of east Auckland and the northern district of Hawke's Bay. *New Zealand Geological Survey, Reports of Geological Explorations during 1886–1887* 18: 183–219.
- Okutani, T., Fujikura, K., and Kojima, S. 2000. New taxa and review of vesicomid bivalves collected from the Northwest Pacific by deep sea research systems of Japan marine Science & Technology. *Venus (Japanese Journal of Malacology)* 59: 83–101.
- Ongley, M. and MacPherson, E.O. 1928. The geology of the Waiapu subdivision, Raukumara division. *New Zealand Geological Survey Bulletin* 30 (New Series): 1–79.
- Saether, K.P. 2011. *A Taxonomic and Palaeobiogeographic Study of the Fossil Fauna of Miocene Hydrocarbon Seep Deposits, North Island, New Zealand*. 479 pp. Unpublished Ph.D. thesis, The University of Auckland, Auckland.
- Saether, K.P., Little, C.T.S., and Campbell, K.A. 2010a. A new fossil provanid gastropod from Miocene hydrocarbon seep deposits, East Coast Basin, North Island, New Zealand. *Acta Palaeontologica Polonica* 55: 507–517.
- Saether, K.P., Little, C.T.S., Campbell, K.A., Marshall, B.A., Collins, M., and Alfaro, A.C. 2010b. New fossil mussels (Mollusca: Bivalvia: Mytilidae) from Miocene hydrocarbon seep deposits, North Island, New Zealand, with general remarks on vent and seep mussels. *Zootaxa* 2577: 1–45.
- Saether, K.P., Little, C.T.S., Marshall, B.A., and Campbell, K.A. 2012. Systematics and palaeoecology of a new fossil limpet (Patellogastropoda: Pectinodontidae) from Miocene hydrocarbon seep deposits, East Coast Basin, North Island, New Zealand with an overview of known fossil seep pectinodontids. *Molluscan Research* 32: 1–15.
- Smith, C.R. and Baco, A.R. 2003. Ecology of whale falls at the deep-sea floor. *Oceanography and Marine Biology: An Annual Review* 41: 311–354.
- Takeda, H. 1953. The Poronai Formation (Oligocene Tertiary) of Hokkaido and South Sakhalin and its fossil fauna. *Studies on Coal Geology, the Hokkaido Association of Coal Mining Technologists* 3: 1–103.
- Taylor, J.D. and Glover, E.A. 2010. Chemosymbiotic bivalves. In: S. Kiel (ed.), *The Vent and Seep Biota. Topics in Geobiology* 33: 107–136.
- Troup, M.J. 2010. *Sedimentology and Petrology of Miocene Cold-seep Carbonates in Southern Hawke's Bay: Geological Evidence for Past Seabed Hydrocarbon Seepage*. 270 pp. Unpublished MSc Thesis, the University of Waikato, Hamilton.
- Woodring W.P. 1925. Miocene mollusks from Bowden, Jamaica. Part I: Pelecypods and Scaphopods. *Carnegie Institution of Washington, Publication* 366: 1–222.
- Xu, F. and Shen, S. 1991. A new species of Vesicomidae from Nansha Islands waters [in Chinese with English abstract]. *Papers on Marine Biology of Nansha Islands and Adjacent Seas* 1: 164–166.