

## **Dispharynxiasis in a Captive Princess Parrot**

Author: Bolette, David P.

Source: Journal of Wildlife Diseases, 34(2) : 390-391

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-34.2.390>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Dispharynxiasis in a Captive Princess Parrot

David P. Bolette, University of Pittsburgh, Laboratory Animal Resources, S1040 Biomedical Science Tower, Pittsburgh, Pennsylvania 15261, USA (e-mail: dbolette@vms.cis.pitt.edu).

**ABSTRACT:** The acuariid nematode *Dispharynx nasuta* was found in a princess parrot, *Polytelis alexandrae*, at the National Aviary in Pittsburgh (Pennsylvania, USA). This is the first report of *D. nasuta* from the host order Psittaciformes, and was the presumed cause of death in this parrot.

**Key words:** Case report, *Dispharynx nasuta*, new host, *Polytelis alexandrae*, princess parrot, record.

An adult male princess parrot, *Polytelis alexandrae* (Aves: Psittaciformes: Psittacidae) was received at the National Aviary in Pittsburgh (Allegheny County, Pennsylvania, USA) from a private collector on 17 September 1991. Approximately 16 mo after its arrival, the bird began to exhibit signs of debilitation and wasting thought to be due to a confirmed gastrointestinal helminth infection. Numerous anthelmintic regimes were implemented over a period of time, but all proved unsuccessful. The bird's clinical condition continued to deteriorate. It was unresponsive to therapeutic intervention, and after approximately 90 days from the onset of clinical signs, the bird became severely debilitated and died on 27 April 1993. This host specimen was deposited in the Carnegie Museum of Natural History (Section of Birds, Pittsburgh, Pennsylvania; catalog number S-15071, accession number 35664).

Necropsy revealed large numbers of a single species of acuariid nematode firmly attached to the proventricular mucosa. Twelve specimens were provided for preliminary identification and assigned to the genus *Dispharynx*. Subsequent to this generic assignment, the specimens (four male, eight female) were specifically identified as *D. nasuta* (Nematoda: Acuariidae). All remaining specimens were formalin fixed in situ, and embedded in paraffin for histological sectioning. Nematodiasis was the only significant post-mortem

finding. All other organs were normal, on gross and histological examination. Additionally, there was no evidence of gastric inflammation, papillomatous proliferation, or necrosis which is often associated with severe infections of *D. nasuta*. Despite this, the large number of parasites may have been sufficient to have caused nutritional compromise, debilitation, and subsequent death of this host.

Helminths were placed in 10% buffered formalin and stored in 70% ethyl alcohol. Specimens were cleared by the ethyl alcohol-glycerin evaporation technique, examined as temporary slide mounts in glycerin, and deposited in the United States National Parasite Collection (USDA, Beltsville, Maryland, USA; accession number 87067).

*Dispharynx nasuta* has been described in numerous avian hosts; these are primarily from the Columbiformes, Galliformes and Passeriformes (Goble and Kutz, 1945; Baruš and Garrido, 1968; Baruš, 1969; Rickard, 1985; Silva et al., 1990). Additional reports of *D. nasuta* from other avian orders include isolated records from the Anseriformes (Wang and Liew, 1991), Charadriiformes (Eckman, 1968), Ciconiiformes (Gupta and Kumar, 1978), Cuculiformes (Macko et al., 1981), Falconiformes (Gupta and Kumar, 1978; Samedov, 1978), Gruiformes (Forrester et al., 1974, 1975), and Piciformes (Baruš, 1971; Bolette, 1998). The present report of *D. nasuta* from *P. alexandrae* is the first from the host order Psittaciformes.

This infection of *D. nasuta* in *P. alexandrae* occurred opportunistically within a captive situation, and was the presumed cause of death in this host. Morbidity and (or) mortality attributed to *D. nasuta* infections have been previously reported from hosts of the Columbiformes (Hwang

et al., 1961; Lindquist and Straffuss, 1980); and from the Passeriformes and Galliformes (Allen, 1924; Cram, 1928; Goble and Kutz, 1945; Jensen, 1962).

I acknowledge the National Aviary in Pittsburgh for providing nematode specimens; J. Bonner for providing clinical information; and E. Klein for providing histopathologic expertise.

#### LITERATURE CITED

- ALLEN, A. A. 1924. The grouse disease. American Game Bulletin 13: 12–14.
- BARUŠ, V. 1969. Resumen sobre la fauna de los nematodos en las aves del orden Galliformes en Cuba. Torreia 5: 1–9.
- . 1971. A survey of parasitic nematodes of piciform birds in Cuba. Folia Parasitologica 18: 315–321.
- , AND O. H. GARRIDO. 1968. Nematodes parasitic in birds of the order Passeriformes in Cuba. Folia Parasitologica 15: 147–160.
- BOLETTE, D. P. 1998. Gastrointestinal helminths of some yellow-shafted flickers, *Colaptes auratus luteus* (Aves: Picidae), from Allegheny County, Pennsylvania. Journal of the Helminthological Society of Washington 65: 114–116.
- CRAM, E. B. 1928. Nematodes of pathological significance found in some economically important birds in North America. United States Department of Agriculture Technical Bulletin 49: 1–10.
- ECKMAN, M. K. 1968. Helminth parasites of the killdeer in Colorado. The Journal of Parasitology 54: 1143.
- FORRESTER, D. J., A. O. BUSH, AND L. E. WILLIAMS, JR. 1975. Parasites of the Florida sandhill cranes. *Grus canadensis patensis*. The Journal of Parasitology 61: 547–548.
- , ———, ———, AND D. J. WEINER. 1974. Parasites of greater sandhill cranes (*Grus canadensis tabida*) on their wintering grounds in Florida. Proceedings of the Helminthological Society of Washington 41: 55–59.
- GOBLE, F. C., AND H. L. KUTZ. 1945. The genus *Dispharynx* (Nematoda: Acuariidae) in galliform and passeriform birds. The Journal of Parasitology 31: 323–331.
- GUPTA, S. P., AND P. KUMAR. 1978. Studies on some nematode parasites of birds from Uttar Pradesh. Indian Journal of Helminthology 28: 86–109.
- HWANG, J. C., N. TOLGAY, W. T. SHALKOP, AND D. S. JACQUETTE. 1961. *Dispharynx nasuta* causing severe proventriculitis in pigeons. Avian Disease 5: 60–65.
- JENSEN, D. N. 1962. The pathological effects of infections of *Dispharynx nasuta* (Nematoda: Spiruroidea) on the blue grouse *Dendragapus obscurus* (Say). Ph.D. dissertation, University of British Columbia, Vancouver, Canada, 113 pp.
- LINDQUIST, W. D., AND A. C. STRAFUSS. 1980. (*Dispharynx nasuta*) may cycle within avian zoo populations. Journal of Zoo Animal Medicine 11: 120–122.
- MACKO, J. R., V. BIROVA, AND L. ESPAINE. 1981. Epizootiologia de *Dispharynx nasuta* (Rudolphi, 1819) en Cuba. 1. Binomia. Academia de Ciencias de Cuba Informe Científico-Técnico 172: 1–80.
- RICKARD, L. G. 1985. Proventricular lesions associated with natural and experimental infections of *Dispharynx nasuta* (Nematoda: Acuariidae). Canadian Journal of Zoology 63: 2663–2668.
- SAMEDOV, G. A. 1978. Nematodes of predatory birds of Azerbaidjan. Izvestiya Akademii Nauk Azerbaidzhanskoi SSR Seriya Biologicheskikh Nauk 1978: 70–73.
- SILVA, C. C. D., D. G. D. MATTOS JR., AND P. M. RAMIRES. 1990. Helminth parasites of *Columba livia* (Gm) in Sao Goncalo, Rio de Janeiro, Brazil. Arquivo Brasileiro de Medicina Veterinaria e Zootecnia 42: 391–394.
- WANG, J. S., AND S. M. LIEW. 1991. A survey on the parasitic infections in ducks. Journal of the Chinese Society of Veterinary Science 17: 117–123.

Received for publication 24 July 1997.