

First Record of *Alcathoe carolinensis* (Lepidoptera: Sesiidae) Collected in Tennessee

Authors: Hansen, Jason A., Klingeman, William E., and Moulton, John K.

Source: Florida Entomologist, 93(2) : 321-322

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.093.0230>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

FIRST RECORD OF *ALCATHOE CAROLINENSIS* (LEPIDOPTERA: SESIIDAE) COLLECTED IN TENNESSEE

JASON A. HANSEN¹, WILLIAM E. KLINGEMAN² AND JOHN K. MOULTON¹

¹Department of Entomology and Plant Pathology, University of Tennessee, Knoxville, TN 37996

²Department of Plant Sciences, University of Tennessee, Knoxville, TN 37996

Known mostly from male specimens, *Alcathoe carolinensis* Engelhardt (Fig. 1) has been reported as rare, but is more likely to be infrequently collected (Thomas D. Eichlin, Senior Insect Biosystematist, retired, California Department of Food and Agriculture, personal communication), thus the full extent of its native range is poorly documented. Though most captures have been incidental and typically consist of 1 to 3 specimens, only 2 studies have reported captures of 10 or more males through use of *E,Z*-3,13-ODDA and *Z,Z*-3,13-ODDA blends (Reed et al. 1981; Snow et al. 1985). Based on other *Alcathoe* species host plant preferences, larval host plants are assumed to be *Clematis* spp., though *A. carolinensis* remains the only North American member of its genus not reared from any species or cultivar of *Clematis* (Engelhardt 1925; Eichlin & Duckworth 1988). *Alcathoe carolinensis* was once listed as a

synonym of *A. autumnalis* Engelhardt, but later the two were recognized as distinct species (Eichlin & Duckworth 1977; Eichlin & Duckworth 1988). When questioned about the lack of label data on the type specimen, Beutenmüller recalled collecting it on *Clematis* flowers somewhere in “the Black Mountains of North Carolina” (Engelhardt 1946). Engelhardt (1946) reported that subsequent visits to the collecting site showed Beutenmüller’s plant identification to be inaccurate, though he did not clarify what the mistakenly identified plant was. Morphological similarities between *A. carolinensis* and 2 species in the western U.S., *A. pepsoides* Engelhardt and *A. autumnalis*, also cast some doubt on the capture of *A. carolinensis* in a state so disjunct from other similar *Alcathoe* populations (Engelhardt 1946, Eichlin & Duckworth 1977, 1988). Currently, *A. carolinensis* specimens have been collected as far north as Indiana and south to Florida (Sharp et al. 1978; Reed et al. 1981). A lone male captured in Missouri extended the western boundary of its known range and is the most recent reported capture of this species (Brown 1986).

In 2007, as part of an on-going survey of clearwing moth presence in eastern Tennessee, a Multipher-1 moth trap (Les Services BioContrôle, Ste.-Foy, Quebec) was baited with a commercial yellow-legged clearwing moth (*Synanthedon vespiformis* (L.)) lure (Great Lakes IPM, Vestaburg, MI). This modified trap was placed just outside the boundaries of the Great Smoky Mountains National Park in Sevier Co., Tennessee. The trap was retrofitted with an ethanol collection chamber, thus preserving DNA for analyses and preventing damage to important morphological characters. Specimens were captured about 70 meters from a mountain stream in a wooded area on a western-facing slope approximately 600 meters above sea level. Several hemlock, pine, oak trees, and rhododendron shrubs had recently been removed from the site. The lure-baited trap was placed on the edge of this canopy opening, where a single male was collected between 29-VI and 5-VII-2007.

In 2009, the same modified trap style was deployed with 2 viburnum borer (*S. viburni* Engelhardt) lures (Scentry Biologicals, Inc., Billings, Montana) along the wooded edge of a roadside park in Oak Ridge, Tennessee approximately 100 kilometers west of the original Norton Creek site and at



Fig. 1. Adult male *Alcathoe carolinensis* Engelhardt, illustrating the hyaline area at base of hind wing (h) and caudal appendage (ca), which are characteristic of *Alcathoe* males.

about 260 meters in elevation. Canopy mid- and overstory consisted predominantly of oaks (*Quercus* sp.), loblolly pine (*Pinus taeda*), bush honeysuckle (*Lonicera maaackii*), privet (*Ligustrum sinense*), tulip poplar (*Liriodendron tulipifera*), sweetgum (*Liquidambar styraciflua*), and rusty blackhaw viburnum (*V. rufidulum*). The trap yielded 8 *A. carolinensis* males between 10 and 20-VIII-2009. Identification of the sesiid was verified by Thomas D. Eichlin. Although *Clematis* species were not found within about 160 meters of deployed traps, several other vining forbs were found within 33 meters, including honeysuckle (*Lonicera* sp.), trumpet creeper (*Campsis radicans*), greenbriar (*Smilax* sp.), grape (*Vitis* sp.), and Carolina moonseed (*Cocculus carolina*).

In Tennessee, *A. carolinensis* responded to lures which attract *S. viburni* and *S. vespiformis*, both known to be drawn to Z,Z-3,13-octadecadienyl acetate (ODDA)/E,Z-3,13-octadecadienyl acetate at a ratio of 9:1 (Greenfield & Karandinos 1979; Voerman et al. 1983). The commercial lures used were confirmed by vendors as containing the same ratio of isomers reported in the literature. With the exception of a solitary account in which E,Z-3,13-ODDA alone was used, previous *A. carolinensis* captures were accomplished with a 50:50 or 75:25 blend of the 2 previously mentioned isomers (Reed et al. 1981; Snow et al. 1985; Sharp et al. 1978; Brown 1985, 1986). Regardless of the exact isomer blend used, we expect *A. carolinensis* will continue to be infrequently collected until larval host plant resources are identified and trapping is focused around habitats containing key plant species.

Although larvae of other *Alcathoe* species rely on *Clematis* plants for development, no specimens of the plant genus were found at the Oak Ridge, Tennessee location. A less common species, *C. glaucophylla*, is listed on Tennessee's rare plant list as endangered (Crabtree 2008) and is reported from North Carolina, Florida, Georgia, and eastern Tennessee. However, older reports of this clematis species may have confused it with closely-related *C. viorna*, thus the range of *C. glaucophylla* may be more restricted than is reported (Estes 2006).

Because many ornamental *Clematis* species are economically important and popular landscape specimens, further efforts to find larvae and rear adults from *Clematis* species are warranted. Engelhardt (1925) noted use of horticultural *Clematis* varieties by *A. caudata* (Harris), but mentioned no specific varieties. We caution that at least 1 native *Clematis* species, *C. glaucophylla* is endangered in Tennessee, thus care should be taken to assess any protected status this plant may have when sampling plants to confirm *A. carolinensis* presence from other localities in the state.

SUMMARY

The first captures of *Alcathoe carolinensis* Engelhardt in Tennessee are reported from phero-

mone-baited trap yields taken in 2007 and 2009 from eastern Tennessee locations 100 kilometers apart and at different elevations. Traps were baited with a different pheromone combination than reported in other published accounts. Its capture at 600 meters elevation in the Great Smoky Mountains adjacent to GSM National Park boundaries marks the highest elevation at which *A. carolinensis* has been recorded and is similar to the original type locality described by Beutenmüller for North Carolina.

REFERENCES CITED

- BROWN, L. N. 1985. New records of the rare clearwing moth, *Alcathoe carolinensis* Engelhardt, (Sesiidae) in Florida. Florida Entomol. 68(4): 700-701.
- BROWN, L. N. 1986. First record of the rare clearwing moth, *Alcathoe carolinensis* Engelhardt (Lepidoptera: Sesiidae) west of the Mississippi river. J. Kansas Entomol. Soc. 59(3): 560.
- CRABTREE, T. 2008. Tennessee Natural Heritage Program Rare Plant List. Tennessee Department of Environment and Conservation, Division of Natural Areas. 46 p.
- DUCKWORTH, W. D., AND EICHLIN, T. D. 1977. A Classification of the Sesiidae of America North of Mexico (Lepidoptera: Sesiioidea). California Department of Food and Agriculture, Occasional Papers in Entomol. 2: 1-54.
- EICHLIN, T. D., AND DUCKWORTH, W. D. 1988. The moths of America North of Mexico, Fascicle 5.1, Sesiioidea, Sesiidae. The Wedge Entomol. Res. Foundation, Washington, D.C., 176 p., illustr.
- ENGELHARDT, G. P. 1925. Studies of North American Aegeriidae (Lepidoptera). Bull. Brooklyn Entomol. Soc. 20(4): 153-158.
- ENGELHARDT, G. P. 1946. The North American clearwing moths of the family Aegeriidae. Smithsonian Institution, U.S. Natl. Mus. Bull. 190, 222 p.
- ESTES, D. 2006. A new narrowly endemic species of *Clematis* (Ranunculaceae: Subgenus *Viorna* from northeastern Texas. J. Bot. Res. Inst. Texas. 22(1): 65-77.
- GREENFIELD, M. D., AND KARANDINOS, M. G. 1979. Resource partitioning of the sex communication channel in clearwing moths (Lepidoptera: Sesiidae) of Wisconsin. Ecol. Monographs. 49: 403-426.
- REED, D. K., EICHLIN, T. D., AND REED, G. L. 1981. Effectiveness of blends of synthetic sex attractants and comparison with virgin female lesser peachtree borers as bait for capture of Sesiidae. Environ. Entomol. 10: 488-491.
- SHARP, J. L., MCLAUGHLIN, J. R., JAMES, J., EICHLIN, T. D., AND TUMLINSON, J. H. 1978. Seasonal occurrence of male Sesiidae in north central Florida determined with pheromone trapping methods. Florida Entomol. 61(4): 245-250.
- SNOW, J. W., EICHLIN, T. D., AND TUMLINSON, J. H. 1985. Seasonal captures of clearwing moths (Sesiidae) in traps baited with various formulations of 3, 13-Octadecadienyl acetate and alcohol. J. Agr. Entomol. 2: 73-84.
- VOERMAN, S., AUDEMARD, H. AND PRIESNER, E. 1983. Sex attractants for clearwing moths: *Synanthedon vespiformis* and *Chamaesphecia tenthrediniformis* (and/or *C. empiformis*). Ent. Exp. Appl. 34: 203-205.