

A New Species of Aeolothrips (Thysanoptera: Aeolothripidae) from Mango Crops in Oaxaca, Mexico

Authors: Cruz, Javier Ruiz-De La, Vásquez-López, Alfonso, Retana-Salazar, Axel P., Mora-Aguilera, José Antonio, and Johansen-Naime, Roberto

Source: Florida Entomologist, 96(1) : 29-35

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0104>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A NEW SPECIES OF *AEOLOTHRIPS* (THYSANOPTERA: AEOLOTHRIPIDAE) FROM MANGO CROPS IN OAXACA, MEXICO

JAVIER RUIZ-DE LA CRUZ¹, ALFONSO VÁSQUEZ-LÓPEZ¹, AXEL P. RETANA-SALAZAR², JOSÉ ANTONIO MORA-AGUILERA³
AND ROBERTO JOHANSEN-NAIME⁴

¹Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional (CIIDIR), Unidad Oaxaca, Instituto Politécnico Nacional. Calle Hornos 1003. 71230, Santa Cruz Xoxocotlán, Oaxaca, México

²Programa de Especial Interés Institucional en Biología Aplicada (PUA), Centro de Investigación en Estructuras Microscópicas (CIEMIC), Ciudad de la Investigación, Universidad de Costa Rica

³Colegio de Postgraduados (CP), Instituto de Fitosanidad, km 36.5, Carr. México-Texcoco, Montecillo, Edo. de México, México 56230

⁴Instituto de Biología, Universidad Nacional Autónoma de México (IBUNAM), Coyoacán, México DF

*Corresponding author: E-mail: ruizj777@gmail.com

ABSTRACT

We describe *Aeolothrips romanruizi* **sp. nova** that has been recently discovered in mango orchards in the Isthmus region of Oaxaca in southern Mexico. *Aeolothrips romanruizi* **sp. nova** exhibits an ornamentation of the mesonotum and metanotum very different from others in this genus, except *A. microstriatus*, which is similarly ornamented; but these 2 species differ in forewing color pattern, body size and some other characters of chaetotaxy. A key to the *Aeolothrips* species of Central America and Mexico is provided.

Key Words: fruit, *Aeolothrips microstriatus*, predator, SEM

RESUMEN

Se describe una nueva especie del género *Aeolothrips* que ha sido recolectado en el cultivo de mango en la región del Istmo de Oaxaca en el sur de México. En las especies descritas del género *Aeolothrips* la ornamentación del meso y metanoto es completamente distinto de la descrita aquí, solo *A. microstriatus* es similar y la nueva especie difiere de esta en el patrón de color del ala anterior, el tamaño corporal y otros caracteres referentes a la quetotaxia. Se anexa una clave para las especies de *Aeolothrips* de Centro América y México.

Palabras Clave: fruta, *Aeolothrips microstriatus*, depredador, SEM

Insect predators are one of the most important agents in insect pest management practices, especially in those scenarios where there are few known parasitoids to control microarthropod pests (Sánchez-Ruiz et al. 1997). Predatory thrips species of Thysanoptera: Aeolothripidae have been studied as potential biological control agents. For example, *Franklinothrips orizabensis* Johansen 1974 has received much attention as a possible biological control agent of thrips species that are harmful to crops (Hoddle 2003b).

Members of Aeolothripidae are medium-sized, usually with dark brown bodies about 2.5mm long (Mound & Marullo 1996). Adults and larvae of many species in this family appear to be facultative predators of other small arthropods, in that they feed on both floral tissues as well as on thrips and mites that live in flowers. Some species are almost certainly exclusively phytophagous (Tyagi et al. 2008), but in the warmer parts

of the world, a considerable number of species are predators (Hoddle 2003a). Worldwide, about 250 species are recognized in 26 genera of Aeolothripidae (Mound & Marullo 1996).

Studies in avocado groves in search of natural enemies of *Scirtothrips perseae* Nakahara 1997 have identified predator species in *Aeolothrips*, *Aleurodothrips*, *Franklinothrips*, *Leptothrips*, *Scolothrips*, and *Karnyothrips* as possible biological control agents (Hoddle et al. 2002; Cambero-Campos et al. 2011) for this pest.

Here we describe a new species of the genus *Aeolothrips* that has recently been discovered in mango crops in the Isthmus region of Oaxaca in southern Mexico. This predatory thrips may be important in controlling many species of phytophagous thrips that attack this crop.

Aeolothrips romanruizi **sp. nova.**

MATERIAL

HOLOTYPE ♀. MEXICO, Oaxaca, San Pedro Tapanatepec, N 16°27'18.5" W 94°13'23.5", 46 m asl. Paratypes: 14 ♀♀, same data the holotype. Holotype and 4 paratypes deposited in the collection IBUNAM of the Universidad Autonoma de Mexico, 5 paratypes deposited in the collection of the Universidad de Costa Rica and 5 paratypes deposited in the Centro Interdisciplinario de In-

vestigación para el Desarrollo Integral Regional (CIIDIR), IPN-Unidad Oaxaca.

Color. Body mostly dark brown. Legs with dark brown femora and tibiae. Wings pale translucent with two brown bands longer than wide (Fig. 1A). Major setae dark brown.

Antennal segments I-IX dark brown, except segment III (and extreme base of segment IV) which is completely yellow (Fig. 1B). Antennal segments III-V longer than wide and cylindrical,




Fig. 1. Phase contrast photographs of *Aeolothrips romanruizi* sp. nova. A) Ventral view of the body, B) Antenna, C) Dorsal view of the head, and D) Mouth cone and maxillary palpi well developed, with 3 segments.

with parallel sides and with several rows of microtrichia. Segments VI-IX forming a style, sensoria on III-IV well developed, linear and longer than half the length of the segment (Fig. 3A)

Head. As long as wide, with many randomly-placed, long, thin setae in the postocular region. Ocular setae short. Back of the head with widely-spaced striations. Well-developed eyes with multiple ommatidia, the eye is longer in the ventral face than on the dorsal face (Figs. 1C and 3B). Mouth cone long and strong, maxillary palpi well developed, with 3 segments (Fig. 1D).

Prothorax. Wider than long, with many discal setae, anteromarginals setae reduced, posteromarginal setae stronger and longer than the oth-

er setae on the pronotum. Posteromarginal setae I more developed than posteromarginal II and III. Discal region without ornamentation with several discal setae shorter than posteromarginals I and II (Figs. 2A and 3B). Fema entire.

Pterothorax. Meso and metanotum with many closely spaced striations (Fig. 2B). Ventral region with many discal setae on the meso- and metasternum (Figs. 2C and 3C).

Wings. Broad with transverse veins evident and well developed between the longitudinal veins (Figs. 2D and 3D).

Abdomen. Terga smooth, sterna IV-VI without discal setae (Fig. 3D). Ovipositor well developed with many very small teeth (Fig. 3C).




Fig. 2. Phase contrast photographs of *Aeolothrips romanruizi* **sp. nova**. A) Prothorax, B) Pterothorax, mesonotum and metanotum with many closely spaced striations, C) Mesosternum and metasternum with many discal setae, and D) Wing, pale translucent with transverse veins evident and well developed between the longitudinal veins.




Fig. 3. Scanning electron micrographs of *Aeolothrips romanruizi* sp. nova. A) Antennal segments with parallel sides and with several rows of microtrichias, B) Head and Prothorax. Well-developed eyes with multiple ommatidia, ocellar setae short. Pronotum with anteromarginals setae reduced, C) Ventral view of the female, D) Dorsal view of the female.

Sizes in μm of morphological traits of the holotype: total body length= 1850, total length of the antenna = 372.5; I = 30, II = 55, III = 95, IV = 92.5, V = 57.5, VI-VIII = 10, IX = 12.5, pedicel length of antennal segments III and IV, 7.5 and

2.5 respectively; am = 15, amm = 15, PMI = 25, PMII = 20, PMIII = 15, = 430 ovipositor.

Male

Unknown.

Etymology

The species name is dedicated to the memory of Roman Ruiz Sánchez in recognition of his dedication and hard work as an evangelical pastor.

DISCUSSION

Specimens of *Aeolothrips romanruizi* **sp. nova** collected in Oaxaca possess the following characters (those in italics are considered diagnostic of *Aeolothrips* by Mound & Marullo (1996)): antenna with 9 segments, sensory in segments III-IV linear VI-IX short and condensed to form a unit, compound eyes prolonged ventrally, *maxillary palpi with 3 segments*, head and pronotum without setae, posteromedial pair of setae on the metanotum, *forewing with 1 or 2 bands of dark color with clear apex*, *abdominal sternites IV-VI with setae and without discal setae marginal or ancillary*, sternite VII with 2 pairs of accessory setae submarginal.

The previously described species of *Aeolothrips* have ornamentation of the mesonotum and metanotum different from that of *A. romanruizi* **sp. nova**. In most species the ornamentation of the mesonotum is striated and with widely spaced lines, and usually the metanotum is reticulated. The only described species showing ornamentation similar to that of *A. romanruizi* **sp. nova** is *microstriatus* (Hood 1935). *Aeolothrips romanruizi* **sp. nova** is known only from the holotype, from a male collected in Panama, and from a specimen collected in Brazil by an anonymous reviewer that has not been described because of insufficient material. Hood (1935) stated that

the species, *A. microstriatus*, differs from others by having a bicolored body, especially in the abdomen, antennae with light brown segments, a single band on the anterior part of the wing I, and unique ornamentation of the pterothorax. The new species *A. romanruizi* **sp. nova** has an entirely dark brown body (Fig. 1A), the segments of the antenna are dark brown with the exception of III, which is bright yellow (Fig. 1B). The wing has 2 dark brown bands and the many striations on the mesonotum and metanotum of the pterothorax with many closely spaced together (Fig. 2D), similar to *microstriatus*.

Color pattern in *Aeolothrips*

The color pattern in many groups of thrips is critical in separating species, as evidenced by keys developed by multiple authors (zur Strassen 1997). Recently it has been shown that in some groups like in *Frankliniella* some color patterns may be of phylogenetic importance and may be useful in determining groups of species (Retana-Salazar 2010). The key of Mound & Marullo (1996) is still the most complete for the determination the New World *Aeolothrips* species. In this key 60% of the items involve the use the color pattern in the separation of species, and this is valid for both males and females.

A review of the main *Aeolothrips* species found in America (Table 1) indicates that the greatest variation in color pattern between males and females is manifested in the coloration of the antennomeres, especially in the coloration of antennomere III. In some species antennomere III is

TABLE 1. COMPARISON BETWEEN THE PRINCIPAL CHARACTERS RELATING TO SEXUAL DIMORPHISMS IN THE MOST ABUNDANT *AEOLOTHrips* SPECIES IN AMERICA, NAMELY, THE COLOR PATTERNS OF THE BODY, ANTENNAE AND WINGS, AS WELL AS CLASPERS, SETAE AND TUBERCLES IN MALES.

Species	Claspers in male at IX	Body (Color)			Antennomere III color		Color of Wings
		♀ compared to ♂	Setae in male	Tubercles in male	♀ compared to ♂	♀ compared to ♂	
<i>Aeolothrips fasciatus</i>	✓	Similar	X	✓	♂ Darker	Similar	
<i>Aeolothrips intermedius</i>	✓	Similar	—	—	Similar	Similar	
<i>Aeolothrips albicinctus</i>	X	Similar	—	—	Similar	Similar	
<i>Aeolothrips vittipennis</i>	X	Similar	X	X	♂ Darker	Similar	
<i>Aeolothrips melaleucus</i>	✓	Similar	—	✓	Similar	Similar	
<i>Aeolothrips fuscus</i>	✓	Similar	X	X	♂ Darker	Similar	
<i>Aeolothrips kuwanaii</i>	✓	Similar	✓	✓	♂ Darker	Similar	
<i>Aeolothrips crucifer</i>	✓	Similar	✓	✓	Similar	Similar	
<i>Aeolothrips hartleyi</i>	✓	Similar	X	X	♂ Darker	Similar	
<i>Aeolothrips hesperus</i>	✓	Similar	X	X	Similar	Similar	
<i>Aeolothrips collaris</i>	✓	Similar	✓	✓	Similar	Similar	
<i>Aeolothrips auricestus</i>	✓	Similar	X	X	Similar	Similar	
<i>Aeolothrips brunneipictus</i>	Condition of material is not adequate to make reliable assessments						

✓ = Present, X = Absent, and — = No data.

REFERENCES CITED

- CAMBERO-CAMPOS, O. J., JOHANSEN-NAIME, R., GARCÍA-MARTÍNEZ, O., CANTU-SIFUENTES, M., CERNA-CHÁVEZ, E., AND RETANA-SALAZAR, A. P. 2011. Especies depredadoras de trips (Thysanoptera) asociadas a huertas de aguacate en Nayarit, México. *Acta Zoológica Mexicana* (n.s.) 27(1): 115-121.
- HODDLE, M. S. 2003a. Predation behaviours of *Franklinothrips orizabensis* (Thysanoptera: Aeolothripidae) towards *Scirtothrips perseae* and *Heliothrips haemorrhoidalis* (Thysanoptera: Thripidae). *Biological Control*, 27: 323-328.
- HODDLE, M. S. 2003b. The effect of prey species and environmental complexity on the functional response of *Franklinothrips orizabensis*: a test of the fractal foraging model. *Ecol. Entomol.* 28: 309-318.
- HODDLE, M. S., NAKAHARA, S., AND PHILLIPS, A. P. 2002. Foreign exploration for *Scirtothrips perseae* Nakahara (Thysanoptera: Thripidae) and associated natural enemies on avocado (*Persea americana* Miller). *Biological Control* 24(3): 251-265.
- HOOD, D. J. 1935. Five new thysanoptera of the genus *Aeolothrips* (Aeolothripidae). *Trans. Entomol. Soc.* 61(3): 103-111.
- MOUND, L. A., AND MARULLO, R. 1996. The Thrips of Central and South America: An Introduction. *Memoirs on Entomology, International* 6: 1-487.
- SÁNCHEZ-RUIZ, M., FONTAL-CAZALLA, F. M., SÁNCHEZ-RUIZ, A., AND LÓPEZ-COLÓN, J. I. 1997. El uso de insectos depredadores en el control biológico aplicado. *Boletín de la Sociedad Entomológica Aragonesa* 20: 141-149.
- STRASSEN, ZUR, R. 1997. How to classify the species of the genus *Thrips* (Thysanoptera)? *Folia Entomol. Hungarica* 58: 227-235.
- TYAGI, K., KUMAR, V., AND MOUND, L. A. 2008. Sexual dimorphism among Thysanoptera Terebrantia, with a new species from Malaysia and remarkable species from India in Aeolothripidae and Thripidae. *Insect Syst. Evol.* 39: 155-170.