

First Record of *Chalcodemus bicolor* (Coleoptera: Curculionidae) Enemies in South America

Authors: Cedeño, P. Emilio, Anjos, N., and Costa, V. A.

Source: Florida Entomologist, 100(1) : 211-212

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0138>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First record of *Chalcodermus bicolor* (Coleoptera: Curculionidae) enemies in South America

P. Emilio Cedeño^{1,2,*}, N. Anjos¹, and V. A. Costa³

Weevils (Coleoptera: Curculionidae) are an ecologically and economically important group of eucalypt pests, and outbreaks have increased in the recent years, mainly of species in the subfamily Molitinae (Garlet et al. 2011; Souza 2011; Cedeño & Flowers 2012; Sweeney et al. 2012; Mafia et al. 2013). Damage by Molitinae weevils in general includes defoliation and chewing holes and pruning the apical sprout and lateral branches, which affects tree growth and wood quality.

The Brazilian eucalypt pruner, *Chalcodermus bicolor* Fiedler (Coleoptera: Curculionidae), causes high levels of damage to the Brazilian eucalypt plantations (Souza et al. 2011). After the female has cut through main and lateral tree sprouts, further damage is caused when the female oviposits in the branch tips.

Chemical insecticide treatment is required to control this species, but pesticides cannot be used by eucalypt producers because many forest enterprises are certified by international forest regulations (e.g., Forest Stewardship Council), which restrict pesticide use. The aim of this work was to find natural enemies affecting *C. bicolor* immature stages.


This study was carried out in Mucuri County, in the southern region of Bahia State, Brazil. To survey for natural enemies of the eucalypt pruner, we collected 440 tips from hybrid *Eucalyptus urophylla* × *E. grandis* (Myrtaceae) trees that had been pruned by the weevils. Samples from the approximately 1 ha site were collected in 2011 on 24 Feb, 4 Mar, and 19 Mar. In the laboratory, branch tips were stored in 18 cm Petri dishes until parasitoids emerged, which were collected daily and stored in 70% ethanol. Specimens were identified by V. A. Costa.

Photos were taken using a Leica M165C stereomicroscope equipped with a Leica DFC420 camera and the Leica Application Suite software.

We obtained 257 adult parasitoids from the collected tips. Among these insects, we could identify 3 morphospecies in the genus *Euderus* Haliday (Hymenoptera: Eulophidae) (Figs. 1 and 2); these morphospecies represented 98.4% ($n = 253$) of all parasitoids that emerged. Four specimens were identified as *Eurydinoteloides* sp. Girault (Hymenoptera: Pteromalidae) (Fig. 3). Specimens were deposited in the Biological Institute Museum, São Paulo County, São Paulo State, Brazil. Percentage of adult parasitoid emergence from the sampled branch tips were: 63.2% ($n = 120$; 24 Feb), 57.0% ($n = 87$; 4 Mar), and 50.0% ($n = 50$; 19 Mar), which suggests an estimate of the natural parasitism level in *C. bicolor* populations of 56.7%, mainly caused by the 3 *Euderus* species.

Previous reports of *Euderus* parasitism of Curculionidae species include *Cylindrocopturus eatoni* Buchanan and *Ceutorhynchus rapae* (Gyllenhal) (Yoshimoto 1971); *Ceutorhynchus obstrictus* (Marshall) (Doddall et al. 2009; Mason et al. 2011); *Cylas formicarius* (F.) (Jansson & Lecrone 1991), and *Goniapterus scutellatus* Gyllenhal (Loch 2008). Similarly, *Eurydinoteloides* species attack various Curculionidae species, including *Eutinobothrus brasiliensis* (Hambleton) (Hambleton & Sauer 1938; Monte 1944; Sauer 1946), species of *Chalcodermus* Schönherr, and 10 additional genera (Noyes 2015).

In summary, we report for the first time that 3 *Euderus* species and 1 species in the genus *Eurydinoteloides* are acting as natural enemies of *C. bicolor* in Brazilian eucalypt plantations. We suggest that *Euderus* species should be studied further as candidates for biological control



Figs. 1–3. Hymenoptera natural enemies of *Chalcodermus bicolor*. **1 and 2:** *Euderus* spp. (Eulophidae), scale bars = 1.33 mm; and **3:** *Eurydinoteloides* sp. (Pteromalidae), scale bar = 1.40 mm.

¹Laboratório de Manejo Integrado de Insetos Florestais/Casa dos Cupins, Departamento de Entomologia, Universidade Federal de Viçosa—UFV, 36570-900 Viçosa, Minas Gerais, Brazil; E-mail: emilioloja@hotmail.com (P. E. C.), nanjos@ufv.br (N. A.)

²Universidad Técnica Estatal de Quevedo, Av. Quito Km 1.5 vía Santo Domingo de los Tsáchilas, Quevedo, Los Ríos, Ecuador. ZIP. 73

³Instituto Biológico/APTA, 13092-543, Campinas, SP-Brasil; E-mail: valmircosta@gmail.com (V. A. C.)

*Corresponding author; E-mail: emilioloja@hotmail.com (P. E. C.)

programs against the eucalypt pruner. Mass rearing and inundative release of adults of *Euderus* may be a useful strategy against this important weevil in the Brazilian eucalypt plantations.

We thank Suzano Papel e Celulose for allowing access to eucalypt plantations to carry out this study.

Summary

Chalcodermus bicolor Fiedler (Coleoptera: Curculionidae) is the most recent pest in Brazilian *Eucalyptus* (Myrtaceae) plantations. Sampling in 2011 for parasitoids of this weevil recovered 4 Hymenoptera species, 3 in the genus *Euderus* (Eulophidae), comprising 98.4% of specimens, and *Eurydinoteloides* sp. (Pteromalidae). This is the first report of natural enemies of *C. bicolor*.

Key Words: forest entomology; eucalypt pruner; biological control; *Euderus*; *Eurydinoteloides*

Sumario

Chalcodermus bicolor Fiedler (Coleoptera: Curculionidae) es la más reciente plaga en las plantaciones de *Eucalyptus* (Myrtaceae) en Brasil. Este gorgojo poda las ramas principales y laterales de los eucaliptos para poner sus huevos. Un muestreo en 2011 evaluó la incidencia de parásitos este gorgojo. Incidencia de géneros de parásitos fueron evaluados. Se observó que 58,29% de los cortes (o ramas cortadas) contenía enemigos naturales del podador-del-eucalipto. Entre ellos, tres especies del género *Euderus* (Hymenoptera: Eulophidae) comprendido 98,46% de los parásitos; otros eran una especie del género *Eurydinoteloides* Girault (Hymenoptera: Pteromalidae). Este es el primer reporte sobre la presencia de enemigos natural de *C. bicolor*.

Palabras Clave: entomología forestal; podador de eucalipto; control biológico; *Euderus*; *Eurydinoteloides*

References Cited

- Cedeño PE, Flowers RW. 2012. *Heilipodus unifasciatus* (Champion) (Coleoptera: Curculionidae: Molytinae: Hylobiini) attacking plantations of *Ochroma pyramidale* (Cavanilles ex Lamarck) Urban (Malvaceae) in Ecuador. *Coleopterists Bulletin* 66: 344–346.
- Dosdall LM, Gibson GAP, Olfert O, Manson PG. 2009. Responses of Chalcidoidea (Hymenoptera) parasitoids to invasion of the cabbage seedpod weevil (Coleoptera: Curculionidae) in western Canada. *Biological Invasions* 11: 109–125.
- Garlet J, Costa EC, Magistrali IC, Boscardina J, Borges JN. 2011. First report of *Heilipodus dorsosulcatus* (Boheman, 1843) (Coleoptera: Curculionidae) in a plantation of *Eucalyptus* L'Heritier in Brazil. *Coleopterists Bulletin* 65: 243–245.
- Hambleton EJ, Sauer HFG. 1938. Observações sobre as pragas da cultura de algodoeira no nordeste e norte do Brasil. *Arquivos do Instituto Biológico* 9: 319–330.
- Jansson RK, Lecrone SH. 1991. *Euderus purpureas* (Hymenoptera: Eulophidae): a parasitoid of sweetpotato weevil in southern Florida. *Florida Entomologist* 74: 596–598.
- Loch AD. 2008. Parasitism of the eucalyptus weevil, *Gonipterus scutellatus* Gyllenhal, by the egg parasitoid, *Anaphes nitens* Girault, in *Eucalyptus globulus* plantations in southwestern Australia. *Biological Control* 47: 1–7.
- Mafia RG, Silva JB, Ramos JF. 2013. Caracterização dos danos causados por *Heilipodus naevulus* em plantios de eucalipto no Espírito Santo, Brasil. *Ciência Rural* 43: 258–261.
- Mason PJ, Miall JH, Bouchard P, Gillespie DR, Broadbent AB, Gibson GAP. 2011. The parasitoid communities associated with an invasive canola pest, *Ceutorhynchus obstrictus* (Coleoptera: Curculionidae), in Ontario and Quebec, Canada. *The Canadian Entomologist* 143: 524–537.
- Monte O. 1944. Curculionídeos do Algodoeiro. *O Biológico* 10: 278–293.
- Noyes JS. 2015. Universal Chalcidoidea database [online], <http://www.nhm.ac.uk/research-curation/research/projects/chalcidoids/> (last accessed 21 Feb 2015).
- Sauer HFG. 1946. Constatação de himenópteros e dípteros entomófagos no estado de São Paulo. *Boletim Fitossanitário* 3: 7–23.
- Souza RM, Anjos N, Mafia RG, Silva JB. 2011. Primeiro registro de *Chalcodermus bicolor* (Coleoptera: Curculionidae) em plantios de eucalipto. *Ciência Rural* 41: 630–633.
- Sweeney J, Anderson RS, Webster RP. 2012. First records of *Orchestes fagi* (L.) (Coleoptera: Curculionidae: Curculioninae) in North America, with a checklist of the North American Rhamphini. *The Coleopterists Bulletin* 66: 297–304.
- Yoshimoto CM. 1971. Revision of the genus *Euderus* of America north of Mexico (Hymenoptera: Eulophidae). *The Canadian Entomologist* 103: 541–578.