

# Development of the First Chloroplast Microsatellite Loci in Ginkgo biloba (Ginkgoaceae)

Authors: Xie, Chun-Xiang, Zhao, Ming-Shui, Fu, Cheng-Xin, and Zhao, Yun-Peng

Source: Applications in Plant Sciences, 1(8)

Published By: Botanical Society of America

URL: https://doi.org/10.3732/apps.1300019

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



PRIMER NOTE

## Development of the first chloroplast microsatellite loci in *Ginkgo biloba* (Ginkgoaceae)<sup>1</sup>

CHUN-XIANG XIE<sup>2,3</sup>, MING-SHUI ZHAO<sup>4</sup>, CHENG-XIN FU<sup>3</sup>, AND YUN-PENG ZHAO<sup>2,3,5</sup>

<sup>2</sup>Key Laboratory of Conservation Biology for Endangered Wildlife of the Ministry of Education, College of Life Sciences, Zhejiang University, Hangzhou 310058, People's Republic of China; <sup>3</sup>Laboratory of Systematic and Evolutionary Botany and Biodiversity, Institute of Ecology, and Conservation Center for Gene Resources of Endangered Wildlife, Zhejiang University, Hangzhou 310058, People's Republic of China; and <sup>4</sup>Administration Bureau of Tianmu Mountain National Nature Reserve, Lin'an 311311, People's Republic of China

- Premise of the study: To investigate population genetics, phylogeography, and cultivar origin of Ginkgo biloba, chloroplast microsatellite primers were developed.
- *Methods and Results:* Twenty-one chloroplast microsatellite markers were identified referring to the two published chloroplast genomes of *G. biloba.* Polymorphisms were assessed on four natural populations from the two refugia in China. Eight loci were detected to be polymorphic in these populations. The number of alleles per locus ranged from three to seven, and the unbiased haploid diversity per locus varied from 0.441 to 0.807.
- *Conclusions:* For the first time, we developed 21 chloroplast microsatellite markers for *G. biloba*, including 13 monomorphic and eight polymorphic ones within the assessed natural populations. These markers should provide a powerful tool for the study of genetic variation of both natural and cultivated populations of *G. biloba*, as well as cultivars.

Key words: chloroplast microsatellite; cpSSR; genetic diversity; Ginkgo biloba; Ginkgoaceae; gymnosperm; molecular marker.

Ginkgo biloba L., a "living fossil" and one of the most mysterious plant species, is the only extant representative of the isolated gymnosperm order Ginkgoales (Zhou, 2009). It became extinct in North America and Europe, and only survived the recurrent glaciation as a relic in China (Kwant, 2013). Recent molecular and ecological evidence strongly supports the existence of two refugia in China (Gong et al., 2008a, 2008b; Tang et al., 2012). Molecular data also confirmed an out-of-China distribution history mediated by anthropogenic introduction (Zhao et al., 2010). This fascinating plant has been cultivated worldwide for its medicinal and nutritional uses, its power as a source of artistic and religious inspiration, and its importance as one of the world's most popular street trees. Thus, it is imperative to increase our knowledge of its evolutionary history to support our efforts to conserve the refugial (natural) populations. Chloroplast DNA (cpDNA) sequences are prevalent markers used in population genetic studies. However, the relatively low resolution of cpDNA sequences of Ginkgo L. did not fully satisfy the inference of its evolutionary processes (Gong et al., 2008a, 2008b). Here, we report a set of polymorphic and monomorphic chloroplast microsatellite or simple sequence repeat

<sup>1</sup>Manuscript received 15 March 2013; revision accepted 15 April 2013.

The authors thank Q. Y. Dai, Y. Sun, and L. Zheng for technical assistance with the experiment and suggestions on manuscript preparation. This research was supported by the National Natural Science Foundation of China (no. 31000102) and the Fundamental Research Funds for the Central Universities (no. 2012FZA6001).

5Author for correspondence: ypzhao@zju.edu.cn

doi:10.3732/apps.1300019

(cpSSR) markers for *G. biloba*. They are expected to be particularly valuable, together with the published nuclear microsatellite markers (Yan et al., 2006, 2009), not only in inferring the evolutionary history but also for breeding of *G. biloba*.

### METHODS AND RESULTS

Two complete chloroplast genomes of *G. biloba* (NC\_016986 and AB684440) were downloaded from GenBank and manually aligned using Geneious version 4.8.5 (Drummond et al., 2008). Polymorphic mononucleotide repeats longer than nine nucleotides were spotted directly from the alignment. Di-, tri-, tetra-, and pentanucleotide motifs with a minimum of six repeats were identified from one of the chloroplast genomes using SSRHunter version 1.3 (Li and Wan, 2005). A total of 27 chloroplast simple sequence repeat loci (cpSSRs) were identified, including 22 mononucleotide repeats and five dinucleotide repeats. Primers for these 27 cpSSRs were designed using Primer Premier version 5.0 (Clarke and Gorley, 2001) following the criteria: (1) GC content 40–60%; (2) melting temperature ( $T_m$ ) 50–60°C; (3) primer size 18–22 bp in length; and (4) amplicon size 100–400 bp in length.

Primer pairs were initially screened for amplification success using four individuals of *G. biloba* from a Tianmu Mountain population. Genomic DNA was extracted from silica gel–dried leaf materials using the cetyltrimethylammonium bromide (CTAB) method (Doyle and Doyle, 1987). PCR was performed in a 15-µL reaction mixture containing 60–90 ng genomic DNA, 0.75 U *Taq* polymerase (TaKaRa Biotechnology Co., Dalian, China), 0.8 µL 10× PCR buffer (MgCl<sub>2</sub> free), 2 mM MgCl<sub>2</sub>, 0.12 mM dNTPs, and 0.33 µM of each primer, 6.67 mM bovine serum albumin (BSA, TaKaRa Biotechnology Co.). The amplification programs were as follows: initial denaturation at 94°C for 5 min; 30 cycles of 30 s at 94°C, 30 s at the optimized annealing temperature (Table 1), and extension for 40 s at 72°C; and a final extension step at 72°C for 10 min. Amplification products, along with a DL2000 DNA ladder (TaKaRa Biotechnology Co.), were electrophoresed on 2.0% agarose gels stained with ethidium bromide to assess successful amplification. Twenty-one of the 27 primer pairs

Applications in Plant Sciences 2013 1(8): 1300019; http://www.bioone.org/loi/apps © 2013 Xie et al. Published by the Botanical Society of America. This work is licensed under a Creative Commons Attribution License (CC-BY-NC-SA).

TABLE 1.	Characteristics of 21	chloroplast micros	atellite primers	developed in	Ginkgo biloba.
----------	-----------------------	--------------------	------------------	--------------	----------------

Locus <sup>a</sup>	Gene <sup>b</sup>	Repeat motif		Primer sequences $(5'-3')$	$T_{\rm a}(^{\circ}{\rm C})$	Size range (bp)	GenBank accession no.
GCP5 (HEX)	petN-psbM*	(T) <sub>11</sub>	F:	ATATGACGAACCCCTCCCGC	55	171–175	KC344393
			R:	TTCTCCTGTGACAAATCCGA			
GCP6 (TAMRA)	trnC-GCA-petN*	$(T)_{10}$	F:	CCACTTCTTCCCCATACTAC	55	226-229	KC344394
			R:	GAATCTCTCAGGTTGACTCG			
GCP22 (FAM)	psbK-psbL*	$(G)_8(A)_{12}$	F:	TCCCCATTTATCCGTTTGTT	55	245-250	KC344388
			R:	CGGGATCAGTATTGGAAGTG			
GCP23 (HEX)	trnG-GCC*	$(A)_{12}$	F:	GAAATAGGGAACGAAGTAAC	55	311-320	KC344389
			R:	GTGTAAAGGGGTATGAAGAA			
GCP24 (TAMRA)	atpF-atpH*	(G) <sub>13</sub>	F:	GAACGGACTAAGTATGGAAA	55	260-262	KC344390
			R:	CAGAGGGTAAAATACGAGGT			
GCP20 (FAM)	$chlP^*$	(T) <sub>24</sub>	F:	TTAGACAGAAAAAACGACTC	55	328-336	KC344387
			R:	GAGAACGATTGGGATGGAAA			
GCP18 (HEX)	psbA-trnK-UUU*	$(T)_{18}(A)_9$	F:	CTGTCATTTCTTTCCTTTCG	55	338-347	KC344386
			R:	ATTTCCCCGTTATCTCCCCC			
GCP27 (TAMRA)	psaL-ycf4*	$(A)_8(A)_9$	F:	TGACCCGATAAGGTGTTGTT	55	179-181	KC344391
			R:	CGGCTTCCGAAGTAACTTTT			
GCP4 (FAM)	petN-psbM	$(T)_{10}$	F:	ACGGAGTATTACATAGGTTC	55	225	KC344392
			R:	AATACTGCTCTTTATGCTCC			
GCP9 (TAMRA)	trnE-UUC-trnT-GGU	$(T)_{10}$	F:	ACCCGTAGCGATATTTGTTC	53	184	KC861980
			R:	TTTGTAGTATTCAGCACCCA			
GCP1 (FAM)	cemA-petA	$(TC)_8$	F:	TATTCCTTTCCCTCTTTCTC	51	108	KC861981
			R:	TTACCTGTCCACGATTCTTC			
GCP2 (HEX)	psbH-petB	$(A)_{10}$	F:	ATTCCCGTGAGTTGGGTCTA	55	193	KC861982
			R:	GATTTTAGCAAGTCGCACAC			
GCP3 (TAMRA)	trnN-GUU-chlL	$(TA)_6$	F:	GGAAGGAGAGAGGAGGAATG	53	233	KC861983
			R:	GAAAGGGTTGAATGGTAG			
GCP7 (FAM)	rps18-rpl20	$(A)_9$	F:	CAACAACGCCTAATGACTAT	50	290	KC861984
			R:	TAATGAAATAGGTGGAGCAG			
GCP8 (HEX)	trnE-UUC-trnT-GGU	$(TA)_7$	F:	TTAGTCCTGAAACGATGAAC	50	199	KC861985
			R:	GATTTGAACCGATGACTTAC			
GCP15 (HEX)	rsp7-trnL-CAA	$(T)_9$	F:	CCCACTCGGACTAAGACATA	52	93	KC861986
	_		R:	TAGTTGTTTCTGACCTCGCT			
GCP17 (FAM)	trnH-GUG-psbA	(A) <sub>23</sub>	F:	AGCAACACTGGCAGATAAAG	53	211	KC861987
	_		R:	GTAACGCTCACAACTTCCCT			
GCP19 (TAMRA)	trnG-GCC-trnR-UCU	$(T)_{28}$	F:	CGCTACCCGCTCAGATTCCT	57	233	KC861988
			R:	TGGACGCTTTTTCTCCTTCT			
GCP21 (TAMRA)	petB-petD	$(T)_{25}$	F:	AGGTATTCCAGGTCCTTTAT	50	234	KC861989
	* *		R:	CTCCCATAATCCATCTTCTC			
GCP25 (FAM)	trnT-GGU-psbD	$(TA)_7$	F:	CGTTATGATCTCTTCTCCTC	51	254	KC861990
	*		R:	GTGCTTTTCCCCTATTCGTA			
GCP26 (HEX)	psal-ycf4	$(AT)_9(AT)_7$	F:	GGACACGAACAAAATAGAGA	50	203	KC861991
	* • •		R:	GTCAGTGTGATGGAAATACC			

*Note:*  $T_a$  = annealing temperature.

<sup>a</sup>The fluorescent dye label used for each forward primer is given in parentheses.

<sup>b</sup>Polymorphic loci are marked with an asterisk (\*); all other loci in Table 1 are monomorphic.

produced amplicons matching the expected sizes. Polymorphisms of these 21 cpSSRs were assessed using 39 accessions from four populations, i.e., Jinfo Mountain, Chongqing Municipality (JF); Wuchuan, Guizhou Province (WC); Enshi, Hubei Province (ES); and Tianmu Mountain, Zhejiang Province (TM) (Appendix 1). As our previous studies (Gong et al., 2008a, 2008b) suggested, these populations are located in the two refugia. We repeated the experiments to verify the reproducibility of these amplicons. The forward primer of each pair was labeled with a fluorescent dye (6-FAM, HEX, or TAMRA). Amplification followed the conditions described above. Equal volumes of 10× diluted PCR products with three different dye-labeled primers were mixed with GeneScan 500 ROX Size Standard (Applied Biosystems, Carlsbad, California, USA). Fragment analyses were performed on a MegaBACE 1000 DNA Analysis System (GE Healthcare Biosciences, Pittsburgh, Pennsylvania, USA), and alleles were manually scored using GeneMarker version 1.97 (SoftGenetics, State College, Pennsylvania, USA). The resulting genotype data were analyzed using GenAlEx version 6.4.1 (Peakall and Smouse, 2006) to estimate number of alleles per locus (A) and unbiased haploid diversity (h).

Eight out of the 21 stably amplified cpSSR markers were polymorphic in the four analyzed populations of *G. biloba* (Table 1). At the species level, the number of alleles per locus ranges from three to seven, and the unbiased haploid diversity per locus varies from 0.441 to 0.807 (Table 2). At the population level, the values of these two parameters range from one to five and from 0.250 to

0.810, respectively (Table 2). Population TM exhibits the highest level of genetic diversity (A = 3.9, h = 0.627), exceeding populations WC (2.1 and 0.384) and JF (2.0 and 0.295), while population ES shows the lowest values (1.8 and 0.277).

#### CONCLUSIONS

The first 21 chloroplast microsatellite markers were developed for *G. biloba*, including 13 monomorphic and eight polymorphic ones. The alleles of the eight polymorphic chloroplast loci were verified to be reliable. This set of novel polymorphic and monomorphic cpSSR markers provides a valuable tool for population genetic and phylogeographic investigations of *G. biloba*. They can also be useful for plant breeding companies, seed/sapling testing agencies, and the wider scientific community in identifying maternal parents and distinguishing cultivars of ginkgo. Furthermore, given the conservative nature of cpDNA loci, these markers might also be useful for other gymnospermous

TABLE 2. Characteristics of eight polymorphic chloroplast microsatellite loci in four populations of *Ginkgo biloba*.<sup>a</sup>

	JF	(n = 8)	WC	(n = 8)	ES	(n = 8)	TM	(n = 15)	All (	(n = 39)
Locus	Α	h	Α	h	Α	h	A	h	Α	h
GCP5	2	0.429	2	0.250	2	0.250	3	0.590	4	0.441
GCP6	2	0.250	2	0.429	1	0.000	4	0.733	4	0.656
GCP22	3	0.679	3	0.607	2	0.536	5	0.790	6	0.807
GCP23	2	0.250	2	0.250	1	0.000	5	0.705	7	0.640
GCP24	2	0.250	2	0.429	1	0.000	2	0.248	3	0.521
GCP20	1	0.000	2	0.429	2	0.250	4	0.695	6	0.771
GCP18	2	0.250	2	0.250	2	0.429	5	0.810	6	0.722
GCP27	2	0.250	2	0.429	3	0.750	3	0.448	3	0.582
Mean	2.0	0.295	2.1	0.384	1.8	0.277	3.9	0.627	4.9	0.642

*Note:* A = number of alleles per locus; h = unbiased haploid diversity; n = sample size.

<sup>a</sup>Locality information for the populations: JF = Jinfo Mountain, Chongqing Municipality; WC = Wuchuan, Guizhou Province; ES = Enshi, Hubei Province; TM = Tianmu Mountain, Zhejiang Province.

species, because three polymorphic cpDNA loci (i.e., *trnK* intron, *trnS-trnG* spacer [Gong et al., 2008a], and *atpH-atpI* spacer [Zhao et al., in preparation]) were successfully amplified in ginkgo using the universal markers.

### LITERATURE CITED

- CLARKE, K. R., AND R. N. GORLEY. 2001. Primer v5: User manual/tutorial. Primer-E Ltd., Plymouth, United Kingdom.
- DOYLE, J. J., AND J. L. DOYLE. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11–15.

- DRUMMOND, A. J., B. ASHTON, M. CHEUNG, J. HELED, M. KEARSE, R. MORE, S. STONES-HAVAS, AND A. WILSON. 2008. Geneious version 4.5.5 created by Biomatters. Website http://www.geneious.com [accessed 8 December 2012].
- GONG, W., C. CHEN, C. DOBEŠ, C. X. FU, AND M. A. KOCH. 2008a. Phylogeography of a living fossil: Pleistocene glaciations forced *Ginkgo biloba* L. (Ginkgoaceae) into two refuge areas in China with limited subsequent postglacial expansion. *Molecular Phylogenetics* and Evolution 48: 1094–1105.
- GONG, W., Z. ZENG, Y. Y. CHEN, C. CHEN, Y. X. QIU, AND C. X. FU. 2008b. Glacial refugia of *Ginkgo biloba* and human impact on its genetic diversity: Evidence from chloroplast DNA. *Journal of Integrative Plant Biology* 50: 368–374.
- KWANT, C. 2013. The Ginkgo pages [online]. Website http://kwanten. home.xs4all.nl/ [accessed 28 June 2013].
- LI, Q., AND J.-M. WAN. 2005. SSRHunter: Development of a local searching software for SSR sites. *Hereditas* 27: 808–810. Website http:// www.bio-soft.net/dna/SSRHunter.htm [accessed 3 July 2013].
- PEAKALL, R., AND P. E. SMOUSE. 2006. GenAlEx 6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6: 288–295.
- TANG, C. Q., Y. C. YANG, M. OHSAWA, S. R. YI, A. MOMOHARA, W. H. SU, H. C. WANG, ET AL. 2012. Evidence for the persistence of wild *Ginkgo biloba* (Ginkgoaceae) populations in the Dalou Mountains, southwestern China. *American Journal of Botany* 99: 1408–1414.
- YAN, X. F., C. L. LIAN, AND T. HOGETSU. 2006. Development of microsatellite markers in ginkgo (*Ginkgo biloba* L.). *Molecular Ecology Notes* 6: 301–302.
- YAN, X. L., Y. Y. CHEN, AND C. X. FU. 2009. Eleven novel microsatellite markers developed from the living fossil *Ginkgo biloba* (Ginkgoaceae). *Conversation Genetics* 10: 1277–1279.
- ZHAO, Y. P., J. PAULE, C. X. FU, AND M. A. KOCH. 2010. Out of China: Distribution history of *Ginkgo biloba* L. *Taxon* 59: 495–504.
- ZHOU, Z. Y. 2009. An overview of fossil Ginkgoales. *Palaeoworld* 18: 1–22.

APPENDIX 1. Population localities of the samples of *Ginkgo biloba* used in this study. All vouchers are deposited in the Herbarium of Zhejiang University (HZU), Zhejiang, China.

Population code	Population locality	Voucher	Geographic coordinates	Altitude (m)
ES	Yesanguan, Enshi, Hubei	W. Gong, 2004ES173	30°36'37"N, 110°18'29"E	675–1122
JF	Mt. Jinfo, Chongqing	C. X. Fu, 2003JF135	107°10′03″N, 29°01′03″E	800-1300
WC	Wuchuan, Guizhou	Y. Q. Ge, 2001WC011	108°06'49"N, 28°42'56"E	550-1200
TM	Mt. Tianmu, Zhejiang	C. Chuan, 2006TM301	119°26′15″N, 30°19′51″E	300-1100