

PARASITIDS AND PARASITES OF SPODOPTERA FRUGIPERDA (LEPIDOPTERA: NOCTUIDAE) IN THE AMERICAS AND CARIBBEAN BASIN: AN INVENTORY

Authors: Molina-Ochoa, Jaime, Carpenter, James E., Heinrichs, E. A., and Foster, John E.

Source: Florida Entomologist, 86(3) : 254-289

Published By: Florida Entomological Society

URL: [https://doi.org/10.1653/0015-4040\(2003\)086\[0254:PAPOSF\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2003)086[0254:PAPOSF]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PARASITOIDS AND PARASITES OF *SPODOPTERA FRUGIPERDA* (LEPIDOPTERA: NOCTUIDAE) IN THE AMERICAS AND CARIBBEAN BASIN: AN INVENTORY

JAIME MOLINA-OCHOA¹, JAMES E. CARPENTER², E. A. HEINRICHS³ AND JOHN E. FOSTER³

¹Facultad de Ciencias Biológicas y Agropecuarias, Universidad de Colima
Apartado postal no. 36, Tecomán, Colima 28100, México

²United States Department of Agriculture, Agricultural Research Service
Crop Protection & Management Research Laboratory, P.O. Box 748, Tifton, GA 31793-0748, USA

³University of Nebraska Lincoln, Department of Entomology
312F Plant Industry Building, Lincoln, NE 68583-0816, USA

ABSTRACT

An inventory of parasitoids and parasites of fall armyworm (FAW), *Spodoptera frugiperda* (J. E. Smith), was conducted using references describing parasitized FAW eggs, larvae, pupae and adults collected from different crops or habitats throughout the Americas and the Caribbean Basin. The crops and countries where these parasites were reported occurring in the Americas is also inventoried. Maize was the crop where the FAW was more frequently collected followed by rice. Overall, *Chelonus insularis* (Cresson) had the broadest natural distribution in the Americas. For the North American region *C. insularis*, *Chelonus* sp., and *Euplectrus platyhypenae* (Howard) were the most relevant parasitoids. In Central America, *C. insularis* was the most prevalent parasitoid, and in the South American region the most prevalent parasites were *Archytas incertus* (Macq.), *A. marmoratus* (Tns.), *C. insularis*, and *Meteorus laphygmae* (Viereck). *Diapetimorpha introita* (Cresson) is the most important pupal parasitoid of FAW occurring mainly in North America. An acugutturid, *Noctuidonema guyanense* (Remillet & Silvain), is the most important ectoparasitic nematode attacking adults of FAW and other noctuid moths in South and Southeastern US, and Mexico in North America, Caribbean Basin, Central America, and Northern South America.

Key Words: parasitoids, fall armyworm, *Chelonus insularis*, *Diapetimorpha introita*, *Noctuidonema guyanense*, maize, natural distribution, biological control

RESUMEN

Un inventario de los parasitoides y parásitos del gusano cogollero, *Spodoptera frugiperda* (J. E. Smith) se llevó a cabo usando referencias relacionadas con parásitos de huevos, larvas, pupas y adultos del insecto plaga colectados de diferentes cultivos en su ámbito de distribución. Además, un inventario se realizó de los cultivos y países donde estos parásitos atacaron al gusano cogollero en América. La plaga fue colectada principalmente en el maíz, seguido por el arroz. *Chelonus insularis* (Cresson) fue el parasitoide distribuido más ampliamente en todo el ámbito de distribución del. *C. insularis*, *Chelonus* sp. y *Euplectrus platyhypenae* (Howard) fueron los parasitoides más prevalecientes en Norteamérica. En Centroamérica, *C. insularis* fue el parasitoide más prevalente, y en la región Sudamericana lo fueron *Archytas incertus* (Macq.), *A. marmoratus* (Tns.), *C. insularis* y *Meteorus laphygmae* (Viereck). *Diapetimorpha introita* (Cresson) fue el parasitoide de pupas más importante y éste habitó en Norteamérica, principalmente. Un acuguturido, *Noctuidonema guyanense* (Remillet & Silvain) fue el nematodo ectoparásito más importante atacando adultos de gusano cogollero y otros noctuidos en el Sur y Sureste de los Estados Unidos de América y México en Norteamérica, Cuenca del Caribe, Centroamérica y Norte de Sudamérica.

Descriptores: parasitoides, gusano cogollero, *Chelonus insularis*, *Diapetimorpha introita*, *Noctuidonema guyanense*, maíz, ámbito de distribución, control biológico

Biodiversity in agro-ecosystems can be as varied as the crops, weeds, arthropods, and microorganisms themselves, and may differ according to geographical location, climate, soil, and human factors. Experimental evidence suggests that

biodiversity can be used for improved pest management (Altieri 1991). A major problem in all areas of agriculture is the lack of basic research on taxonomy of insect pests and their natural enemies. This problem is greatest in tropical coun-

tries where the needs are also greater (Claridge 1991). Waage (1991) stated that agricultural systems are generally simpler than the natural habitats from which they are developed. These systems have fewer plant species, fewer primary consumers, and generally fewer natural enemy species. In tropical systems, despite their great biodiversity, natural enemies could be particularly susceptible to local extinction as a result of habitat destruction and unfavorable cropping practices such as indiscriminate use of pesticides (Claridge 1991; Waage 1991).

More than 200 years ago, the fall armyworm (FAW), *Spodoptera frugiperda* (J. E. Smith), was recognized as a destructive pest of many agricultural crops (Luginbill 1928). In the continental United States the costs for chemical control and losses due to this pest exceeded \$300,000,000 during 1977 (Gross & Pair 1986). The biological control of FAW in areas of overwintering and throughout its annual geographical distribution is a highly desirable alternative to conventional control methods (Gross & Pair 1986). Luginbill (1928) and Vickery (1929) recognized the value of parasitoids in reducing larval populations of FAW. Parasitoids and parasites can be highly effective at little or no cost, serve as biotic insecticides in place of chemicals, provide long-term control without the target pest developing significant resistance to them, and impose minimal or no harm to humans or the environment (Wilson & Huffaker 1976; Stary & Pike 1999).

Sivasubramaniam et al. (1997), and Dent (2000) advocated that the first step in any investigation of the role of natural enemies in pest control should involve a field survey to determine which species are present and how their numbers vary in relation to those of the pest insects. Surveys on the FAW parasitoids and other natural enemies in different parts of its range have been conducted because of increasing economic and environmental concerns (Carrillo 1980; Ashley 1986; Castro et al. 1989; Gross & Pair 1991; Cave 1993; Lezama-Gutiérrez et al. 2001; Molina-Ochoa et al. 2001). However, information about distributions and host plants of the FAW and the accompanying parasites and parasitoids are scattered throughout the published literature, and most reviews of FAW parasitoids have emphasized those attacking the egg and larval stages (Ashley 1986). The aim of this paper is to summarize the information and provide an inventory of the known FAW parasites and parasitoids occurring in the Americas and the Caribbean, indicating the host stage attacked, the crops from which parasitized fall armyworm were collected, and the country of collection. In addition, because parasites and parasitoids of the pupal and adult stages of FAW have received little attention in most previous reviews on this subject, we provide an expanded discussion of these natural enemies.

MATERIALS AND METHODS

Sources of Information

Research was conducted to obtain information, papers, and bibliographic references reporting the collection of parasitized FAW from the field. In the US, we used the Agricola and CAB Abstracts database at the University of Nebraska-Lincoln in Lincoln, Nebraska, and the University of Georgia, Coastal Plain Experiment Station in Tifton, Georgia. We collected references cited in reviews and catalogs of FAW parasites (Guimarães 1971; Marsh 1978; Ashley 1979; Ashley 1986; Andrews 1988) to verify information concerning host plant of parasitized FAW, location of the collected FAW, and stage of FAW attacked. We also used various internet search engines to identify published reports of parasitized FAW collected from the field. In México, we collected papers from Latin American journals and proceedings from meetings of International and Mexican Entomological and Biological Control Societies. Information was also obtained from the libraries in the Facultad de Ciencias Biológicas y Agropecuarias (School of Biological, Agricultural & Animal Sciences) of the Universidad de Colima, and in the Centro Nacional de Referencia de Control Biológico (CNRCB)-Comisión Nacional de Sanidad Agropecuaria-SAGARPA (National Center for Reference on Biological Control) in Tecmán, Colima, México, during 2001 and 2002.

Organization of the Information

The classification (order, family, genus and species) of each parasite and parasitoid collected from FAW is presented in Table 1. The crop from which parasitized FAW were collected, the FAW stage attacked, the country from which parasitized FAW were collected, and the bibliographic references for each record were included in Table 1 whenever the information was available. We preferred to list only original references that report the collection of FAW from the field. Therefore, review articles that contain lists of insects collected by other authors (Guimarães 1971; Marsh 1978; Ashley 1979; Ashley 1986; Andrews 1988) usually are omitted for the entries in Table 1. Because of the quantity involved, all references listed in Table 1 were not cited in the text. Data on FAW parasitoids and parasites from Table 1 have been summarized as the number of species in each taxon reported from different geographical regions (Table 2), Number of species and FAW stage attacked reported from different countries (Table 3), and number of species in each taxon reported from FAW collected from different host plants (Table 4). Omission of pertinent literature from this paper is the authors' responsibility and was unintentional.

TABLE 1. FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
Diptera: Bombyliidae				
<i>Poecilanthrax (Anthrax) lucifer</i> (Fabricius)	L	US	BG	Allen 1921
Diptera: Phoridae				
<i>Megaselia</i> sp.	L	Honduras	(N/G)	Cave 1993
		Nicaragua	M	Maes 1989
Diptera: Sarcophagidae				
<i>Helicobia morionella</i> (Aldrich)	L	Honduras	(N/G)	Cave 1993
Syn: <i>Sarcophaga morionella</i> Aldrich		Nicaragua	M	Maes 1989
<i>Rivinia assidua</i> (Walker)	L	US	(N/G)	Luginbill 1928
Syn: <i>Sarcophaga assidua</i> (Walker)				
<i>Sarcophaga georgiana</i> (Weideman)	L	US	M	Dew 1913
<i>Sarcophaga lambens</i> (Weideman)	L	Lesser Antilles	M	Fennah 1947
<i>Sarcophaga</i> sp.	L	US	(N/G)	Enkerlin 1975
		Venezuela	(N/G)	Terán 1974
<i>Sarcodexia sternodontis</i> (Townsend)	L	Honduras	M	Maes 1989; Cave 1993
Diptera: Tachinidae				
<i>Tachinidae</i> sp.	L	Mexico	M,S	Lacayo 1977
		Nicaragua	M	Ryder & Pulgar 1969; Ashley 1986
<i>Acroglossa vetula</i> (Reinhard)	L	Brazil	(N/G)	Guimarães 1977
Syn: <i>Spallanzania vetula</i> (Reinhard)		Honduras	(N/G)	Cave 1993
		Venezuela	M	Notz 1972
			(N/G)	Terán 1974
<i>Admontia degeerioides</i> (Coquillett)	L	US	(N/G)	Luginbill 1928
<i>Archytas analis</i> Fabricius	L	Argentina	M	Virla et al. 1999
		Barbados	M	Alam 1979
		Honduras	M,R,O,T	Cave 1993
		Mexico	M	Ravlin & Stehr 1984
		Nicaragua	M	Maes 1989
		US	(N/G)	Luginbill 1928
		Venezuela	(N/G)	Terán 1974
<i>Archytas apicifer</i> (Walker)	L	US	(N/G)	Ravlin & Stehr 1984
<i>Archytas incasana</i> Townsend	L	Argentina	M	Virla et al. 1999
Syn: <i>Archytas divisus</i> (Walker)		Brazil	(N/G)	Guimarães 1977
		Chile	M	Etcheverry 1957
		Venezuela	(N/G)	Terán 1974

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Archytas incertus</i> (Macquart)	L	Argentina	M	Virla et al. 1999
			SY	Molinari & Avalos 1997
			(N/G)	Parker et al. 1953
		Barbados	M	Alam 1979
			M	Lucchini & Almeida 1980; Patel & Habib 1982, 1984, 1986; Silveira et al. 1987; Valicente 1989; Silva et al. 1997
			(N/G)	Parker et al. 1953; Guimarães 1977; Milward et al. 1991a,b,c,d,e
		Chile	M	Etcheverry 1957
			SC	Curran 1927
			SC	Van Dine 1913
		Mexico	SC,M	Jones 1913
			M,PN	Segeren & Sharma 1979
			M	Hynes 1942
		Puerto Rico	M,PN	Vickery 1929
			(N/G)	Luginbill 1928
			(N/G)	Parker et al. 1953; Silveira & Ruffinelli 1956
		Suriname	M	Valicente & Barreto 1999; Virla et al. 1999
			SY	Molinari & Avalos 1997
			(N/G)	Avalos 1988
<i>Archytas marmoratus</i> (Townsend)	L	Barbados	M	Alam 1979
			M	Valicente 1989
			(N/G)	Guimarães 1977
		Brazil	(N/G)	Valencia & Valdivia 1973
			(N/G)	Bruner et al. 1975
			(N/G)	Bruner et al. 1975
		Chile	Ve	Benzing et al. 2000
			M	Malausa 1981, 1983
			M	Canas & O'Neil 1998
		Cuba	P,S,M	Cave 1993
			M	Fennah 1947
			M,S	Pair et al. 1986
		Ecuador	M	Huis 1981; Maes 1989; Lacayo 1977
			M	Sarmiento & Razuri 1978
			M	Pantoja et al. 1985; Pantoja & Fuxa 1992
		Guadeloupe	R	Van Dither 1960
			(N/G)	Yaseen 1979
			M	Soteres et al. 1984
		Honduras	M	
			A	

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Archytas marmoratus</i> (Townsend)	L		(M M,C M,S C,A,M,S M,BG,Mi (N/G)	Campos 1965; Enkerlin 1975; Hogg et al. 1982; Gross & Pair 1986, 1991; Riggins et al. 1992, 1993 Tingle et al. 1994 Rohlf & Mack 1985; Pair et al. 1986; McCutcheon 1991 Butler 1958a Reed 1980 Ravlin & Stehr 1984
		Venezuela	M (N/G)	Notz 1972; Fernández & Clavijo 1984 Terán 1974
<i>Archytas plangens</i> Curran	L	Argentina Brazil Honduras	M (N/G) M (N/G)	Virla et al. 1999 Guimarães 1977 King & Saunders 1984 Cave 1993
		Nicaragua	M	Estrada 1960
		Trinidad	M	Hynes 1942
<i>Archytas</i> sp.	L	Argentina	M SY (N/G)	Vera et al. 1995 Molinari & Avalos 1997 Guimarães 1977
		Brazil	PN	Enkerlin 1975
		Chile	M	Wheeler et al. 1989
		Honduras	M	Carrillo 1980
		Mexico	M	Estrada 1960
		Nicaragua	PN	Nickle 1976
		US	M	Cave 1993
<i>Chetogena</i> sp.	L	Honduras	(N/G)	Parker et al. 1953
<i>Cuphocerini</i> sp.	L	Argentina	(N/G)	Bruner et al. 1975
<i>Eucelatoria armigera</i> (Coquillett)	L	Cuba	PN	Wall & Berberet 1975; Nickle 1976
		US	(N/G)	Terán 1974
<i>Eucelatoria australis</i> (Townsend)	L	Peru	(N/G)	Enkerlin 1975
<i>Eucelatoria bryani</i> Sabrosky	L	Honduras	S	Cave 1993
		Nicaragua	M	Maes 1989
		US	S,PN	Sabrosky 1981
<i>Eucelatoria guimaraesi</i>	L	Brazil	S,PN	Sabrosky 1981
<i>Eucelatoria rubentis</i> (Coquillett)	L	US	M M,C S,PN	Ashley et al. 1980 Tingle et al. 1994 Sabrosky 1981

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Eucelatoria</i> sp.	L	Barbados Brazil	M M (N/G)	Alam 1979 Silveira et al. 1987; Valicente 1989 Guimaraes 1977
		Chile Nicaragua Venezuela	PN M M (N/G)	Enkerlin 1975 Maes 1989; Gladstone 1991 Fernández & Clavijo 1984 Terán 1974
<i>Euphorocera floridensis</i> Townsend	L	Honduras	(N/G)	Cave 1993
<i>Euphorocera tachinomoides</i> (Townsend) Syn: <i>Chetogena tachinomoides</i>	L	US	PN	Wall & Berberet 1975
<i>Euphorocera</i> sp.	L	Brazil	M (N/G)	Goncalves & Goncalves 1973; Silva et al. 1997 Guimarães 1977
		US	(N/G)	Nickle 1976
<i>Exorista mella</i> (Walker)	L	US	PN	Wall & Berberet 1975
<i>Gonia capitata</i> DeGeer	L	US	(N/G)	Luginbill 1928
<i>Gonia crassicornis</i> (Fabricius)	L	Brazil Honduras Lesser Antilles Puerto Rico	(N/G) S M SC	Goncalves & Goncalves 1973; Guimarães 1977 Cave 1993 Fennah 1947 Van Dine 1913
		US Venezuela	SC,M (N/G) (N/G)	Jones 1913 Luginbill 1928 Terán 1974
<i>Gonia (Reaumuria) pacifica</i> Townsend	L	Brazil Peru	(N/G) (N/G)	Guimarães 1977 Enkerlin 1975
<i>Gonia texensis</i> Reinhard	L	Cuba	(N/G)	Bruner et al. 1975
<i>Gonia</i> sp.	L	Chile Nicaragua	(N/G) (N/G)	Enkerlin 1975 Maes 1989
<i>Hyphantrophaga hyphantriae</i> (Townsend) Syn: <i>Exorista ceratomiae</i> (Coquillett)	L	US	(N/G)	Luginbill 1928
<i>Hyphantrophaga collina</i> (Reinhard) Syn: <i>Zenillia blanda</i>	L	Cuba	(N/G)	Bruner et al. 1975
<i>Incamyia chilensis</i> (Aldrich)	L	Argentina Brazil Chile	(N/G) (N/G) M (N/G)	Blanchard 1963 Guimarães 1977 Etcheverry 1957 Enkerlin 1975
		Uruguay	(N/G)	Parker et al. 1953; Silveira & Ruffinelli 1956
<i>Lespesia affinis</i> (Townsend)	L	Brazil	M	Guimarães 1983; Silva et al. 1997

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Lespesia aletiae</i> (Riley)	L	Honduras US	(N/G) PN M,S M,BG,Mi	Cave 1993 Wall & Berberet 1975 Pair et al. 1986 Reed 1980
<i>Lespesia archippivora</i> (Riley) Syn: <i>Achaetoneura archippivora</i> (Wile) Syn: <i>Frontina archippivora</i> (Scudder)	L	Argentina Brazil	M M	Virla et al. 1999 Patel & Habib 1984, 1986; Guimarães 1983; Valicente 1989
		Chile	(N/G) M	Guimarães 1977 Etcheverry 1957
		Cuba	(N/G)	Parker et al. 1953; Ryder & Piedra 1968; Bruner et al. 1975
		Guadeloupe	M	Malausa 1981, 1983
		Guatemala	M,Te	Painter 1955
		Honduras	M	Canas & O'Neil 1998
			M,S	Cave 1993
		Lesser Antilles	M	Fennah 1947
		Mexico	M	Carrillo 1980
		Nicaragua	M	Estrada 1960; Huis 1981; Maes 1989; Gladstone 1991; Lacayo 1977
		Puerto Rico	SC SC,M	Van Dine 1913 Jones 1913
		US	A C M PN PN, M M,S C,A,M,S M,BG,Mi,S (N/G)	Soteres et al. 1984 Tingle et al. 1994 Gross & Pair 1986; Riggin et al. 1992, 1993 Wall & Berberet 1975; Nickle 1976 Vickery 1929 Pair et al. 1986 Butler 1958a Reed 1980 Luginbill 1928
		Uruguay	M	Silveira & Ruffinelli 1956 Notz 1972; Fernández & Clavijo 1984
		Venezuela	(N/G)	Terán 1974, 1977
<i>Lespesia frenchi</i> (Williston)	L	US	(N/G)	Luginbill 1928
<i>Lespesia grioti</i> (Blanchard)	L	Argentina	M	Virla et al. 1999
			(N/G)	Blanchard 1963
		Brazil	(N/G)	Guimarães 1977

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Lespesia</i> sp.	L	Argentina	M (N/G)	Virla et al. 1999 Parker et al. 1953
		Brazil	M (N/G)	Lucchini & Almeida 1980 Goncalves & Goncalves 1973; Guimarães 1977
		Colombia	R	Vargas & Sanchez 1983
		Cuba	(N/G)	Ryder & Pulgar 1969
		Honduras	M	Wheeler et al. 1989
		Nicaragua	M	Maes 1989
		Puerto Rico	R	Pantoja et al. 1985; Pantoja & Fuxa 1992
		US	M M,S (N/G)	Ashley et al. 1980 McCutcheon 1991 Parker et al. 1953
		Uruguay	M (N/G)	Notz 1972 Terán 1974
		Venezuela	M	Terán 1977
<i>Linnaemya annalis</i> (Townsend)	L	Venezuela	M	Terán 1977
<i>Linnaemya comta</i> (Fallen)	L	Honduras	(N/G)	Cave 1993
<i>Linnaemya</i> sp.	L	Ecuador	M	Benzing et al. 2000
<i>Nemorilla pyste</i> (Walker)	L	US	A	Soteres et al. 1984
Syn: <i>Exorista pyste</i> (Walker)			C (N/G)	Wilson 1923 Luginbill 1928
<i>Parasetigena</i> sp.	L	Brazil	(N/G)	Guimarães 1977
		Uruguay	(N/G)	Parker et al. 1953; Silveira & Ruffinelli 1956
<i>Patelloa similis</i> (Townsend)	L	Brazil	M (N/G)	Patel & Habib 1984, 1986 Guimarães 1977
<i>Patelloa</i> sp.	L	Argentina	M (N/G)	Virla et al. 1999 Parker et al. 1953
		Uruguay	(N/G)	Parker et al. 1953; Silveira & Ruffinelli 1956
<i>Peleteria robusta</i> Wiedeman	L	Chile	(N/G)	Etcheverry 1957
<i>Phorocera claripennis</i> (Macquart)	L	US	A (N/G)	Soteres et al. 1984 Luginbill 1928
Syn: <i>Chetogena claripennis</i> (Macquart)		Venezuela	(N/G)	Terán 1974
<i>Phorocera floridensis</i> (Townsend)	L	Brazil	M	Valicente 1989
Syn: <i>Chetogena floridensis</i> (Townsend)		Honduras	(N/G)	Cave 1993
		Nicaragua	(N/G)	Maes 1989
		US	PN (N/G)	Enkerlin 1975 Luginbill 1928
		Venezuela	(N/G)	Terán 1974

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Pronemorilla mima</i> Townsend	L	Venezuela	M (N/G)	Notz 1972 Terán 1974
<i>Pseudokea</i> sp.	L	Brazil	(N/G)	Costa-Lima 1949
<i>Voria ruralis</i> (Fallen)	L	Chile	M	Etcheverry 1957
		Argentina	M	Virla et al. 1999
			SY	Molinari & Avalos 1997
			(N/G)	Parker et al. 1953
		Brazil	M	Guimarães 1971
			(N/G)	Guimarães 1977
		US	C,A,M,S	Butler 1958a
<i>Winthemia leucanae</i> (Kirkpatrick)	L	Uruguay	M	Silveira & Ruffinelli 1956
Syn: <i>Nemorea leucanae</i> (Kirkpatrick)		US	M,S,BG,SY,C	Dew 1913
<i>Winthemia mima</i> (Reinhard)	L	Argentina	M	Virla et al. 1999
		Brazil	(N/G)	Guimarães 1977
<i>Winthemia quadripustulata</i> Fabricius	L	Chile	M	Etcheverry 1957
		US	(N/G)	Walton 1913; Luginbill 1928
		Venezuela	(N/G)	Terán 1974
<i>Winthemia reliqua</i>	L	Chile	(N/G)	Valencia & Valdivia 1973
<i>Winthemia roblesi</i>	L	Chile	(N/G)	Valencia & Valdivia 1973
<i>Winthemia rufopicta</i> (Bigot)	L	US	A	Soteres et al. 1984
			C	Tingle et al. 1994
			M	Hofmaster & Greenwood 1949; Ashley et al. 1980; Riggins et al. 1992, 1993
			PN	Wall & Berberet 1975
			M,S	Rohlf's & Mack 1985; Pair et al. 1986
			PN,BG	Enkerlin 1975
			C,A,M,S	Danks 1975
			M,BG,Mi,S	Reed 1980
<i>Winthemia sinuata</i> (Reinhard)	L	US	PN	Wall & Berberet 1975
<i>Winthemia trinitatis</i> (Thompson)	L	Argentina	M	Virla et al. 1999
		Brazil	M	Valicente 1989
			(N/G)	Guimarães 1977
<i>Winthemia</i> sp.	L	Argentina	M	Vera et al. 1995; Virla et al. 1999
		Brazil	M	Escalante 1974; Patel & Habib 1984, 1986; Silva et al. 1997
			(N/G)	Guimaraes 1977

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Winthemia</i> sp.	L	Chile	M	Etcheverry 1957; Campos 1965
			M,PN	Enkerlin 1975
		Colombia	R	Vargas & Sánchez 1983
		Honduras	M,S	Cave 1993
		Lesser Antilles	(N/G)	Fennah 1974
		Mexico	M	Guevara et al. 1979
		Peru	M	Sarmiento & Razuri 1978
		Trinidad	(N/G)	Yaseen 1979; Hynes 1942
		US	PN	Wall & Berberet 1975
			M,S	McCutcheon 1991
		Venezuela	(N/G)	Terán 1974
Hymenoptera: Bethyridae				
<i>Perisierola</i> sp.	L	US	M	Bianchi 1944
Hymenoptera: Braconidae				
<i>Agathis stigmatera</i> (Cresson)	E	Argentina	M	De Santis 1967; De Santis & Esquivel 1966; Virla et al. 1999
			(N/G)	Parker et al. 1953
<i>Aleiodes caphimal</i>	L	Nicaragua	(N/G)	Andrews 1988
Syn: <i>Rogas caphimal</i>				
<i>Alieodes laphygmae</i> (Viereck)	L	Brazil	M	Cruz et al. 1997b
Syn: <i>Rogas laphygmae</i> (Viereck)		Chile	A,M,C	Etcheverry 1957
		Honduras	M	Wheeler et al. 1989; Canas & O'Neil 1998
			M,S	Cave 1993
		Mexico	M	Molina-Ochoa et al. 2001
			M,S	Pair et al. 1986
		Nicaragua	M	Estrada 1960; Huis 1981; Maes 1989; Gladstone 1991
		Puerto Rico	R	Pantoja & Fuxa 1992
		US	M	Vickery 1929; Bianchi 1944; Ashley et al. 1982; Mitchell et al. 1984; Gross & Pair 1986; Riggins et al. 1992, 1993
			S	Rohlf & Mack 1985
			M,S	Reed 1980; Pair et al. 1986; Isenhour 1988; McCutcheon 1991
			M,BG,PG	Ashley et al. 1983
			(N/G)	Luginbill 1928; Muesebeck et al. 1951
<i>Aleiodes terminalis</i> (Cresson)	L	Canada	A,M,C	Muesebeck et al. 1951
Syn: <i>Rogas terminalis</i> (Cresson)		US	A,M,C	Muesebeck et al. 1951
			(N/G)	Luginbill 1928; Marsh & Shaw 2001

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Aleiodes vaughani</i> (Muesebeck) Syn: <i>Rogas vaughani</i> Muesebeck	L	Honduras	M	Cave 1993; Passoa 1983; Wheeler et al. 1989; Canas & O'Neil 1998
		Nicaragua	M	Huis 1981; Maes 1989
<i>Aleiodes</i> sp. Syn: <i>Rogas</i> sp.	L	Argentina	M	Virla et al. 1999
		Cuba	M	Ryder & Pulgar 1969; Ryder & Piedra 1968
		Honduras	M	Passoa 1983
		Nicaragua	M	Estrada 1960; Maes 1989; Lacayo 1977
		US	A	Soteres et al. 1984
<i>Bassus</i> sp.	L	Argentina	(N/G)	Parker et al 1953
		Honduras	(N/G)	Cave 1993
<i>Bracon kirkpatricki</i> Wilkinson	L	Mexico	(N/G)	Moya 1980; Pena 1980
<i>Cardiochiles nigriceps</i> (Viereck)	L	US	C	Tingle et a. 1994
<i>Chelonus antillarum</i> (Marshall)	E	Barbados	M	Alam 1979
		Nicaragua	M	Ryder & Pulgar 1969
<i>Chelonus cautus</i> Cresson Syn: <i>Microchelonus cautus</i> (Cresson)	E	Honduras	M	Cave 1993; Canas & O'Neil 1998
		Mexico	M,S	Molina-Ochoa et al. 2001
		Nicaragua	M	Huis 1981
<i>Chelonus formosanus</i> (Sonan)	E	Barbados	(N/G)	Alam 1979
		Trinidad	M	Yaseen 1979
<i>Chelonus insularis</i> (Cresson) Syn: <i>Chelonus texanus</i> (Cresson)	E	Argentina	M	Virla et al. 1999
			(N/G)	Parker et al. 1953
		Barbados	(N/G)	Alam 1979
		Brazil	M	Patel & Habib 1982, 1984, 1986; Valicente 1989; Rezende et al. 1995a,b; Cruz et al. 1997a,b; Silva et al. 1997
			M,C	Rezende et al. 1994
		Chile	A,M,C	Etcheverry 1957
		Colombia	M	Medina et al. 1988
			R	Vargas & Sánchez 1983
		Cuba	BG	Myers 1932
			M	Ryder & Pulgar 1969
			(N/G)	Ryder & Piedra 1968; Bruner et al. 1975;
		Haiti	BG	Myers 1932
		Honduras	M	Wheeler et al. 1989; Canas & O'Neil 1998
			M,S	Castro et al. 1989; Cave 1993
		Lesser Antilles	M	Fennah 1947

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Chelonus insularis</i> (Cresson) Syn: <i>Chelonus texanus</i> (Cresson)	E	Mexico	M	Bahena & García 1991; Nava & Castro 1991; Arce & Garcia 1995; Molina-Ochoa et al. 2001
			M, S	Pair et al. 1986
		Nicaragua	M	Estrada 1960; Huis 1981; Ashley 1986; Maes 1989; Gladstone 1991
			M,S	Lacayo 1977
		Puerto Rico	R	Pantoja & Fuxa 1992
			M	Yaseen 1979
		Trinidad	(N/G)	Yaseen et al. 1981
			A	Soteres et al. 1984
		US	C	Tingle et al. 1994
			M	Pierce & Holloway 1912; Vickery 1929; Bianchi 1944; Muesebeck et al. 1951; Ashley et al. 1980, 1982; Ashley 1983, 1986; Mitchell et al. 1984; Gross & Pair 1986; Riggins et al. 1992
			PN	Wall & Berberet 1975
			M,BG	Ashley et al. 1983
			M,PN	Enkerlin 1975
			M,S	Waddill & Whitcomb 1982; Rohlf & Mack 1985; Pair et al. 1986; McCutcheon 1991
			C,A,M,S	Butler 1958b
			(N/G)	Luginbill 1928
			(N/G)	Parker et al. 1953; Silveira & Ruffinelli 1956
			M	Notz 1972; Fernández & Clavijo 1984
<i>Chelonus</i> sp.	E	Brazil	M	Valicente 1989
			M	Loya-Ramírez 1978; Coronado & Ruíz 1991; Cortez & Trujillo 1994; Sánchez-García et al. 1998; Carrillo 1980
		Mexico	M,S	Molina-Ochoa et al. 2001
			C	Herrera-Arangüena 1998
		Peru	(N/G)	Cave 1993
		Honduras	M	Maes 1989
		Nicaragua	M	Alam 1979
		Barbados	M	Wheeler et al. 1989; Virla et al. 1999
		Argentina	M	Lucchini & Almeida 1980; Patel & Habib 1982, 1984, 1986; Cruz et al. 1997b
		Brazil	M	Etcheverry 1957
<i>Cotesia (Apanteles) congregata</i> (Say)	E,L ³	Chile	A,M,C	Cave 1993
		Honduras	M	Fennah 1947
		Lesser Antilles	M,S	
<i>Cotesia (Apanteles) glomeratus</i> (Linnaeus)	E,L ³			
<i>Cotesia (Apanteles) marginiventris</i> (Cresson)	E,L ³			
Syn: <i>Apanteles grenadensis</i> (Ashmead)				
Syn: <i>Protapanteles harnedi</i> (Viereck)				
Syn: <i>Apanteles laphygmae</i> (Ashmead)				

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Cotesia (Apanteles) marginiventris</i> (Cresson) Syn: <i>Apanteles grenadensis</i> (Ashmead) Syn: <i>Protapanteles harnedi</i> (Viereck) Syn: <i>Apanteles laphygmae</i> (Ashmead)	E,L ³	Mexico	M M,S	Marsh 1978 Pair et al. 1986; Molina-Ochoa et al. 2001
		Nicaragua	M	Estrada 1960; Huis 1981; Maes 1989; Gladstone 1991
		Puerto Rico	R	Pantoja & Fuxa 1992
		Suriname	M,PN (N/G)	Segeren & Sharma 1979 Van Dither 1960
		US	A C M	Soteres et al. 1984 Tingle et al. 1994 Muesebeck 1921; Vickery 1929; Bianchi 1944; Hofmaster & Greenwood 1949; Muesebeck et al. 1951; Marsh 1978; Ashley et al. 1980, 1982; Hogg et al. 1982; Ashley 1983, 1986; Mitchell et al. 1984; Gross & Pair 1986; Riggins et al. 1992, 1993, 1994; Hamm et al. 1994; Ruberson & Whitfield 1996
			M,S PN M,BG M,S,C,Mi	Rohlf & Mack 1985; Pair et al. 1986; McCutcheon 1991 Wall & Berberet 1975; Nickle 1976 Ashley et al. 1983 Reed 1980
		Uruguay	(N/G)	Luginbill 1928
		Venezuela	(N/G) M SY,To	Parker et al. 1953; Silveira & Ruffinelli 1956 Notz 1972; Fernández & Clavijo 1984 Terán 1980
<i>Cotesia (Apanteles) ruficrus</i> (Haliday) Syn: <i>Microplitis manillee</i> (Ashmead)	E,L ³	Trinidad & Tobago	M	Yaseen 1979*
<i>Cotesia (Apanteles)</i> sp.	E,L ³	US	(N/G)	McCutcheon et al. 1983**; Rajapakse et al. 1985***
		Barbados	M	Alam 1979
		Brazil	M	Patel & Habib 1984, 1986; Silva et al. 1997
		Colombia	R	Vargas & Sánchez 1983
			M	Medina et al. 1988
		Guyana	M	Sinha 1982
		Nicaragua	M M,S	Maes 1989 Lacayo 1977
		Peru	C	Herrera-Aranguena 1998
		Trinidad	M	Yaseen 1979
		US	M	Hofmaster & Greenwood 1949
<i>Distatrix</i> sp.	L	Honduras	M	Cave 1993
<i>Glyptapanteles militaris</i> (Walsh)	L	Honduras	M	Cave 1993
		US	S (N/G)	Rohlf & Mack 1985 Muesebeck et al. 1951

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Gnathopleura</i> sp.	L	Honduras	(N/G)	Cave 1993
<i>Homolobus truncator</i> (Say)	L	Honduras	M,S	Cave 1993
Syn: <i>Zele mellea</i> (Cresson)		Nicaragua US	M	Huis 1981; Maes 1989
			A	Soteres et al. 1984
			M	Vickery 1929; Wall & Berberet 1975; Riggins et al. 1992, 1993
			M,S	Pair et al. 1986; McCutcheon 1991
			M,S,C	Reed 1980
<i>Macrocentrus</i> sp.	L	Barbados Brazil	(N/G)	Luginbill 1928
			(N/G)	Alam 1979
			M	Silva et al. 1997
<i>Meteorus arizonensis</i> (Muesebeck)	L	Honduras	(N/G)	Cave 1993
<i>Meteorus autographae</i> (Muesebeck)	L	Nicaragua Mexico US	M	Maes 1989
			M,S	Pair et al. 1986
			A	Soteres et al. 1984
			C	Tingle et al. 1994
			M	Ashley et al. 1980, 1982; Mitchell et al. 1984; Gross & Pair 1986; Riggins et al. 1992, 1993
			PN	Nickle 1976
			S	Rohlf & Mack 1985
			M,S	Pair et al. 1986; McCutcheon 1991
			M,BG,Mi	Reed 1980
			M,BG,PG	Ashley et al. 1983
<i>Meteorus laphygmae</i> (Viereck)	L	Chile	(N/G)	Luginbill 1928; Muesebeck et al. 1951
			M	Etcheverry 1957
			M	Medina et al. 1988
		Colombia	R	Vargas & Sánchez 1983
			C,S	Ortegón et al. 1988
			M,S	Cave 1993
		Honduras Mexico	M	Molina-Ochoa et al. 2001
			S	Ciceros et al. 1995
		Nicaragua	(N/G)	Gladstone 1991
		Suriname	M,PN	Segeren & Sharma 1979
		US	(N/G)	Van Dither 1960
			M	Vickery 1929; Bianchi 1944
			M,BG	Enkerlin 1975
			(N/G)	Luginbill 1928

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Meteorus laphygmae</i> (Viereck)	L	Venezuela	M	Notz 1972; Terán 1980; Fernández & Clavijo 1984; Fernández & Terán 1990a,b
<i>Meteorus vulgaris</i> (Cresson)	L	US	(N/G)	Luginbill 1928; Muesebeck et al. 1951
<i>Meteorus</i> sp.	L	Mexico	M	Peraza-Lizarraga 1982; Coronado & Ruíz 1991
		Peru	M, S	Molina-Ochoa et al. 2001
			C	Herrera-Aranguena 1998
<i>Microplitis</i> sp.	L	Uruguay	(N/G)	Parker et al. 1953; Silveira & Ruffinelli 1956
<i>Palinzele</i> sp.	L	Trinidad	M	Yaseen 1979
<i>Stantonina</i> sp.	L	Honduras	(N/G)	Cave 1993
		Nicaragua	M	Maes 1989
Hymenoptera: Chalcididae				
<i>Brachymeria ovata</i> (Say)	P	Argentina	M	Virla et al. 1999
		US	(N/G)	Parker et al. 1953
			M	Ashley 1979
<i>Brachymeria robusta</i> (Cresson)	P	US	C	Wilson 1923
			(N/G)	Luginbill 1928
<i>Conura (Spilochalcis) femorata</i> (Fabricius)	L	Honduras	(N/G)	Cave 1993
		Nicaragua	(N/G)	Maes 1989
		US	C	Wilson 1923
			(N/G)	Luginbill 1928
<i>Conura (Ceratosmicra) hirtifemora</i> (Ashmead)	L	US	M	Riggin et al. 1992, 1993
Syn: <i>Splilochalcis hirtifemora</i> (Ashmead)				
<i>Conura (Ceratosmicra) immaculata</i> (Cresson)	L	Honduras	(N/G)	Cave 1993
Syn: <i>Conura fulvomaculata</i> (Cameron)		Nicaragua	M	Maes 1989
<i>Conura (Ceratosmicra) meteori</i> (Burks)	L	US	M	Hofmaster & Greenwood 1949
<i>Conura igneoides</i> (Kirby)	L	US	C	Wilson 1923
Syn: <i>Splilochalcis vittata</i> (Ashmead)			(N/G)	Luginbill 1928; Muesebeck et al. 1951
Hymenoptera: Eulophidae				
<i>Euplectrus comstockii</i> Howard	L	Nicaragua	M	Maes 1989
		US	(N/G)	Luginbill 1928
<i>Euplectrus furnicus</i> Walker	L	Argentina	M	De Santis 1979; De Santis 1989; De Santis & Fidalgo 1994; Virla et al. 1999
		Puerto Rico	R	Pantoja & Fuxa 1992
<i>Euplectrus hircinus</i> (Say)	L	Panama	(N/G)	Andrews 1988
<i>Euplectrus insularis</i> Howard	L	Honduras	M	Wheeler et al. 1989
		Nicaragua	(N/G)	Huis 1981

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Euplectrus marginatus</i> Ashmead	L	Nicaragua	M	Maes 1989
<i>Euplectrus plathypenae</i> Howard	L	Barbados	M	Alam 1979
		Brazil	(N/G)	Costa-Lima 1962
		Chile	M	Etcheverry 1957
		Colombia	M	Vargas & Sánchez 1983
		Cuba	Pm&Pb	Myers 1932
			(N/G)	Ryder & Piedra 1968; Bruner et al. 1975
		Guyana	M	Sinha 1982
		Lesser Antilles	M	Fennah 1947
		Mexico	M	Guevara et al. 1979; Montoya-Burgos 1980; Gutiérrez-Rodriguez 1982; Molina-Ochoa et al. 2001
		Nicaragua	M	Ryder & Pulgar 1969
		Puerto Rico	R	Pantoja & Fuxa 1992
		Trinidad	(N/G)	Yaseen 1979
		US	M	Vickery 1929; Muesebeck et al. 1951; Ashley et al. 1980, 1982; Hogg et al. 1982; Riggins et al. 1992, 1993
			PN	Wall & Berberet 1974, 1975
			(N/G)	Luginbill 1928
		Venezuela	M	Marín-Acosta 1966
			S,C	Guagliami 1962
<i>Euplectrus ronnai</i> (Brethes)	L	Brazil	M	De Santis 1980
<i>Euplectrus</i> sp.	L	Brazil	M	Lucchini & Almeida 1980
		Cuba	BG	Myers 1932
		Honduras	M	Cave 1993
		Mexico	M	Montoya-Burgos 1979, 1980; Cortez & Trujillo 1994
			M,S	Pair et al. 1986
		Nicaragua	M	Huis 1981
			M,S	Lacayo 1977
		US	BG	Reed 1980
			M	Keller 1980
			M,S	Pair et al. 1986
			(N/G)	Enkerlin 1975
<i>Trichodischia caerulea</i>	L	Brazil	(N/G)	Guimaraes 1977
<i>Trichodischia soror</i> (Bigot)	L	Argentina	M	Virla et al. 1999
			(N/G)	Parker et al. 1953; Blanchard 1963
		Brazil	(N/G)	Cortes 1980
<i>Trichospilus pupivora</i> (Ferriere)	P	Barbados	M	Alam 1979
<i>Trichospilus</i> sp.	L	Argentina	(N/G)	Parker et al. 1953

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
Hymenoptera: Ichneumonidae				
<i>Amblyteles</i> sp.	L	Brazil	(N/G)	Costa-Lima 1949
		Chile	M	Etcheverry 1957
<i>Ancyloneura</i> sp.	L	Argentina	(N/G)	Parker et al. 1953
<i>Anomalon ejuncidum</i> (Say)	L	US	M,S	Pair et al. 1986
<i>Campoletis chloridae</i> (Vierech)	L	Barbados	M	Alam 1979
<i>Campoletis curvicauda</i> (Blanchard)	L	Peru	M	Ayqui-Vilca 1993
<i>Campoletis flavicincta</i> (Ashmead)	L	Brazil	M	Patel & Habib 1982, 1984, 1986; Cruz et al. 1997a,b
		Honduras	(N/G)	Cave 1993
		Mexico	M,S	Molina-Ochoa et al. 2001
		Nicaragua	M	Huis 1981; Maes 1989
		US	A	Soteres et al. 1984
			M	Hogg et al. 1982
			PN	Wall & Berberet 1975
		Uruguay	(N/G)	Parker et al. 1953; Yaseen et al. 1981
<i>Campoletis grioti</i> (Blanchard)	L	Argentina	M	Virla et al. 1999
		Brazil	M	Lucchini & Almeida 1980; Silveira et al. 1987; Cruz et al. 1997b
		US	M	Ashley 1983
		Uruguay	M	Morey 1971
<i>Campoletis oxylus</i> (Cresson)	L	US	M	Muesebeck et al. 1951
Syn: <i>Sagaritis oxylus</i> (Cresson)			M,S	Pair et al. 1986
			(N/G)	Luginbill 1928
<i>Campoletis sonorensis</i> (Cresson)	L	Brazil	M	Cruz et al. 1997b
		Chile	M	Machuca et al. 1989
		Honduras	M	Canas & O'Neil 1998
			M,S	Cave 1993
		Mexico	M,S	Pair et al. 1986
		US	A	Soteres et al. 1984
			M	Gross & Pair 1986; Riggins et al. 1992, 1993
			M,S	Pair et al. 1986; Isenhour 1988
<i>Campoletis</i> sp.	L	Argentina	M	Vera et al. 1995
		Brazil	M	Silva et al. 1997; Silveira et al. 1987; Valicente & Barreto 1999
			(N/G)	Cruz et al. 1999
		Nicaragua	M	Maes 1989
		US	M,S	McCutcheon 1991

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Campoplex</i> sp.	L	Brazil	M	Silva et al. 1997
<i>Cryptus albitarsis</i> (Cresson)	P	US	M	Pair & Gross 1989
<i>Diadegma</i> sp.	L	Argentina	M	Porter 1998; Virla et al. 1999
		Brazil	M	Silva et al. 1997
		Cuba	M	Ayala-Sifontes et al. 1978
		Mexico	M	Flores-Dávila et al. 1991
<i>Diapetimorpha introita</i> (Cresson)	P	Honduras	M	Cave 1993
		US	M	Pair & Gross 1984, 1989
			M,S	Pair et al. 1986
<i>Eiphosoma annulatum</i> (Cresson)	L	Venezuela	M	Notz 1972
<i>Eiphosoma vitticole</i> (Cresson)	L	Bolivia	M	Ashley 1979
			(N/G)	Yaseen et al. 1981
		Brazil	M	Patel & Habib 1982, 1984, 1986; Valicente 1989; Silveira et al. 1987; Cruz et al. 1997b
		Colombia	M	Medina et al. 1988
		Honduras	M	Wheeler et al. 1989; Canas & O'Neil 1998
			M,S	Cave 1993
		Mexico	M,S	Pair et al. 1986
		Nicaragua	M	Huis 1981
		US	M	Ashley 1983
		Venezuela	(N/G)	Giraldo-Vanegas & Garcia-R. 1992, 1994a,b, 1995
<i>Eiphosoma</i> sp.	L	Brazil	M	Cruz et al. 1997a; Silva et al. 1997
		Mexico	M	Cortez & Trujillo 1994
		Trinidad	M	Hynes 1942
			(N/G)	Yaseen et al. 1981
		Venezuela	M	Fernández & Clavijo 1984
<i>Enicospilus flavus</i> (Fabricius)	L	US	C	Wilson 1923
Syn: <i>Enicospilus concolor</i> (Cresson)			(N/G)	Luginbill 1928
<i>Enicospilus merdarius</i> (Gravenhorst)	L	Argentina	M	Muesebeck et al. 1951
Syn: <i>Enicospilus purgatus</i> (Say)		Cuba	M	Bruner et al. 1975
		Honduras	(N/G)	Cave 1993
		Nicaragua	M	Maes 1989
		US	C,M	Dew 1913
			(N/G)	Luginbill 1928
<i>Goryphina</i> sp.	L	Brazil	M	Silva et al. 1997
<i>Hyposoter</i> sp.	L	Honduras	M	Cave 1993
		Uruguay	(N/G)	Parker et al. 1953; Silveira & Ruffinelli 1956

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Ichneumon ambulatorius</i> Fabricius	P	US	M	New record (JEC)
<i>Ichneumon promissorius</i> (Cresson)	P	US	M	New record (JEC)
<i>Isdromas lycaenae</i> (Howard)	L	US	M	Riggin et al. 1992, 1993
<i>Mesochorus disceitergus</i> (Say)	L	US	M	Hofmaster & Greenwood 1949; Riggin et al. 1992, 1993
<i>Microcharops anticarsiae</i> (Gupta)	L	Honduras	M S	Wheeler et al. 1989 Cave 1993
<i>Netelia sayi</i> (Cushman)	L	US	C	Tingle et al. 1994
<i>Netelia</i> sp.	L	Peru	M	Escalante 1974
		US	A	Soteres et al. 1984
<i>Parania (Atrometus) tricolor</i> (Morley)	L	Uruguay	(N/G)	Parker et al. 1953; Silveira & Ruffinelli 1956
<i>Pristomerus spinator</i> (Fabricius)	L	Brazil	M	Patel & Habib 1984, 1986
Syn: <i>Neopristomerus appalachianus</i> (Viereck)		Honduras	M	Wheeler et al. 1989; Canas & O'Neil 1998
			M,S	Cave 1993
		Mexico	M	Carrillo 1980
			M,S	Pair et al. 1986; Molina-Ochoa et al. 2001
		Nicaragua	M	Estrada 1960; Huis 1981; Gladstone 1991; Maes 1989; Lacayo 1977
		US	A	Soteres et al. 1984
			C	Tingle et al. 1994
			M	Vickery 1929; Bianchi 1944
			PN	Wall & Berberet 1975
			M,S	Pair et al. 1986
			(N/G)	Luginbill 1928
<i>Ophion ancyloneura</i> (Wichsee)	L	Argentina	M	De Santis 1967; Virila et al. 1999
			(N/G)	Parker et al. 1953
		Uruguay	(N/G)	Silveira & Ruffinelli 1956
<i>Ophion bilineatus</i> (Say)	L	Chile	M	Etcheverry 1957
		US	M	Vickery 1929
			PN	Enkerlin 1975
			(N/G)	Luginbill 1928
<i>Ophion flavidus</i> (Brulle)	L	Argentina	M	Virila et al. 1999
		Brazil	M	Goncalves 1973; Patel & Habib 1982, 1984, 1986; Silveira et al. 1987
		Honduras	M	Wheeler et al. 1989; Canas & O'Neil 1998
			M,S	Cave 1993
			(N/G)	Passoa 1983

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Ophion flavidus</i> (Brulle)	L	Mexico	M	Molina-Ochoa et al. 2001
			M,S	Pair et al. 1986
		Nicargua	M	Maes 1989; Gladstone 1991
		US	M	Hogg et al. 1982; Gross & Pair 1986, 1991; Riggin et al. 1992, 1993
			M,S	Rohlf's & Mack 1985; Pair et al. 1986
			M,BG,Mi,S	Reed 1980
<i>Ophion merdarius</i> (Gravenhorst)	L	Nicaragua	M	Maes 1989
Syn: <i>Enicospilus purgatus</i> (Gravenhorst)		US	(N/G)	Luginbill 1928
<i>Ophion</i> sp.	L	Argentina	M	Vera et al. 1995
		Brazil	M	Silva et al. 1997
		Mexico	M	Ashley 1986
		Nicaragua	M	Huis 1981; Maes 1989
		Peru	M	Escalante 1974
		US	C	Tingle et al. 1994
			M	Mitchell et al. 1984
			PN	Nickle 1976
			M,PG	Ashley et al. 1983
		Uruguay	(N/G)	Parker et al. 1953
<i>Sagaritis dubitatus</i> (Cresson)	L	US	M	Vickery 1929
			(N/G)	Luginbill 1928
<i>Trachysphyrus cleonis</i> (Viereck)	L	Peru	M	Escalante 1974
<i>Temelucha difficilis</i> (Dasch.)	L	US	M	Mitchell et al. 1984; Gross & Pair 1986
				Pair et al. 1986
			M,S	
<i>Temelucha grapholithae</i> (Cushman)	L	Honduras	M	Canas & O'Neil 1998
			M,S	Cave 1993
		Nicaragua	M	Gladstone 1991
<i>Temelucha</i> sp.	L	Honduras	M	Wheeler et al. 1989
		Nicaragua	M	Maes 1989
		US	M	Ashley et al. 1980, 1982, 1983
<i>Vulgichneumon brevicintor</i> (Say)	P	US	M	Pair & Gross 1989
Hymenoptera: Perilampidae				
<i>Perilampus hyalinus</i> (Say)	L	Honduras	M	Cave 1993
(Reported as hiperparasitoid)		Venezuela	M	Notz 1972

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
Hymenoptera: Pteromalidae				
<i>Catalaccus aeneoviridis</i> (Girault)	L	US	M	Hofmaster & Greenwood 1949
<i>Trichomalopsis viridescens</i> (Walsh) Syn: <i>Eupteromalus viridescens</i>	L	US	M	Hofmaster & Greenwood 1949; Riggan et al. 1992, 1993
Hymenoptera: Scelionidae				
<i>Telenomus remus</i> (Nixon)	E	Antigua	M	Irving 1978
		Barbados	M	Alam 1979
		Brazil	M	Correa-Figueiredo et al. 1999
		Colombia	M	Alvarez & Roa 1995
		Dominican Republic	(N/G)	Yaseen et al. 1981
		Guadeloupe	(N/G)	Yaseen et al. 1981
		Guyana	M,S	Sinha 1982
		Honduras	M	Cave & Acosta 1999
		Nicaragua	M	Maes 1989
		Puerto Rico	(N/G)	Wojcik et al. 1976
		Suriname	M	Segeren & Sharma 1979
		Trinidad	M	Yaseen 1979
		US	M,S	Waddill 1977; Waddill & Whitcomb 1982
			(N/G)	Wojcik et al. 1976
		Venezuela	M	Hernández et al. 1989; Hernandez & Díaz 1995, 1996a,b; Gonzalez-Narvaez & Zocco 1996; Ferrer 1998a,b
<i>Telenomus</i> sp.	E	Brazil	M	Cruz et al. 1999
		Colombia	M	Alvarez & Roa 1995
			R	Vargas & Sánchez 1983
		Cuba	M	Armas-García & Ayala-Sifontes 1987; Ayala-Sifontes et al. 1992
		Guadeloupe	M	Malausa 1983
		Mexico	M	Montoya-Burgos 1979; Morales-Pérez 1982; Canseco-Román 1988; García-Lagunas 1988; Barilla-Vera 1989
Hymenoptera: Trichogrammatidae				
<i>Trichogramma demoraesi</i> (Zucchi)	E	Chile	M	Pratissoli et al. 1999
<i>Trichogramma fasciatum</i> Perkins	E	Barbados	M	Alam 1979
		Nicaragua	M	Maes 1989
<i>Trichogramma minutum</i> Riley	E	Nicaragua	M	Maes 1989
		US	M	Muesebeck et al. 1951
			(N/G)	Luginbill 1928

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Trichogramma pretiosum</i> (Riley)	E	Brazil	M	De Sa & Parra 1994; Zucchi et al. 1991
		Nicaragua	M	Huis 1981
<i>Trichogramma</i> sp.	E	Argentina	M	Virla et al. 1999
		Brazil	M	De Sa & Parra 1994
			(N/G)	Cruz et al. 1999
		Colombia	R	Vargas & Sánchez 1983
		Cuba	M	Armas-García & Ayala-Sifontes 1987; Ayala-Sifontes et al. 1988
		Guadeloupe	M	Malausau 1983
			(N/G)	Yaseen et al. 1981
		Mexico	M	Loya-Ramírez 1978; Montoya-Burgos 1979, 1980; Guevara et al. 1979; Rodríguez-Luna 1982; Wong-Arevalo 1982; Johannes-Toonders & Carrillo-Sánchez 1987
				Bahena & García 1991
			(N/G)	Huis 1981; Maes 1989; Mulock et al. 1990
		Nicaragua	M	Waddill & Whitcomb 1982
		US	M,S	
Nematoda: Acugutturidae				
<i>Noctuidonema guyanense</i> Remillet & Silvain	A	Bahamas	(N/G)	Simmons & Rogers 1990b
		Belize	(N/G)	Simmons & Rogers 1990b
		Bermuda	(N/G)	Simmons & Rogers 1990b; Simmons et al. 1991
		Cayman Islands	(N/G)	Simmons & Rogers 1990b
		Colombia	(N/G)	Simmons & Rogers 1990b
		Costa Rica	(N/G)	Simmons & Rogers 1990b
		Dominica	(N/G)	Simmons & Rogers 1990b
		El Salvador	(N/G)	Simmons & Rogers 1990b
		French Guiana	(N/G)	Remillet & Silvain 1988; Marti et al. 1990; Rogers et al. 1990a, 1991, 1993; Silvain & Remillet 1993; Marti et al. 2000
		Grenada	(N/G)	Simmons & Rogers 1990b; Simmons et al. 1991
		Guadeloupe	(N/G)	Marti et al. 2000
		Honduras	(N/G)	Simmons & Rogers 1990b
		Martinique	(N/G)	Simmons & Rogers 1990b; Marti et al. 1990
		Mexico	(N/G)	Simmons & Rogers 1990b
		Panama	(N/G)	Simmons & Rogers 1990b; Simmons et al. 1991
		Puerto Rico	(N/G)	Simmons & Rogers 1990b; Simmons et al. 1991

TABLE 1. (CONTINUED) FALL ARMYWORM, *Spodoptera frugiperda* (J. E. SMITH) (LEPIDOPTERA: NOCTUIDAE), PARASITES AND PARASITOIDS IN THE AMERICAS AND CARIBBEAN BASIN WITH ACCOMPANYING REFERENCES INDICATING FAMILY, HOST STAGE ATTACKED, CROPS FROM WHICH FALL ARMYWORM WERE COLLECTED AND COUNTRIES OF COLLECTION.

Classification of parasite and parasitoid	Host stage attacked ¹	Country of collection	Crop ²	Reference
<i>Noctuidonema guyanense</i> Remillet & Silvain	A	Suriname	(N/G)	Simmons & Rogers 1990b
		Trinidad	(N/G)	Simmons & Rogers 1990b
		US	(N/G)	Marti et al. 1990; Simmons & Rogers 1990b; Rogers et al. 1990b, 1991, 1993, 1996; Simmons et al. 1991; Simmons & Rogers, 1990a, b, 1991, 1994; Marti et al. 2000; Rogers & Marti 1994, 1996
			M	Simmons & Marti 1992
		Virgin Island	M, S (N/G)	Rogers & Marti 1992a, b, 1993a; Marti & Rogers 2000 Simmons & Rogers 1990b

*Imported from Pakistan, ** Imported from Australia, *** Imported from Thailand & Imported from India.

¹Host stage attacked: E (Egg), L (Larva), P (Pupa), A (Adult).

²A = Alfalfa (*Medicago sativa* L.), BG = Bermudagrass (*Cynodon dactylon* L.), C = Cotton (*Gossypium* spp.), M = Maize (*Zea mays* L.), Mi = Millet (*Panicum miliaceum* L.), O = Onion (*Allium cepa* L.), P = Hot pepper (*Capsicum annuum* L.), Pb = (*Panicum barbinode* L.), PG = Paragrass (*Brachiarie mutica* L.), Pm = (*P. maximum* L.), PN = Peanut (*Arachis hypogaea* L.), R = Rice (*Oryza sativa* L.), S = Sorghum (*Sorghum bicolor* L. Moench.), SC = Sugarcane (*Saccharum officinarum* L.), SY = Soybean (*Glycine max* L.), T = Tomato (*Lycopersicum esculentum* L.), Te = Teosinte (*Euchlaena mexicana* Schard), To = Tobacco (*Nicotiana tabacum* L.), Ve = Vegetable Crops, (N/G) = Not given.

³Reported attacking eggs and larva (no confirmation).

TABLE 2. NUMBER OF SPECIES IN EACH TAXON REPORTED FROM DIFFERENT REGIONS.

Order	Family	Region		
		North America (Mexico-US)	Central America & Caribbean Basin	South America
Diptera				
	Bombyliidae	1		
	Phoridae		1	
	Sarcophagidae	3	3	1
	Tachinidae	29	24	38
Hymenoptera				
	Bethylidae	1		
	Braconidae	18	22	11
	Chalcididae	6	2	1
	Eulophidae	3	8	7
	Ichneumonidae	27	18	21
	Perilampidae		1	1
	Pteromalidae	2		
	Scelionidae	2	2	2
	Trichogrammatidae	2	4	3
Nematoda				
	Aphelenchoididae	1	1	1

RESULTS AND DISCUSSION

A great diversity of parasitoids and parasites of FAW has been reported occurring in the Americas and the Caribbean basin. In this inventory, approximately 150 species of parasitoids and parasites have been recorded from 14 families, nine in Hymenoptera, four in Diptera and one in Nematoda (Table 1). Ichneumonids and braconids were the most diverse families in Hymenoptera, represented by 36 and 28 species, respectively. The most diverse family in Diptera, as well as overall, was Tachinidae with 55 species.

Distribution among the taxa was highly variable. Three of the 14 families were only reported from one country, and five of the 14 families were reported from two or fewer countries. Bethyloid and bombyliid parasitoids were only reported in the United States (US) during the 1920s and 1940s (Table 1). Of the ≈ 150 species, 74 were reported from only one country, and 102 were reported from only one geographical region. In contrast, nine of the 14 families and 18 of the ≈ 150 species were reported from all geographical regions (North America, Central America and the Caribbean, and South America), and eight species were reported from 10 or more countries (Tables 1 and 2).

The number of parasitoid and parasite species reported from each FAW stage attacked in each country is summarized in Table 3. The highest number of parasitoid species attacking FAW larvae was reported from the US (75), followed by Brazil (45), Honduras (43) and Nicaragua (42).

Parasitoid species that attack FAW eggs were more prevalent in Nicaragua (11) Brazil (8), Barbados (7) and Mexico (6). Parasitoid species that attack FAW pupae were reported from only 4 countries, with the greatest number of these species collected from the US (6).

Parasitized FAW were collected from 19 different host plants (Table 4). FAW collected from corn were parasitized by species from all families except Bombyliidae. The number of parasite and parasitoid species attacking FAW was greatest for FAW collected from corn (134), sorghum (40), cotton (28), peanut (24), alfalfa (17), rice (13), and Bermuda grass (11). Six of the 14 parasitoids and parasite families were recovered from FAW that had been collected from a single host plant species. Ten of the 14 parasitoids and parasite families were recovered from FAW that had been collected from three or fewer host plant species. The most diverse parasitoid families also were recovered from more FAW host plant species. Tachinids, braconids, eulophids, and ichneumonids were recovered from FAW collected from 15, 11, 8 and 8 FAW host plant species, respectively.

A considerable number of parasitoid species ($\approx 33\%$) were reported only from one geographical region (Table 1). These findings emphasize the need for more surveys and taxonomical studies of the natural enemies in the different habitats of the natural distribution of the pest (Molina-Ochoa et al. 2001). It is important to consider the different developmental stages of the pest and the growing season of the crop to determine if differ-

TABLE 3. NUMBER OF SPECIES OF PARASITOIDS AND STAGE HOST ATTACKED REPORTED FROM DIFFERENT COUNTRIES.

Country	Number of parasitoid species			
	Egg	Larval	Pupae	Adult
Antigua	1			
Argentina	4	28	1	
Bahamas				1
Barbados	7	9	1	
Belize				1
Bermuda				1
Bolivia		1		
Brazil	8	45		
Cayman Is.				1
Canada		1		
Chile	3	19		
Colombia	5	6		1
Costa Rica				
Cuba	3	11		
Dominica				1
Dominican Republic	1			
Ecuador		2		
El Salvador				1
French Guinea				1
Grenada				1
Guadeloupe	3	2		1
Guatemala		1		
Guyana	2	2		
Haiti	1			
Honduras	5	43	1	1
Lesser Antilles	2	8		
Martinique				1
Mexico	6	22		
Nicaragua	11	42		
Panama		1		
Peru	2	10		
Puerto Rico	3	9		1
Suriname	2	4		1
Tobago	1	1		
Trinidad	4	9		1
United States	3	75	7	1
Uruguay	1	16		
Venezuela	2	29		
Virgin Is.				1

ences in natural distribution of the parasitoids exist (Ashley 1979; Molina-Ochoa et al. 2001) or if the records related to unique occurrence are due to inadequate surveys. It also is important to determine if the occurrence of FAW parasitoids is associated with the developmental stage of the pest and the host plant. Studies related to tritrophic interactions are needed to elucidate the role of the plant metabolites on the susceptibility of the pest to parasitoids and pathogens, as well as their suitability (Molina-Ochoa et al. 1999). A high diversity of FAW parasitoids has been reported in its natural distribution exerting significant mortal-

ity on egg and larval populations. An understanding of induced parasitization of the complex of parasitoids is needed to determine the species with higher capability to attack each stage of the FAW. It is necessary to unite institutional efforts to establish programs of release of FAW parasitoids in overwintering areas and to reduce the migration of the pest northward (Gross & Pair 1986). Because much of the published work does not include environmental data with the collections, we encourage authors to include this information in the future. These data would help select candidate parasitoids for a specific or broad region.

TABLE 4. NUMBER OF PARASITOID SPECIES IN EACH TAXON REPORTED FROM FAW COLLECTED FROM DIFFERENT HOST PLANTS.

Order	Family	FAW host plants																		
		A	BG	C	M	Mi	O	P	Pb	PG	Pm	PN	R	S	SC	SY	T	Te	To	Ve
Diptera																				
	Bombyliidae		1																	
	Phoridae				1															
	Sarcophagidae				4															
	Tachinidae	6	4	7	34	4	1	1				15	4	13	3	5	1	1		1
Hymenoptera																				
	Bethylidae				1															
	Braconidae	7	4	12	24	2				2		4	5	11		1			1	
	Chalcididae			3	4															
	Eulophidae		1	1	9				1		1	1	2	2						
	Ichneumonidae	4	1	5	36	1				1		4		11						
	Perilampidae				1															
	Pteromalidae				2															
	Scelionidae				2							1	1							
	Trichogrammatidae				5							1	1							
Nematoda																				
	Acuguttuidae				1									1						

A = Alfalfa, BG = Bermuda grass, C = Cotton, M = Maize, Mi = Millet, O = Onion, P = Hot Pepper, Pb = *Panicum barbinode*, PG = Para grass, Pm = *P. maximum*, PN = Peanut, R = Rice, S = Sorghum, SC = Sugarcane, SY = Soybean, T = Tomato, Te = Teosinte, To = Tobacco, Ve = Vegetable crops

Parasites and Parasitoids of *Spodoptera frugiperda* Pupae and Adults

Spodoptera frugiperda pupae and adults are attacked by several parasite and parasitoid species. Five species of Ichneumonidae: *Diapetimorpha introita* (Cresson), *Cryptus albitarsis* (Cresson), *Ichneumon promissorius* (Erichson), *Ichneumon ambulatorius* and *Vulgichneumon brevicinctor* (Say) have been reported attacking pupae of *S. frugiperda* and other noctuids (Bechinski & Pedigo 1983; Pair & Gross 1984; Wilson 1983; Pair & Gross 1989; Fitt & Daly 1990; Pavuk & Stinner 1991). Two generalist pupal parasitoids of Chalcididae, *Brachymeria ovata* (Say) and *B. robusta* (Cresson), have been collected from *S. frugiperda* (Wilson 1923; Luginbill 1928; Parker et al. 1953; Ashley 1979; Virla et al. 1999). Only one eulophid species, *Trichospilus pupivora* (Ferriere), has been reported as a generalist parasitoid of Lepidopteran pupae, including *S. frugiperda* (Alam 1979, Anantanarayanan 1934). Also, only one species (the ectoparasitic nematode, *Noctuidonema guyanense* Remillet & Silvain (Remillet & Silvain 1988)), has been observed parasitizing *S. frugiperda* adults.

Studies on *Diapetimorpha introita*

Diapetimorpha introita, a parasitoid which attacks FAW in the pupal stage, was reported for first time by Pair & Gross (1984) in Tifton, Georgia. The levels of parasitization in six trials ranged from 0.0 to 23.7%, and the percent of intact pupae that were parasitized during that study averaged 13.5%. Later, Gross & Pair (1986) emphasized the need for more efforts to explore the role of species of parasitoids that employ similar strategies. Pair & Gross (1989) reported the seasonal incidence of *D. introita* with rates of parasitism that averaged 5.2% (range 0-23.7%) and 8.4% (range 0-50.0%), respectively, during 1983 and 1984, with the highest rate occurring during September to November of each year. Because male *D. introita* are attracted to chemicals emitted by the female wasps (Jewett & Carpenter 1998) sticky traps baited with live female wasps were used to study the seasonal abundance of *D. introita* (Jewett & Carpenter 2001). The highest number of adult males was caught during early autumn which corresponded to the rates of parasitism reported by Pair and Gross (1989). As the investigation of their importance to biological control of *Spodoptera* spp. proceeds, more convenient methods of monitoring *D. introita* in the field are needed.

Pair (1995) studied the biology and rearing of *D. introita* on host and non-host noctuid pupae, *Spodoptera* spp., *Helicoverpa zea* (Boddie), and *Heliothis virescens* (F.) to identify factors that influence the reproduction and developmental rate of *D. introita*. This parasitoid was successfully

reared in the laboratory on *S. frugiperda* pupae. Carpenter & Greany (1998) compared the developmental time, weight, fecundity, longevity, and ability to parasitize hosts for *D. introita* wasps developing on artificial diet and wasps reared on *S. frugiperda* pupae. They conclude that the ability to rear *D. introita* on an inexpensive, artificial diet significantly enhances the potential of mass rearing this parasitoid for inundative releases against species in the genus *Spodoptera*.

Studies on *Ichneumon promissorius* and Collection of *Ichneumon ambulatorius*

I. promissorius was collected in Australia from *Helicoverpa armigera* (Hübner), and *H. punctigera* (Wallengren) pupae (Chadwick & Nikitin 1976; Wilson 1983; Fitt & Daly 1990). The parasitoid was imported into the US and released in Arkansas, Georgia, Oklahoma, and Texas from 1992 to 1997 (J. E. C. et al., unpublished data). Following releases of *I. promissorius* in ear-stage corn in the lower Rio Grande valley in Texas during 1993 and 1994, feral noctuid pupae, including FAW, were removed from the soil. During 1993, 575 FAW pupae were collected, 8 of which were parasitized by *I. promissorius*. Of the 13 FAW pupae collected in 1994, none were parasitized by *I. promissorius*. A similar study was conducted in a cornfield in Rabun County, Georgia. During 1993, 141 of the 300 FAW pupae collected were parasitized by *I. promissorius* and 24 were parasitized by *I. ambulatorius*, a new host record. Only 5 FAW pupae collected in 1994. Two of these FAW pupae were parasitized by *I. promissorius* and one was parasitized by *I. ambulatorius*.

Carpenter et al. (1994) compared several indigenous lepidopteran species in Tifton, GA, for *I. promissorius* acceptance and development. Pupae of *H. zea*, *H. virescens* (F.), *H. subflexa* (Guenée), *S. frugiperda*, *S. exigua* (Hübner), *Trichoplusia ni* (Hübner), *Agrotis ipsilon* (Hufnagel), and *Anticarsia gemmatilis* (Hübner) were used as hosts. Ninety pupae/species were tested, resulting in 74, 72, 68, 66, and 62 wasps emerging from *T. ni*, *A. ipsilon*, *H. subflexa*, *S. exigua*, and *S. frugiperda*, respectively. Lowest emergence of wasps was obtained on *A. gemmatilis* (6). The most acceptable hosts of *I. promissorius* were *H. zea* and *H. virescens*, as expected because *I. promissorius* was collected in Australia from heliothid species (Fitt & Daly 1990). Carpenter et al. (1994) considered that *H. zea* should be the primary host species of this pupal parasitoid in the US.

Carpenter (1995) examined the influence of host species, host availability, and mating status on *I. promissorius* fecundity and oviposition. The results from this study suggest that the females budget their energy expenditures and regulate oogenesis to maximize their reproductive potential. Host size, as well as the host species, may

have contributed to differences in weight of the wasps, and contributed to differences in fecundity, longevity, and oviposition between wasps reared on *S. exigua* and *H. zea* pupae. Virgin females reared on *S. exigua* pupae laid fewer eggs than virgin females reared on *H. zea* pupae.

Studies on *Vulgicheimon brevicinctor*
and *Cryptus albitarsis*

Two *V. brevicinctor* were recovered from FAW pupae collected in 1982 from a cornfield near Tifton, GA (Pair & Gross 1989). Five *V. brevicinctor* were recovered from a sample of 300 FAW pupae collected from a corn field in Rabun County, GA, in 1993, and one *V. brevicinctor* was recovered from a sample of 3 FAW pupae collected from the same field in 1994 (J. E. C., unpublished data). *V. brevicinctor* also has been reported from other noctuids and species of other lepidopteran families (Carlson 1979). For example, Bechinski & Pedigo (1983) studied the population dynamics of the green cloverworm (*Plathypena scabra* F.) in soybeans in Iowa during 1979 and 1980. They and found that *V. brevicinctor* acted in a delayed density-dependent manner on green cloverworm pupal mortality. Pavuk & Stinner (1991) reported that *V. brevicinctor* was reared from pupae of an arctiid, *Cisseps fulvicollis* (Hübner), and pupae from a pyralid, *Ostrinia nubilalis* (Hübner).

Pair & Gross (1989) collected *C. albitarsis* in Tifton, GA, from a single sample taken November 2, 1984. Four *C. albitarsis* also were recovered from a sample of 300 FAW pupae collected from a corn field in Rabun County, GA, in 1993 (J. E. C., unpublished data). Pair & Gross (1989) reported that *C. albitarsis* had been established in a laboratory colony in Tifton, GA. *C. albitarsis* cohorts from this laboratory colony were successfully reared on an artificial diet devoid of any host components (Greany and Carpenter 1998).

Studies on Ectoparasitic Nematodes Attacking Adults
of *Spodoptera frugiperda*

Remillet & Silvain in 1982 discovered and reported an ectoparasitic nematode, *Noctuidonema guyanense* Remillet & Silvain infecting *Spodoptera androgea* (Cramer) in French Guiana. It was subsequently described as a new genus and species (Remillet & Silvain 1988) most commonly found on moths of FAW, *Spodoptera latifascia* (Walker), *S. marima* (Schaus), *Anicla infecta* (Ochsenheimer), and *Leucania* spp. (Remillet & Silvain 1988).

Rogers et al. (1990a) determined the life cycle and host range for *N. guyanense* in French Guiana. They collected moths using a white sheet illuminated by UV light, by pheromone traps, and sweeping vegetation from a variety of habitats in Northeastern French Guiana. Moths in five families, Lasiocampidae, Noctuidae, Notodontidae,

Pyralidae, and Sphingidae were naturally infested with this nematode species. Twenty-five species of Noctuidae were infected by *N. guyanense*, the hosts most commonly infected were *Lesmone formularis* (Hübner), *S. dolichos* (F.), *S. frugiperda*, and *Xanthopastis timais* (Cramer) (Rogers et al. 1990a). Using the same methodologies, Rogers et al. (1990b) determined that multiple species of *Mocis* and *Spodoptera* were parasitized by the nematode in Florida and Georgia, and it was the first record of this parasite in North America. Simmons & Rogers (1990a) determined the distribution and prevalence of the nematode in tropical Americas, occurring in northern South America, Mexico, Texas, Florida, Bermuda, most of the Caribbean basin countries, and Central America.

Since 1990, numerous studies have been conducted on this nematode including biology (Simmons & Rogers 1994; Marti & Rogers 2000), pathological effects on the host (Marti et al. 1990; Rogers et al. 1993), infestation dynamics (Rogers & Marti 1992a; Silvain & Remillet 1993; Rogers & Marti 1993b), geographical distribution (Rogers et al. 1993; Rogers et al. 1991; Simmons et al. 1991; Rogers et al. 1997; Marti et al. 2000), ecology (Silvain & Remillet 1993), population profiles (Rogers & Marti 1992a, 1994), mating behavior (Simmons & Marti 1992), prevalence (Simmons & Rogers 1990b), maintenance of colonies (Rogers & Marti 1993a, b), host range (Simmons & Rogers 1996; Rogers & Marti 1996; Marti et al. 2000), and the bionomics of host insects of *N. guyanense* (Rogers et al. 1996; Marti et al. 2000). Subsequently however, a study on the speciation in Acuturidae shows that *N. guyanense* is limited to the lepidopteran genera *Spodoptera* and *Pseudaletia* (Marti et al. 2002).

Simmons et al. (1991) studied the seasonal chronology of the nematode in the tropical and subtropical America, and determined that host parasitism and nematode population density varied among locations and over time. Examples include higher parasitism on male moths in Grenada (77%) than in Texas (1%), and higher nematode populations at lower latitudes than higher latitudes. Simmons & Rogers (1991) also studied the dispersal and seasonal occurrence of *N. guyanense* on FAW adults in the US. They found that nematode populations and parasitism of FAW males were higher in Eastern States than in the Plains, Midwestern, and Central states, and that the percent parasitism and the number of nematodes per infested FAW changed over time at each location.

ACKNOWLEDGMENTS

The authors thank J. J. Hamm and O. G. Marti (USDA-ARS-Crop Protection & Management Research Laboratory, Tifton, GA), for their critical review of earlier versions of this manuscript, and thank O. G. Marti for providing information on ectoparasitic nematodes.

The authors appreciate the assistance of Duncan McClusky (University of Georgia Libraries, Tifton, GA), and Marco Antonio Mellín-Rosas (CNRCB, Tecmán, Colima, México) in obtaining scientific literature. The authors gratefully acknowledge and thank Susan Drawdy (USDA-ARS-Crop Protection & Management Research Laboratory, Tifton, GA) for her exceptional technical assistance and dedication to this project. Insect identifications and verification of insect classifications were provided by several scientists associated with the USDA/ARS Systematic Entomology Laboratory, including B. Carlson, E. E. Grissell, P. Marsh, N. E. Woodley, and N. W. Gates. This paper is a contribution of the University of Nebraska Agricultural Research Division, Lincoln, NE 68583. Journal Series No. 13681, Department of Entomology, University of Nebraska-Lincoln. The authors thank Universidad de Colima, and CONACYT-Mexico for supporting the senior author.

REFERENCES CITED

- ALAM, M. M. 1979. Attempts at the biological control of major insect pests of maize in Barbados, W. I. Symp. On maize and peanut. Paramaribo, Suriname, Nov. 13-18, 1978. Proc. Caribbean Food Crops Soc. 15: 127-135.
- ALLEN, H. W. 1921. Notes on a bombylid parasite and a polyhedral disease of the southern grass worm, *La-phygma frugiperda*. J. Econ. Entomol. 14: 510-511.
- ALTIERI, M. A. 1991. Increasing biodiversity to improve insect pest management in agro-ecosystems, Chapter XIV, pp. 165-182. In D. L. Hawksworth [ed.] The Biodiversity of Microorganisms and Invertebrates: Its Role in Sustainable Agriculture. CAB International, Wallingford, UK.
- ALVAREZ, L. R., Y F. G. ROA. 1995. Comportamiento parasítico de *Telenomus* sp. en *Spodoptera frugiperda*. Rev. Col. Entomol. 21: 191-197.
- ANANTANARAYANAN, K. P. 1934. On the bionomics of a eulophid *Trichospilus pupivora* Ferr.), a natural enemy of the coconut caterpillar (*Nephantis serinopa*, Meyr.) in South India. Bull. Ent. Res. 25:55-61.
- ANDREWS, K. L. 1988. Latin American research on *Spodoptera frugiperda* (Lepidoptera: Noctuidae). Florida Entomol. 71: 630-653.
- ARCE, G. F., Y J. GARCÍA G. 1995. Parasitismo natural y biología de *Chelonus insularis* Cresson (Hymenoptera: Braconidae) sobre *Spodoptera frugiperda* (J.E. Smith) Lepidoptera: Noctuidae, en los valles centrales de Oaxaca, pp. 48-49. En XVIII Congreso Nacional de Control Biológico. Soc. Mex. Contr. Biol. Noviembre de 1995. Tapachula, Chiapas, México.
- ARMAS-GARCÍA, J. L., Y J. L. AYALA-SIFONTES. 1987. Parásitos de huevos de la palomilla del maíz, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). Centro-Agrícola 14: 88-89.
- ASHLEY, T. R. 1979. Classification and distribution of fall armyworm parasites. Florida Entomol. 62: 114-123.
- ASHLEY, T. R., E. R. MITCHELL, N. C. LEPPLA, AND E. E. GRISSELL. 1980. Parasites attacking fall armyworm larvae *Spodoptera frugiperda* in late planted field corn. Florida Entomol. 63: 136-142.
- ASHLEY, T. R., V. H. WADDILL, E. R. MITCHELL, AND J. RYE. 1982. Impact of native parasites of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), in South Florida and release of the exotic parasite, *Eiphosoma vitticole* (Hymenoptera: Ichneumonidae). Environ. Entomol. 11: 833-837.
- ASHLEY, T. R. 1983. Growth pattern alterations in fall armyworm, *Spodoptera frugiperda* larvae, after parasitization by *Apanteles marginiventris*, *Campoletis grioti*, *Chelonus insularis*, and *Eiphosoma vitticole*. Florida Entomol. 66: 260-266.
- ASHLEY, T. R., C. S. BARFIELD, V. H. WADDILL, AND E. R. MITCHELL. 1983. Parasitization of fall armyworm larvae on volunteer corn, bermuda grass, and paragrass. Florida Entomol. 66: 267-271.
- ASHLEY, T. R. 1986. Geographical distributions and parasitization levels for parasitoids of the fall armyworm, *Spodoptera frugiperda*. Florida Entomol. 69:516-524.
- AVALOS, D. S. 1988. Moscas tachinidae de la provincia de Córdoba (Argentina). Rev. Peruana Entomol. 31: 48-50.
- AYALA-SIFONTES, J. L., M. GONZÁLEZ-ALVAREZ, H. GRILLO-RAVELO, Y S. CABALLERO-FIGUEROA. 1978. Observaciones preliminares sobre el ciclo biológico de *Diadegma* sp., parásito de *Heliothis virescens* (Fabricius). Centro Agrícola 5: 1-13.
- AYALA-SIFONTES, J. L., J. RAMÍREZ-OBREGÓN, S. RODRÍGUEZ-CUELLAR, Y J. L. ARMAS-GARCÍA. 1988. Hábitos oviposicionales de *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) en plantas de arroz. Rev. Centro-Agrícola 15: 11-15.
- AYALA-SIFONTES, J. L. ARMAS-GARCÍA, T. RODRÍGUEZ, Y M. BORGES-SOTO. 1992. Parasitismo de las puestas de *Spodoptera exigua* (Hubner) por *Telenomus* sp. en condiciones de laboratorio y campo. Rev. Centro-Agrícola 19: 42-46.
- AYQUI-VILCA, S. 1993. Morfología y biología de *Campoletis curvicauda* (Hymenoptera: Ichneumonidae) parasitoide de *Spodoptera frugiperda*. Rev. Peruana Entomol. 35: 31-36.
- BAHENA, J. F., Y O. GARCÍA M. 1991. Enemigos naturales del gusano cogollero del maíz *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) en Morelos, México, pp. 54-61. En XIV Congreso Nacional de Control Biológico. Soc. Mex. Contr. Biol. Octubre 10-11, 1991. Buenavista, Saltillo, Coahuila, México.
- BARILLA-VERA, M. 1989. Técnica de cría del gusano cogollero y sus parasitoids, pp. 59-62. En XII Reunión Nacional de Control Biológico. Secretaría de Agricultura y Recursos Hidráulicos-DGSV, Noviembre 22-24, 1989, Torreón, Coahuila, México.
- BECHINSKI, E. J., AND L. P. PEDIGO. 1983. Green cloverworm (Lepidoptera: Noctuidae) population dynamics: Pupal life table studies in Iowa soybeans. Environ. Entomol. 12: 656-661.
- BENZING, A., R. G. KLEESPIES, Y F. PONCE. 2000. Mortalidad natural de larvas de Noctuidae (Lepidoptera) en los Andes ecuatorianos: un primer acercamiento. Rev. Colombiana Entomol. 26: 57-60.
- BIANCHI, F. A. 1944. The recent introduction of armyworm (*S. exempta*, *S. exigua*, *S. frugiperda*) parasites from Texas. Hawaii Planter's Rec. 48: 203-212.
- BLANCHARD, E. E. 1963. Dípteros parásitos de Noctuidae argentinos. Rev. Inv. Agrop. INTA 17: 129-254.
- BRUNER, S. C., L. C. SCARAMUZZA, AND A. R. OTERO. 1975. Catálogo de los insectos que atacan a las plantas económicas de Cuba. 2nda. Edición. Academia de Ciencias de Cuba. Instituto de Zoología. 401 pp.
- BUTLER, G. D., JR. 1958a. Tachinid flies reared from lepidopterous larvae in Arizona, 1957. J. Econ. Entomol. 51: 561-62.
- BUTLER, G. D., JR. 1958b. Braconid wasps reared from lepidopterous larvae in Arizona, 1957. Pan-Pac. Ent. 34: 221-223.

- CAMPOS, L. 1965. Investigaciones sobre el control biológico del "cogollero" del maíz, *Spodoptera frugiperda* (J.E. Smith) y otros noctuideos. *Rev. Peruana Entomol.* 8:126-131.
- CANAS, L. A., AND R. J. O'NEIL. 1998. Applications of sugar solutions to maize, and the impact of natural enemies on fall armyworm. *Int. J. Pest Management* 44: 59-64.
- CANSECO-ROMÁN, O. 1988. Método para la cría de *Spodoptera frugiperda* (Smith) y sus parasitoides *Telenomus* spp. *En XI Reunión Nacional de Control Biológico*. Secretaría de Agricultura y Recursos Hidráulicos, Agosto 15-17, 1988. Hermosillo, Sonora. 11 pp.
- CARLSON, R. W. 1979. Ichneumonidae, pp. 315-741 *In* Krombein et al., *Catalog of Hymenoptera in America north of Mexico*. Smithsonian Institution Press, Washington.
- CARPENTER, J. E., S. D. PAIR, AND G. P. FITT. 1994. *Ichneumon promissorius* (Hymenoptera: Ichneumonidae): Development on North American hosts. *J. Econ. Entomol.* 87: 929-932.
- CARPENTER, J. E. 1995. *Ichneumon promissorius* (Erichson) (Hymenoptera: Ichneumonidae): Factors affecting fecundity, oviposition, and longevity. *J. Entomol. Sci.* 30: 279-286.
- CARPENTER, J. E., AND P. D. GREANY. 1998. Comparative development and performance of artificially reared versus host-reared *Diapetimorpha introita* (Cresson) (Hymenoptera: Ichneumonidae) wasps. *Biological Control* 11: 203-208.
- CARRILLO, H. 1980. Determinación del parasitismo natural de gusano cogollero, *Spodoptera frugiperda* (J.E. Smith) en Quintana Roo. *Folia Entomol. Méx.* 45: 111-112.
- CASTRO, M. T., H. N. PITRE, AND D. H. MECKENSTOCK. 1989. Populations of fall armyworm, *Spodoptera frugiperda* (J.E. Smith), larvae and associated natural enemies in sorghum and maize cropping systems in Southern Honduras. *Trop. Agric.* 66: 259-264.
- CAVE, R. D. 1993. Parasitoides larvales y pupales de *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) en Centro América con una clave para las especies encontradas en Honduras. *Ceiba* 34: 33-56.
- CAVE, R. D., Y N. M. ACOSTA. 1999. *Telenomus remus* Nixon: un parasitoide en el control biológico del gusano cogollero, *Spodoptera frugiperda* (Smith). *Ceiba* 40: 215-227.
- CHADWICK, C. E., AND M. I. NIKITIN. 1976. Records of parasitism in the families Ichneumonidae, Braconidae and Aulacidae. *J. Entomol. Soc. Aust. (N.S.W.)* 9: 28-38.
- CICEROS, O. J., J. M. MORALES, W. H. KING C., Y G. SÁNCHEZ, G. 1995. *Meteorus laphygmae* (Viereck) endoparasito del gusano cogollero en sorgo, pp. 56. *En XVIII Congreso Nacional de Control Biológico*. Soc. Mex. Contr. Biol. Noviembre de 1995. Tapachula, Chiapas, México.
- CLARIDGE, M. F. 1991. Genetic and biological diversity of insect pests and their natural enemies. Chapter XV, pp. 183-194 *In* D. L. Hawksworth [ed.]. *The Biodiversity of Microorganisms and Invertebrates: Its role in Sustainable Agriculture*. CAB International, Wallingford, UK.
- CORONADO, B. J. M., Y E. RUÍZ, C. 1991. Bracónidos (Hymenoptera) de la Reserva "El Cielo" de Tamaulipas, pp.72-76. *En XIV Congreso Nacional de Control Biológico*. Soc. Méx. Contr. Biol. Octubre 10-11, 1991. Buenavista, Saltillo, Coahuila, México.
- CORREA-FIGUEIREDO, M. DE L. C., I. CRUZ, E T. M. C. D. LUCIA. 1999. Controle integrado de *Spodoptera frugiperda* (Smith & Abbott) utilizando-se o parasitoide *Telenomus remus* Nixon. *Pesq. Agrop. Bras.* 34: 1975-1982.
- CORTES, R. 1980. Neotropical Tachinidae (Diptera) I. Notes, records, distribution and descriptions. *Rev. Bras. Entomol.* 24: 105-110.
- CORTEZ, M. H., Y J. TRUJILLO A. 1994. Incidencia de gusano cogollero y sus enemigos naturales en tres agro-sistemas de maíz. *Turrialba* 44: 1-9.
- COSTA-LIMA, A. 1949. *Insetos do Brasil*, 6to. Tomo. Escola Nacional de Agronomia. Seria Didatica No. 8: 184-185.
- COSTA-LIMA, A. 1962. *Insectos do Brasil*. XII. Himenópteros. Rio de Janeiro, Escola Nac. Agr., 1962. 393 p. (Ser. No. 14).
- CRESSON, E. T. 1869. List of the North American species of the genus *Aleiodes* Wesmael. *Trans. Am. Entomol. Soc.* 2: 377-382.
- CRUZ, I., M. L. C. FIGUEIREDO, E. P. GONCALVES, D. A. N. LIMA, E E. E. DINIZ. 1997a. Efeito da idade de lagartas de *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) no desempenho do parasitoide *Camptoclis flavipectus* (Ashmead) (Hymenoptera: Ichneumonidae) e consumo foliar por lagartas parasitadas e nao-parasitadas. *An. Soc. Entomol. Brasil* 26: 229-234.
- CRUZ, I., M. L. C. FIGUEIREDO, F. H. VALICENTE, E A. C. OLIVEIRA. 1997b. Application rate trials with a nuclear polyhedrosis virus to control *Spodoptera frugiperda* (Smith) on maize. *An. Soc. Entomol. Brasil* 26: 145-152.
- CRUZ, I., M. L. C. FIGUEIREDO, E M. J. MATOSO. 1999. Controle biológico de *Spodoptera frugiperda* utilizando o parasitoide de ovos *Trichogramma*. Circular Técnica No. 30, Centro Nacional de Pesquisa de Milho e Sorgo (CNPMS), Sete Lagoes, Brazil, 40 pp.
- CURRAN, C. H. 1927. A new tachinid parasitic on armyworm in Mexico. *Proc. Hawaii Ent. Soc.* 6: 497-498.
- DANKS, H. V. 1975. Factors determining levels of parasitism by *Winthemia rufopicta* (Diptera: Tachinidae), with particular reference to *Heliothis* spp. (Lepidoptera: Noctuidae) as hosts. *Can. Ent.* 107: 655-684.
- DE SA, L. A. N., AND J. R. P. PARRA. 1994. Natural parasitism of *Spodoptera frugiperda* and *Helicoverpa zea* (Lepidoptera: Noctuidae) eggs in corn by *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae) in Brazil. *Florida Entomol.* 77: 185-188.
- DE SANTIS, L., Y L. ESQUIVEL. 1966. Tercera lista de himenópteros parásitos de los insectos de la República Argentina. *Rev. Mus. La Plata* 9: 47-217.
- DE SANTIS, L. 1967. Catálogo de los himenópteros argentinos de la serie parasítica, incluyendo Bethyloidea. *Com. Invest. Cient. Prov. De Buenos Aires, Gobernación*. La Plata, 337 p.
- DE SANTIS, L. 1979. Catálogo de los himenópteros Calcidoideos de América al Sur de los Estados Unidos. *Com. Invest. Cient. Prov. De Buenos Aires, Gobernación*, 488 p.
- DE SANTIS, L. 1980. Nueva sinonimia, nueva combinación y nuevas citas de Himenópteros Calcidoideos para la República Argentina. *Neotrópica* 26:153-154.
- DE SANTIS, L. 1989. Catálogo de los Himenópteros Calcidoideos (Hymenoptera) al Sur de los Estados Unidos. Segundo Suplemento. *Acta Ent. Chilena* 15: 9-90.
- DE SANTIS, L., Y P. FIDALGO. 1994. Catálogo de los Himenópteros Calcidoideos de América al Sur de los

- Estados Unidos. Tercer Suplemento. Acad. Nac. Agron. Vet. (Buenos Aires, Argentina) 13: 154 pp.
- DENT, D. 2000. Insect pest management. 2nd Ed. CABI Publishing, University Press, Cambridge, UK, 410 pp.
- DEW, J. A. 1913. Fall armyworm *Laphygma frugiperda* (S&A). J. Econ. Entomol. 6: 361-366.
- ENKERLIN, D. 1975. Review of *Spodoptera* in Latin America. Summary of research at Monterrey Tech. Inst., N.L. Mexico. 13 pp.
- ESCALANTE G., J. A. 1974. Contribución al conocimiento de la biología de *Heliothis zea* y *Spodoptera frugiperda*, en el Cusco. Rev. Peruana Entomol. 17: 121-122.
- ESTRADA R., F. A. 1960. Lista preliminar de insectos asociados al maíz en Nicaragua. Turrialba, 10: 68-73.
- ETCHEVERRY, M. 1957. *Laphygma frugiperda* (Abbot and Smith) en Chile (Lepidoptera: Noctuidae). Rev. Chilena Entomol. 5: 183-192.
- FENNAH, R. G. 1947. The insect pests of food-crops in the Lesser Antilles. Dept. Agric. Windward IS., St. George's Grenada, B.W.I. pp. 77-79.
- FERNÁNDEZ, B. R. I., Y A. S. CLAVIJO. 1984. Efectos de dos insecticidas (uno químico y otro biológico) sobre el parasitismo observado en larvas de *Spodoptera frugiperda* (S.) provenientes de parcelas experimentales de maíz. Rev. Fac. Agron. Univ. Central Venezuela 13: 101-109.
- FERNÁNDEZ, B. R. I., Y B. J. B. TERÁN. 1990a. Biología de *Meteorus laphygmae* Viereck (Hymenoptera: Braconidae). Parte I. Fases de huevo, larva y pupa. Rev. Fac. Agron. Univ. Central Venezuela 16: 177-198.
- FERNÁNDEZ, B. R. I., Y B. J. B. TERÁN. 1990b. Biología de *Meteorus laphygmae* Viereck (Hymenoptera: Braconidae). Parte II. Adulto, partenogénesis e interrelaciones parásito-hospedero. Rev. Fac. Agron. Univ. Central Venezuela 16: 199-206.
- FERRER, W. F. 1998a. Alcances sobre el control biológico en la región andina de Sudamérica, pp. 25-26 En Aportes del control biológico en la agricultura sostenible. I Congr. Latinoam. Sec. Reg. Neotrop. Org. Inter. Lucha Biol. Mayo 18-22, 1998, Lima, Perú.
- FERRER, W. F. 1998b. El control biológico en el control de plagas de importancia económica en maíz y caña, pp. 48-49. En Aportes del control biológico en la agricultura sostenible. I Congr. Latinoam. Sec. Reg. Neotrop. Org. Inter. Lucha Biol. Mayo 18-22, 1998, Lima, Perú.
- FITT, G. P., AND J. C. DALY. 1990. Abundance of overwintering pupae and the spring generation of *Helicoverpa* spp. (Lepidoptera: Noctuidae) in Northern New South Wales, Australia: implications for pest management. J. Econ. Entomol. 83: 1827-1836.
- FLORES-DÁVILA, M., L. A. AGUIRRE-URIBE, Y E. RUÍZ-CANCINO. 1991. Ichneumonidae (Hymenoptera) del Sureste de Coahuila, pp. 82-87 En XIV Congreso Nacional de Control Biológico. Soc. Mex. Contr. Biol. Octubre 10-11, 1991. Buenavista, Saltillo, Coahuila, México.
- GARCÍA-LAGUNAS, E. D. 1988. Evaluación del parasitoide *Telenomus solitus* y *Telenomus remus* (Hymenoptera: Scelionidae) sobre huevecillo de *Spodoptera frugiperda* (Smith) en condiciones de campo en Emiliano Zapata, Mor. En XI Reunión Nacional de Control Biológico. Secretaría de Agricultura y Recursos Hidráulicos, Agosto 15-17, 1988. Hermosillo, Sonora. 12 pp.
- GIRALDO-VANEGAS, H., Y J. L. GARCÍA R. 1992. Determinación del número de instares de *Eiphosoma vitticole* Cresson (Hymenoptera: Ichneumonidae). Bol. Entomol. Venezolana 7: 133-137.
- GIRALDO-VANEGAS, H., Y J. L. GARCÍA R. 1994a. Comportamiento, descripción y tiempo de desarrollo de los estadios inmaduros de *Eiphosoma vitticole* Cresson (Hymenoptera: Ichneumonidae), parásito de *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). Agron. Trop. Maracay 44: 645-665.
- GIRALDO-VANEGAS, H., Y J. L. GARCÍA R. 1994b. Comportamiento de los adultos de *Eiphosoma vitticole* Cresson (Hymenoptera: Ichneumonidae), parásito de *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae). Bol. Entomol. Venezolana 9: 15-20.
- GIRALDO-VARGAS, H., Y J. L. GARCÍA R. 1995. Influencia de la alimentación sobre la capacidad reproductiva de *Eiphosoma vitticole* Cresson (Hymenoptera: Ichneumonidae), parásito de *Spodoptera frugiperda* (J.E. Smith). Agron. Trop. Maracay 45: 159-170.
- GLADSTONE, S. H. 1991. Parásitos del cogollero, *Spodoptera frugiperda* Smith (Lepidoptera: Noctuidae) en maíz sembrado en la época seca en Nicaragua. Ceiba 32: 201-206.
- GONCALVES, C. R. 1973. Observacoes sobre hospedeiros de himenopteros da familia Ichneumonidae, no Brasil. An. Soc. Entomol. Brasil. 2: 31-36.
- GONCALVES, C. R., AND A. J. L. GONCALVES. 1973. Novas observacoes sobre insetos hospedeiros de moscas da familia Tachinidae (Diptera). Agronomia 31: 9-15.
- GONZÁLEZ-NARVAEZ, C. E., Y J. L. ZOCCO. 1996. Control integrado de *Spodoptera frugiperda* (Smith) utilizando *Telenomus remus* (Nixon) en *Zea mays* L. Rev. Invest. Agric. DANAC 1: 1-5.
- GREANY, P. D., AND J. E. CARPENTER. 1998. Culture medium for parasitic and predaceous insects. U.S. Patent No. 5799607.
- GROSS, H. R., JR., AND S. D. PAIR. 1986. The fall armyworm: Status and expectations of biological control with parasitoids and predators. Florida Entomol. 69:502-515.
- GROSS, H. R., JR., AND S. D. PAIR. 1991. Seasonal distribution, response and host developmental stage, and screened-cage performance of *Archytas marmoratus* (Diptera: Tachinidae) and *Ophion flavidus* (Hymenoptera: Ichneumonidae) on *Spodoptera frugiperda* (Lepidoptera: Noctuidae). Florida Entomol. 74: 237-245.
- GUAGLIAMI, P. 1962. Las plagas de la caña de azúcar en Venezuela. Maracay, Centro de Investigaciones Agronómicas. 1962. V.2, pp. 520-521. (Monografía 2).
- GUEVARA, M. A., C. G. PÉREZ, Y F. J. GUTIÉRREZ. 1979. Dinámica de poblaciones de insectos benéficos en el cultivo del maíz. VII Reunión Nacional de Control Biológico. Secretaría de Agricultura y Recursos Hidráulicos, Sanidad Vegetal. 24-26 de Abril de 1979, Veracruz, Veracruz, México, 7 pp.
- GUIMARÃES, J. H. 1971. "Fam. Tachinidae (Larvaevoridae)", In A catalogue of the Diptera of the America South of the United States. Mus. Zool. Univ. Sao Paulo 104: 1-333.
- GUIMARÃES, J. H. 1977. Host-parasite and parasite-host catalogue of South American Tachinidae (Diptera). Arq. Zool. Mus. Zool. Sao Paulo 28: 1-131.
- GUIMARÃES, J. H. 1983. Taxonomy of Brazilian flies of the genus *Lespesia* (Robineau-Desvoidy) (Diptera, Tachinidae). Papeis Avulsos de Zoologia 35: 11-30.
- GUTIÉRREZ-RODRÍGUEZ, F. J. 1982. Técnica para cría masiva de *Euplectrus plathypenae* (Howard) (Hymenoptera: Eulophidae) y su hospedero *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae), pp. 17-24. En X Reunión Nacional de Control Biológico.

- Grupo Sectorial Agropecuario y Forestal. Secretaría de Agricultura y Recursos Hidráulicos. Abril 1982, Durango, Durango, México.
- HAMM, J. J., L. D. CHANDLER, AND H. R. SUMNER. 1994. Field tests with a fluorescent brightener to enhance infectivity of fall armyworm (Lepidoptera: Noctuidae) nuclear polyhedrosis virus. *Florida Entomol.* 77: 425-437.
- HERNÁNDEZ, D., F. FERRER, Y B. LINARES. 1989. Introducción de *Telenomus remus* Nixon (Hym: Scelionidae) para controlar *Spodoptera frugiperda* (Lep.: Noctuidae) en Yaritagua-Venezuela. *Agron. Trop. Maracay* 39: 199-205.
- HERNÁNDEZ, D., Y F. DÍAZ. 1995. Efecto de la edad del parasitoide *Telenomus remus* Nixon (Hymenoptera: Scelionidae) sobre su capacidad de oviposición y proporción sexual de la descendencia. *Bol. Entomol. Venezolana* 10: 167-175.
- HERNÁNDEZ, D., Y F. DÍAZ. 1996a. Efecto de la edad del hospedero *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) sobre el parasitismo y la proporción sexual de la descendencia (psd) de *Telenomus remus* Nixon (Hymenoptera: Scelionidae). *Bol. Entomol. Venezolana* 11: 27-32.
- HERNÁNDEZ, D., Y F. DÍAZ. 1996b. Efecto de la temperatura sobre el desarrollo de *Telenomus remus* Nixon (Hymenoptera: Scelionidae) parasitoide de *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae). *Bol. Entomol. Venezolana* 11: 149-153.
- HERRERA-ARANGUENA, J. M. 1998. El control biológico en la represión de plagas de importancia económica en el algodónero, pp. 40-44. *En Aportes del control biológico en la agricultura sostenible. I Congr. Latinoam. Sec. Reg. Neotrop. Org. Inter. Lucha Biol.* Mayo 18-22, 1998, Lima, Perú.
- HOFMASTER, R. N., AND D. E. GREENWOOD. 1949. Fall armyworm control on forage and truck crops. *J. Econ. Ent.* 42: 502-506.
- HOGG, D. B., R. E. ANDERSON, AND H. N. PITRE. 1982. Early-season parasitization of fall armyworm (Lepidoptera: Noctuidae) larvae in Mississippi. *Florida Entomol.* 65: 584-585.
- HUIS, A. V. 1981. Integrated pest management in the small farmer's maize crop in Nicaragua. *Med. Landb. Wageningen* 81: 221 pp.
- HYNES, H. B. N. 1942. Lepidopterous pests of maize in Trinidad. *Tropical Agric.* 29: 194-202.
- IRVING, N. S. 1978. Report on the Antigua, Monserrat and St. Kitts-Nevis-Anguilla entomology programme. Centre for Overseas pest Research, London, United Kingdom, 55 pp.
- ISENHOUR, D. J. 1988. Interactions between two hymenopterous parasitoids of the fall armyworm (Lepidoptera: Noctuidae). *Environ. Entomol.* 17: 616-620.
- JEWETT, D. K., AND J. E. CARPENTER. 1998. Chemical communication between males and females of a pupal parasitoid *Diapetimorpha introita* (Hymenoptera: Ichneumonidae). *Ann. Entomol. Soc. Am.* 91: 748-753.
- JEWETT, D. K., AND J. E. CARPENTER. 2001. Seasonal abundance of pupal parasitoid, *Diapetimorpha introita* (Hymenoptera: Ichneumonidae). *Florida Entomol.* 84: 50-54.
- JOHANNES-TOONDERS, T., Y J. L. CARRILLO-SÁNCHEZ. 1987. Evaluación de la efectividad de *Trichogramma* spp. (Hymenoptera: Trichogrammatidae) en el combate de *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae). *Recomendaciones para su uso. Agrocencia* 67: 75-84.
- JONES, T. H. 1913. Some notes on *Laphygma frugiperda* S. and A. in Porto Rico. *J. Econ. Entomol.* 6: 230-236.
- KELLER, M. S. 1980. Effects of temperature and corn phenology on Fall Armyworm Biology. Masters Thesis, University of Florida, 83 pp.
- KING, A. B. S., AND J. L. SAUNDERS. 1984. Las plagas invertebradas de cultivos anuales alimenticios en América Central. Overseas Development Administration, London, 62 pp.
- LACAYO, L. 1977. Especies parásitas de *Spodoptera frugiperda* Smith, *Diatraea lineolata* (Wer.) y *Trichoplusia ni* (Hbn) en zonas de Managua y Masatepe. XXIII Reunión Annual del PCCMCA, Panamá, Panamá 1977, 28 pp.
- LEZAMA-GUTIÉRREZ, R., J. J. HAMM, J. MOLINA-OCHOA, M. LÓPEZ-EDWARDS, A. PESCADOR-RUBIO, M. GONZÁLEZ-RAMÍREZ, AND E. L. STYER. 2001. Occurrence of entomopathogens of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the Mexican states of Michoacan, Colima, Jalisco and Tamaulipas. *Florida Entomol.* 84: 23-30.
- LOYA-RAMÍREZ, J. 1978. Estudio sobre el gusano cogolero, *Spodoptera frugiperda* (J.E. Smith), en Morelos. Instituto Nacional de Investigaciones Agrícolas (INIA), México. Informe Técnico de la Coordinación del Apoyo Entomológico 3: 45-55.
- LUCCHINI, F., E A. A. ALMEIDA. 1980. Parasitas da *Spodoptera frugiperda* (Smith & Abbot, 1797) (Lep., Noctuidae), lagarta do cartucho do milho, encontrado em Ponta Grossa, PR. *An. Soc. Entomol. Brasil.* 9: 115-121.
- LUGINBILL, P. 1928. The fall armyworm. U.S. Dept. Agric. Tech. Bull. 34, 91 pp.
- MACHUCA, J. R., P. ARRETZ, Y J. ARAYA. 1989. Presencia de *Campoleptis sonorensis* (Cameron, 1986) (Hymenoptera: Ichneumonidae) en Chile. *Acta Entomol. Chilena* 15: 269-270.
- MAES, J. M. 1989. Catálogo de los insectos controladores biológicos en Nicaragua. Vol. III. Insectos Parasitoides. *Rev. Nicar. Entomol.* 10: 1-138.
- MALAUSSA, J. C. 1981. Etude de la dynamique des populations des chenilles de *Spodoptera frugiperda* Abbot & Smith et d'*Heliothis zea* Boddie (Lepidoptera: Noctuidae) sur deux varietes de mais en Guadeloupe (Antilles Francaises). *Agronomie* 1: 701-705.
- MALAUSSA, J. 1983. Etude bioecologique des noctuelles (Lepidoptera: Noctuidae) des Antilles Francaises. *Bol. Soc. Portuguesa Entomol.* 2: 49-67.
- MARÍN-ACOSTA, J. C. 1966. Nota sobre *Euplectrus platypeneae* HOW., ectoparásito de *Laphygma frugiperda* S. y A. y *Prodenia eridania* (Cramer). *Agron. Trop.* 16: 155-159.
- MARSH, P. M. 1978. The braconid parasites (Hymenoptera) of *Heliothis* species (Lepidoptera: Noctuidae). *Proc. Entomol. Soc. Wash.* 80: 15-36.
- MARSH, P. M., AND S. R. SHAW. 2001. Revision of North American *Aleiodes* Wesmael (Part 6): the Gasterator (Jurine) and Unipunctator (Thunberg) species-groups (Hymenoptera: Braconidae: Rogadinae). *Proc. Entomol. Soc. Wash.* 103: 291-307.
- MARTÍ, O. G., C. E. ROGERS, J. F. SILVAIN, AND A. M. SIMMONS. 1990. Pathological effects of an ectoparasitic nematode *Noctuidonema guyanense* (Nematoda: Aphelenchoididae) on adults of the fall armyworm (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.* 83: 956-960.
- MARTÍ, O. G. JR., AND C. E. ROGERS. 2000. Effect of *Noctuidonema guyanense* (Nematoda: Acugutturidae)

- on the longevity of feral male *Spodoptera frugiperda* (Lepidoptera: Noctuidae) moths. *J. Entomol. Sci.* 35: 259-266.
- MARTI, O. G. JR., B. LALANNE-CASSOU, J. F. SILVAIN, A. KERMARREC, AND A. M. SIMMONS. 2000. Ectoparasitic nematodes (Aphelenchoididae: Acugutturidae) of Lepidoptera and Blattodea in Guadeloupe. *Nematology* 2: 669-684.
- MARTI, O. G., JR., J. F. SILVAIN, AND B. J. ADAMS. 2002. Speciation in the Acugutturidae (Nematoda: Aphelenchida). *Nematology* 4: 489-504.
- MCCUTCHEON, G. S., W. Z. SALLEY, JR., AND S. G. TURNIPSEED. 1983. Biology of *Apanteles ruficrus*, an imported parasitoid of *Pseudoplusia includens*, *Trichoplusia ni*, and *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Environ. Entomol.* 12: 1055-1058.
- MCCUTCHEON, G. S. 1991. Late-season parasitoids of the fall armyworm in South Carolina. *J. Agric. Entomol.* 8: 219-221.
- MEDINA, T. M. C., D. P. CAMACHO, Z. J. E. LUQUE, Y P. A. SIABATTO. 1988. Ciclo de vida y descripción de *Chelonus insularis* Cresson (Hymenoptera: Braconidae), parásito de *Spodoptera* spp. *Rev. Colombiana Ent.* 14: 13-21.
- MILWARD-DE AZEVEDO, E. M. V., J. R. POSTALI PARRA, J. H. GUIMARAES, E R. DE PAULA-ALMEIDA. 1991a. Aspectos da biologia de *Archytas incertus* (Diptera: Tachinidae) e de suas inter-relacoes com *Spodoptera frugiperda* (Lepidoptera: Noctuidae). 1. Metodologia de criacao e determinacao do instar mais adequado para a producao do parasitoide. *Rev. Bras. Entomol.* 35: 485-497.
- MILWARD-DE AZEVEDO, E. M. V., J. H. GUIMARAES, E J. R. POSTALI-PARRA. 1991b. Aspectos da biologia de *Archytas incertus* (Diptera: Tachinidae) e de suas inter-relacoes com *Spodoptera frugiperda* (Lepidoptera: Noctuidae). 2. Carga parasitariae de regio inoculacao no corpo do hospedeiro. *Rev. Bras. Entomol.* 35: 499-507.
- MILWARD-DE AZEVEDO, E. M. V., E J. R. POSTALI-PARRA. 1991c. Aspectos da biologia de *Archytas incertus* (Diptera: Tachinidae) e de suas inter-relacoes com *Spodoptera frugiperda* (Lepidoptera: Noctuidae). 3. Influencia da temperatura na ontogenia do parasitoide. *Rev. Bras. Entomol.* 35: 509-516.
- MILWARD-DE AZEVEDO, E. M. V., E J. R. POSTALI-PARRA. 1991d. Aspectos da biologia de *Archytas incertus* (Diptera: Tachinidae) e de suas inter-relacoes com *Spodoptera frugiperda* (Lepidoptera: Noctuidae). 4. Sincronismo. *Rev. Bras. Entomol.* 35: 517-520.
- MILWARD-DE AZEVEDO, E. M. V., J. R. POSTALI PARRA, E J. H. GUIMARAES. 1991e. Aspectos da biologia de *Archytas incertus* (Diptera: Tachinidae) e de suas inter-relacoes com *Spodoptera frugiperda* (Lepidoptera: Noctuidae). 5. Fase reproductiva e longevidade. *Rev. Bras. Entomol.* 35: 521-530.
- MITCHELL, E. R., V. H. WADDILL, AND T. R. ASHLEY. 1984. Population dynamics of the fall armyworm (Lepidoptera: Noctuidae) and its larval parasites on the whorl stage corn in pheromone-permeated field environments. *Environ. Entomol.* 13: 1618-1623.
- MOLINA-OCHOA, J., R. LEZAMA-GUTIÉRREZ, J. J. HAMM, B. R. WISEMAN, AND M. LÓPEZ-EDWARDS. 1999. Integrated control of fall armyworm (Lepidoptera: Noctuidae) using resistant plants and entomopathogenic nematodes (Rhabditida: Steinernematidae). *Florida Entomol.* 82: 263-271.
- MOLINA-OCHOA, J., J. J. HAMM, R. LEZAMA-GUTIÉRREZ, M. LÓPEZ-EDWARDS, M. GONZÁLEZ-RAMÍREZ, AND A. PESCADOR-RUBIO. 2001. A survey of fall armyworm (Lepidoptera: Noctuidae) parasitoids in the Mexican states of Michoacan, Colima, Jalisco, and Tamaulipas. *Florida Entomol.* 84:31-36.
- MOLINARI, A. M., Y S. D. AVALOS. 1997. Contribución al conocimiento de taquinidos (Diptera) parasitoides de defoliadores (Lepidoptera) del cultivo de la soja. *Rev. Soc. Entomol. Argentina* 56: 131-136.
- MONTOYA-BURGOS, J. A. 1979. Estudio preliminar del ectoparásito del género *Euplectrus* del gusano cogollero *Spodoptera frugiperda*. VII Reunión Nacional de Control Biológico. Secretaría de Agricultura y Recursos Hidráulicos, Sanidad Vegetal. 24-26 de Abril de 1979, Veracruz, Veracruz, México, 4 pp.
- MONTOYA-BURGOS, J. A. 1980. Observaciones sobre la importancia de *Euplectrus* sp. en la integración de los métodos de control de plagas del maíz. VIII Reunión Nacional de Control Biológico. Secretaría de Agricultura y Recursos Hidráulicos, Sanidad Vegetal. 22-25 de Abril de 1980, Manzanillo, Colima, México, 7 pp.
- MORALES-PÉREZ, A. 1982. Cría masiva del parásito *Telenomus* sp. usando como huesped huevecillos de *Spodoptera exigua*, pp. 13-16. *En X Reunión Nacional de Control Biológico. Grupo Sectorial Agropecuario y Forestal. Secretaría de Agricultura y Recursos Hidráulicos. Abril 1982, Durango, Durango, México.*
- MOREY, C. S. 1971. Biología de *Campoletis grioti* (Blanchard) (Hymen.: Ichneumonidae) parásito de la "lagarta cogollera del maíz" *Spodoptera frugiperda* (J.E. Smith). Sociedad Entomológica del Perú, pp. 263-271. *En Anales del Primer Congreso Latinoamericano de Entomología, Cusco Perú, 12-18 de Abril de 1971.*
- MOYA, M. 1980. Parasitismo de Bracon Kirkpatricki Wilkinson (Hymenoptera-Braconidae) en larvas de *Spodoptera frugiperda* Smith. *Ibid.* 14 pp.
- MUESEBECK, C. F. W. 1921. A revision of the North American species of Ichneumon-flies belonging to the genus *Apanteles*. *U.S. Nat. Mus., Proc.* 58: 483-576.
- MUESEBECK, C. F. W., K. V. KROMBEIN, AND H. K. TOWNES (Eds.). 1951. Hymenoptera of America North of Mexico. Synoptic Catalog. USDA Agric. Monogr. No. 2 1420 pp.
- MULOCK, B. S., S. S. SWEZEY, C. NARVAEZ, P. CASTILLO, AND C. M. RIZO. 1990. Development of baculoviruses as a contribution to biological control of lepidopterous pests of basic grains in Nicaragua, pp. 179-183. *In Proc. Vth. Inter. Colloq. Invertebr. Pathol. Microb. Control. Adelaide, Australia, 20-24 August 1990.*
- MYERS, J. G. 1932. Biological observations on some neotropical parasitic hymenoptera. *Trans. Ent. Soc. Lond.* 80: 121-136.
- NAVA, R. A., Y N. S. CASTRO. 1991. Parasitismo del gusano cogollero *Spodoptera frugiperda* (J.E. Smith) en la región central del Estado de Tamaulipas., pp. 62-64 *En XIV Congreso Nacional de Control Biológico. Soc. Mex. Contr. Biol. Octubre 10-11, 1991. Buenavista, Saltillo, Coahuila, México.*
- NICKLE, D. A. 1976. The peanut agroecosystem in Central Florida. Ph.D. dissertation, University of Florida, Gainesville, 131 pp.
- NOTZ, P. A. 1972. Parasitismo de Diptera e Hymenoptera sobre larvas de *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) recolectadas en

- maíz, Maracay, Venezuela. Rev. Fac. Agron. Univ. Central. Venezuela 6: 5-16.
- ORTEGÓN, E. J., N. C. TORRES, E. LUQUE, Y A. SIABATTO. 1988. Estudio sobre la longevidad, hábitos, progenie y evaluación preliminar de *Meteorus laphygmae* (Viereck), parásito de *Spodoptera* spp. Rev. Colombiana Entomol. 14: 7-12.
- PAINTER, R. H. 1955. Insects on corn and teosinte in Guatemala. J. Econ. Ent. 48: 36-42.
- PAIR, S. D. 1995. Biology and rearing of *Diapetimorpha introita* (Cresson) (Hymenoptera: Ichneumonidae) on host and non-host noctuid pupae. J. Entomol. Sci. 30: 468-480.
- PAIR, S. D., AND H. R. GROSS, JR. 1984. Field mortality of pupae of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith), by predators and a newly discovered parasitoid, *Diapetimorpha introita*. J. Georgia Entomol. Soc. 19:22-26.
- PAIR, S. D., J. R. RAULSTON, A. N. SPARKS, AND P. B. MARTIN. 1986. Fall armyworm (Lepidoptera: Noctuidae) parasitoids: Differential spring distribution and incidence on corn and sorghum in the southern United States and Northeastern Mexico. Environ. Entomol. 15: 342-348.
- PAIR, S. D., AND H. R. GROSS, JR. 1989. Seasonal incidence of fall armyworm (Lepidoptera: Noctuidae) pupal parasitism in corn by *Diapetimorpha introita* and *Cryptus albitarsis* (Hymenoptera: Ichneumonidae). J. Entomol. Sci. 24: 339-343.
- PANTOJA, A., C. M. SMITH, AND J. F. ROBINSON. 1985. Natural control agents affecting *Spodoptera frugiperda* (Lepidoptera: Noctuidae) infesting rice in Puerto Rico. Florida Entomol. 68: 488-490.
- PANTOJA, A., AND J. R. FUXA. 1992. Prevalence of biotic control agents in the fall armyworm *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae). Folia Entomol. Mex. 84: 79-84.
- PARKER, H., P. BERRY, AND A. S. GUIDO. 1953. Host-parasite and parasite-host lists of insects reared in the South American parasite laboratory. Montevideo, Republica Oriental del Uruguay, 100 p.
- PASSOA, S. 1983. Lista de los insectos asociados con los granos básicos y otros cultivos selectos en Honduras. CEIBA 25. 97 pp.
- PATEL, P. N., E M. E. M. HABIB. 1982. Ocorrência natural de *Aspergillus parasiticus* em populações de *Spodoptera frugiperda* (Abbot & Smith, 1797) (Lepidoptera: Noctuidae) e sua transmissão por insetos parasitos. Rev. Agric. Piracicaba 57: 223-232.
- PATEL, P. N., E M. E. M. HABIB. 1984. Levantamento e eficácia de insetos parasitos de *Spodoptera frugiperda* (Abbot & Smith, 1797) (Lepidoptera: Noctuidae). Rev. Agric. Brasil 59: 229-237.
- PATEL, P. N., E M. E. M. HABIB. 1986. Levantamento e eficácia de insetos parasitos de *Spodoptera frugiperda* (Abbot & Smith, 1797) (Lepidoptera: Noctuidae). Rev. Agric. Piracicaba 61: 93-100.
- PAVUK, D. M., AND B. R. STINNER. 1991. New lepidoptera-parasitoid associations in weedy corn plantings: a potential alternate host for *Ostrinia nubilalis* (Lepidoptera: Pyralidae) parasitoids. Great Lakes Entomol. 24:219-223.
- PENA, G. 1980. Estudio de Bracon Kirkpatricki Wilkinson en diferentes huéspedes, con fines de control biológico. VIII Reunión Nacional de Control Biológico, Manzanillo, Col., Mex. 17 pp.
- PERAZA-LIZARRAGA, J. C. 1982. Dinámica de poblaciones de insectos benéficos en cultivo de maíz en Durango, Durango, pp. 91-95. En X Reunión Nacional de Control Biológico. Grupo Sectorial Agropecuario y Forestal. Secretaría de Agricultura y Recursos Hidráulicos. Abril 1982, Durango, Durango, México.
- PIERCE, W. D., AND T. E. HOLLOWAY. 1912. Notes on the biology of *Chelonus texanus* Cress. J. Econ. Entomol. 5: 425-428.
- PORTER, C. C. 1998. Guía de los géneros de Ichneumonidae en la región Neártica del Sur de Sudamérica. Opera Lilloana 42: 234 pp.
- PRATISSOLI, D., J. B. TORRES, Y J. C. ZANUNCIO. 1999. Selectividad de *Trichogramma demoraesi* (Hymenoptera: Trichogrammatidae) en huevos de tres lepidópteros inactivados. Agro-Ciencia 15: 75-80.
- RAJAPAKSE, R. H. S., T. R. ASHLEY, AND V. H. WADDILL. 1985. Biology and host acceptance of *Microplitis manilae* (Hymenoptera: Braconidae) raised on fall armyworm larvae *Spodoptera frugiperda*. Florida Entomol. 68: 653-657.
- RAVLIN, F. W., AND F. W. STEHR. 1984. Revision of the genus *Archytas* (Diptera: Tachinidae) for America North of Mexico. Misc. Pub. Entomol. Soc. Am. 58, 59 pp.
- REED, D. J. 1980. Toxicological and parasitological studies on the fall armyworm, *Spodoptera frugiperda* (J.E. Smith), in Alabama. M.S. Thesis, Auburn University, Auburn, Alabama, 55 pp.
- REMILLET, M., AND J. F. SILVAIN. 1988. *Noctuidonema guyanense* n.g., n. sp. (Nematoda: Aphelenchoididae) ectoparasite de noctuelles du genre *Spodoptera* (Lepidoptera: Noctuidae). Rev. Nematol. 11: 21-24.
- REZENDE, M. A. A., I. CRUZ, E T. M. C. DELLA LUCIA. 1994. Consumo foliar de milho e desenvolvimento de lagartas de *Spodoptera frugiperda* (Smith) parasitadas por *Chelonus insularis* (Cresson) (Hymenoptera: Braconidae). An. Soc. Entomol. Brasil 23: 473-478.
- REZENDE, M. A. A., I. CRUZ, E T. M. C. DELLA LUCIA. 1995a. Aspectos biológicos do parasitoide *Chelonus insularis* (Cresson) (Hymenoptera, Braconidae) criados em ovos de *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae). Rev. Bras. Zool. 12: 779-784.
- REZENDE, M. A. A., T. M. C. DELLA LUCIA, E I. CRUZ. 1995b. Comportamento de lagartas de *Spodoptera frugiperda* (Lepidoptera: Noctuidae) parasitadas por *Chelonus insularis* (Hymenoptera, Braconidae), sobre plantas de milho. Rev. Bras. Entomol. 39: 675-681.
- RIGGIN, T. M., B. R. WISEMAN, D. J. ISENHOUR, AND K. E. ESPELIE. 1992. Incidence of fall armyworm (Lepidoptera: Noctuidae) parasitoids on resistant and susceptible corn genotypes. Environ. Entomol. 21: 888-895.
- RIGGIN, T. M., K. E. ESPELIE, B. R. WISEMAN, AND D. J. ISENHOUR. 1993. Distribution of fall armyworm (Lepidoptera: Noctuidae) parasitoids on five corn genotypes in South Georgia. Florida Entomol. 76: 292-302.
- RIGGIN, T. M., B. R. WISEMAN, D. J. ISENHOUR, AND K. E. ESPELIE. 1994. Functional response of *Cotesia marginiventris* (Cresson) (Hym., Braconidae) to *Spodoptera frugiperda* (J.E. Smith) (Lep., Noctuidae) on meridic diet containing resistant or susceptible corn genotypes. J. Appl. Ent. 117: 144-150.
- RODRÍGUEZ-LUNA, S. J. 1982. Parasitismo natural de *Trichogramma* sp. sobre huevecillos de *Spodoptera frugiperda* (Smith) en maíz de temporal de Jiutepec, Emiliano Zapata y Xotitepec, Mor., pp. 22-33. En X Reunión Nacional de Control Biológico. Secretaría de Agricultura y Recursos Hidráulicos, Abril 1982, Durango, Durango, México.

- ROGERS, C. E., O. G. MARTI, A. M. SIMMONS, AND J. F. SILVAIN. 1990a. Host range of *Noctuidonema guyanense* (Nematoda: Aphelenchoididae): An ectoparasite of moths in French Guiana. *Environ. Entomol.* 19: 795-798.
- ROGERS, C. E., A. M. SIMMONS, AND O. G. MARTI. 1990b. Parasitism of Lepidoptera adults by *Noctuidonema guyanense* Remillet and Silvain (Nematoda: Aphelenchoididae) in Southeastern United States. *J. Agric. Entomol.* 7: 241-245.
- ROGERS, C. E., A. M. SIMMONS, AND O. G. MARTI. 1991. *Noctuidonema guyanense*: an ectoparasitic nematode of fall armyworm adults in the tropical Americas. *Florida Entomol.* 74: 246-257.
- ROGERS, C. E., AND O. G. MARTI, JR. 1992a. *Noctuidonema guyanense* (Nematoda: Aphelenchoididae): population profiles on male and female fall armyworm moths. *J. Entomol. Sci.* 27: 354-360.
- ROGERS, C. E., AND O. G. MARTI, JR. 1992b. Infestation and dispersal of *Noctuidonema guyanense* (Nematoda: Aphelenchoididae) on *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Environ. Entomol.* 21: 417-421.
- ROGERS, C. E., AND O. G. MARTI, JR. 1993a. Infestation dynamics and distribution of *Noctuidonema guyanense* (Nematoda: Aphelenchoididae) on adults of *Spodoptera frugiperda* and *Mocis latipes* (Lepidoptera: Noctuidae). *Florida Entomol.* 76: 326-333.
- ROGERS, C. E., AND O. G. MARTI, JR. 1993b. Maintenance of an ectoparasitic nematode, *Noctuidonema guyanense* (Nematoda: Aphelenchoididae), on *Spodoptera frugiperda* (Lepidoptera: Noctuidae) moths. *J. Entomol. Sci.* 28: 335-337.
- ROGERS, C. E., O. G. MARTI, JR., AND A. M. SIMMONS. 1993. *Noctuidonema guyanense* (Nematoda: Aphelenchoididae): Host range and pathogenicity to the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), pp. 27-32. In *Nematodes and the biological control of insect pests*. Bedding, R., R. Akhurst & H. Kaya [eds.] CSIRO, Australia.
- ROGERS, C. E., AND O. G. MARTI, JR. 1994. Population structure and transfer success of *Noctuidonema guyanense* (Nematoda: Aphelenchoididae) on moths of *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.* 87: 327-330.
- ROGERS, C. E., AND O. G. MARTI, JR. 1996. Beet armyworm (*Spodoptera exigua*) as a host for the ectoparasitic nematode, *Noctuidonema guyanense*. *J. Agric. Entomol.* 13: 81-86.
- ROGERS, C. E., O. G. MARTI, JR., L. D. CHANDLER, AND A. M. SIMMONS. 1996. Bionomics of the fall armyworm (Lepidoptera: Noctuidae) and its ectoparasitic nematode, *Noctuidonema guyanense*, in South Georgia. *J. Entomol. Sci.* 31: 209-217.
- ROGERS, C. E., O. G. MARTI, JR., AND J. A. CLAYTON. 1997. Report of the ectoparasitic nematode *Noctuidonema guyanense* (Acugetturidae) infesting Lepidoptera in the Fiji Islands. *Nematologica* 43: 505-506.
- ROHLFS, W. M., AND T. P. MACK. 1985. Seasonal parasitism rates, host size, and adult emergence pattern of parasitoids of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith), with emphasis on *Ophion flavidus* Brulle (Hymenoptera: Ichneumonidae). *Ann. Entomol. Soc. Am.* 78: 217-220.
- RYDER, W. D., AND N. PIEDRA. 1968. Field investigations into the control for some insects pests of maize in Cuba with high-volume sprays of DDT, diazinon, malathion, and phosdrin. *Ibid* 2: 295-303.
- RYDER, W. D., AND N. PULGAR. 1969. A note on parasitism of the fall armyworm, *Spodoptera frugiperda*, on maize. *Rev. cubana Cienc. Agric. (English ed.)* 3: 267-271.
- SABROSKY, C. W. 1981. A partial revision of the genus *Eucelatoria* (Diptera: Tachinidae), including important parasites of *Heliothis*. *Tech. Bull. Sci. & Educ. Adm. USDA*, No. 1635, 18 pp.
- SÁNCHEZ-GARCÍA, J. A., J. ROMERO-NAPOLES, S. RAMARIZ-ALARCÓN, S. ANAYA-ROSALES, Y J. L. CARILLO-SÁNCHEZ. 1998. Géneros de Braconidae del estado de Guanajuato (Insecta: Hymenoptera). *Acta Zool. Mex.* 74: 59-137.
- SARMIENTO, M. J., Y R. V. RAZURI. 1978. *Bacillus thuringiensis* en el control de *Spodoptera frugiperda* y de *Diatraea saccharalis* en maíz. *Rev. Peruana Entomol.* 21: 121-124.
- SEGEREN, P. A., AND S. R. SHARMA. 1979. Insect control on maize in Suriname. *Proc. Caribbean Food Crops Soc.* 15: 142-155.
- SILVA, F. M. A., H. G. FOWLER, E R. N. S. LEMOS. 1997. Parasitismo em lagarta-do-cartucho, *Spodoptera frugiperda* (Smith), na região do Triângulo Mineiro, MG. *An. Soc. Entomol. Brasil* 26: 235-241.
- SILVAIN, J. F., AND M. REMILLET. 1993. Ecology and biology of *Noctuidonema guyanense* (Nematoda: Aphelenchoididae), an ectoparasite of *Spodoptera frugiperda* (Lep., Noctuidae), in French Guiana. *Entomophaga* 38: 465-474.
- SILVEIRA, G. A., Y A. RUFFINELLI. 1956. Primer catálogo de los parásitos y predadores encontrados en el Uruguay. *Bol. Fac. Agron. Montev.* 32: 80 pp.
- SILVEIRA, J. C. F. DA, E. T. SASAKI, M. A. FORNER, M. S. HONDA, E M. H. CALAFIORI. 1987. Ocorrência de parasitoides de *Spodoptera frugiperda* (J.E. Smith, 1797) em cultura de milho, em Espírito Santo do Pinhal, SP. *Ecossistema* 12: 41-44.
- SIMMONS, A. M., AND C. E. ROGERS. 1990a. Temperature and humidity effects on *Noctuidonema* (Nematoda: Aphelenchoididae), an ectoparasite of adult *Spodoptera frugiperda* (Lepidoptera: Noctuidae), and transfer success during host mating. *Ann. Entomol. Soc. Am.* 83: 1084-1087.
- SIMMONS, A. M., AND C. E. ROGERS. 1990b. Distribution and prevalence of an ectoparasitic nematode, *Noctuidonema guyanense*, on moths of the fall armyworm (Lepidoptera: Noctuidae) in the tropical Americas. *J. Entomol. Sci.* 25: 510-518.
- SIMMONS, A. M., AND C. E. ROGERS. 1991. Dispersal and seasonal occurrence of *Noctuidonema guyanense*, an ectoparasitic nematode of adult fall armyworm (Lepidoptera: Noctuidae), in the United States. *J. Entomol. Sci.* 26: 136-148.
- SIMMONS, A. M., C. E. ROGERS, K. U. BUCKMIRE, B. GRAY, K. D. MONKMAN, A. PANTOJA, J. R. RAULSTON, AND V. H. WADDILL. 1991. Seasonal chronology of *Noctuidonema*, an ectoparasitic nematode of adult moths, in tropical and subtropical America. *Florida Entomol.* 74: 311-319.
- SIMMONS, A. M., AND O. G. MARTI, JR. 1992. Mating by the fall armyworm (Lepidoptera: Noctuidae): frequency, duration, and effect of temperature. *Environ. Entomol.* 21: 371-375.
- SIMMONS, A. M., AND C. E. ROGERS. 1994. Effect of an ectoparasitic nematode, *Noctuidonema guyanense*, on adult longevity and egg fertility in *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Biological Control* 4: 285-289.

- SIMMONS, A. M., AND C. E. ROGERS. 1996. Ectoparasitic acugutturid nematodes of adult Lepidoptera. *J. Nematol.* 28: 1-7.
- SINHA, A. K. 1982. Integrated pest control programme in intermediate savannahs of Guyana. *Caribbean Plant Protec. Newsl.* 1: 2.
- SIVASUBRAMANIAM, W., S. D. WRATTEN, AND J. KLIMASZEWSKI. 1997. Species composition, abundance, and activity of predatory arthropods in carrot fields, Canterbury, New Zealand. *New Zealand J. Zool.* 24: 205-212.
- SOTERES, K. M., R. C. BERBERET, AND R. W. MCNEW. 1984. Parasitic insects associated with lepidopterous herbivores on alfalfa in Oklahoma. *Environ. Entomol.* 13: 787-793.
- STARY, P., AND K. S. PIKE. 1999. Uses of beneficial insect diversity in agroecosystem management. Chapter 4, pp. 49-67 *In* Biodiversity in Agroecosystems. W. W. Collins & C. O. Qualset [eds.]. CRC Press, Boca Raton, Florida, USA.
- TERÁN, B. J. 1974. Lista preliminar de dípteros parasíticos de otros insectos en Venezuela. *Rev. Fac. Agron. Alcance (Maracay)* 23: 64-65.
- TERÁN, B. J. 1977. Dípteros parasíticos de otros insectos en Venezuela. *Rev. Fac. Agron. (Maracay)* 9: 134-135.
- TERÁN, B. J. 1980. Lista preliminar de Hymenoptera parásitos de otros insectos en Venezuela. *Rev. Fac. Agron.* 11: 283-389.
- TINGLE, F. C., E. R. MITCHELL, AND J. R. MCLAUGHLIN. 1994. Lepidopterous pest of cotton and their parasitoids in a double-cropping environment. *Florida Entomol.* 77: 334-341.
- VALENCIA, L., AND R. VALDIVIA. 1973. Noctuides del Valle de Ica, sus plantas hospederas y enemigos naturales. *Rev. Peru. Entomol.* 16: 94-101.
- VALICENTE, F. H. 1989. Levantamento dos inimigos naturais de *Spodoptera frugiperda* (J.E. Smith, 1797) (Lepidoptera: Noctuidae) em diferentes regiões de Minas Gerais. *An. Soc. Entomol. Brasil* 18: 119-130.
- VALICENTE, F. H., E M. R. BARRETO. 1999. Levantamento dos inimigos naturais da lagarta do cartucho do milho, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae), na região de Cascavel, PR. *An. Soc. Entomol. Brasil* 28: 333-337.
- VAN DINE, D. L. 1913. The insects affecting sugar cane in Porto Rico. *J. Econ. Entomol.* 6: 251-257.
- VAN DITHER, J. B. M. 1960. Insect pests of cultivated plants in Surinam. *Bull. Landouwproefstation. Suriname* 76. 131 pp.
- VARGAS, M. L., Y G. G. SÁNCHEZ. 1983. Control natural de algunas plagas de arroz en las variedades IR-22 y CICA-6. *Rev. Colombiana Entomol.* 9: 50-54.
- VERA, M. L., L. VALVERDE, S. B. POPICH, Y Z. D. AJMAT-DE TOLEDO. 1995. Evaluación preliminar de los enemigos naturales de *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) en Tucuman, Argentina. *Acta Entomol. Chilena* 19: 135-141.
- VICKERY, R. A. 1929. Studies on the fall armyworm in the Gulf Coast District of Texas. *USDA Tech. Bull.* 138, 63 pp.
- VIRLA, E. G., M. V. COLOMO, C. BERTA, Y L. VALVERDE. 1999. El complejo de los parasitoides del "gusano cogollero" del maíz, *Spodoptera frugiperda*, en la República Argentina (Insecta: Lepidoptera). *Neotrópica* 45: 3-12.
- WAAGE, J. K. 1991. Biodiversity as a resource for biological control. Chapter XIII, pp. 149-163. *In* D. L. Hawksworth [ed.]. *The Biodiversity of Microorganisms and Invertebrates: Its Role in Sustainable Agriculture*. CAB International, Wallingford, UK.
- WADDILL, V. I. 1977. Shadow sampling: A fast, painless method for collecting fall armyworm egg masses. *Florida Entomol.* 60: 215-216.
- WADDILL, V. H., AND W. H. WHITCOMB. 1982. Release of *Telenomus remus* (Hym. Scelionidae) against *Spodoptera frugiperda* (Lep.:Noctuidae) in Florida, USA. *Entomophaga* 27: 159-162.
- WALL, R., AND R. C. BERBERET. 1974. The life cycle of *Euplectrus platyhypenae* a gregarious external parasitoid of peanut foliage feeders in Oklahoma. *Environ. Entomol.* 3: 744-746.
- WALL, R., AND R. C. BERBERET. 1975. Parasitoids associated with lepidopterous pests on peanuts; Oklahoma fauna. *Environ. Entomol.* 4: 877-882.
- WALTON, W. R. 1913. Efficiency of a tachinid parasite on the last instar of *Laphygma*. *Proc. Ent. Soc. Wash.* 15:128-131.
- WHEELER, G. S., T. R. ASHLEY, AND K. L. ANDREWS. 1989. Larval parasitoids and pathogens of the fall armyworm in Honduran maize. *Entomophaga* 34: 331-340.
- WILSON, A. G. L. 1983. Abundance and mortality of overwintering *Heliothis* spp. *J. Aust. Entomol. Soc.* 22: 191-199.
- WILSON, C. E. 1923. Insect pests of cotton in St. Croix and a means of combating them. *Virgin Islands Agric. Exp. Stn. Bull.* 3: 1-20.
- WILSON, F., AND C. B. HUFFAKER. 1976. The philosophy, scope, and importance of biological control, pp. 3-15. *In* C. B. Huffaker and P. S. Messenger [eds.] *Theory and Practice of Biological Control*. Academic Press, New York, USA.
- WOJCIK, B., W. H. WHITCOMB, AND D. H. HABECK. 1976. Host range testing of *Telenomus remus* (Hymenoptera: Scelionidae). *Florida Entomol.* 59: 195-198.
- WONG-ARÉVALO, B. 1982. Un nuevo sistema de liberación del parásito *Trichogramma* spp en el Siconusco Chiapas, México, pp. 34-46 *En* X Reunión Nacional de Control Biológico. Secretaría de Agricultura y Recursos Hidráulicos, Abril 1982, Durango, Durango, México.
- YASEEN, M. 1979. Introduction of exotic parasites for control of *Spodoptera* and *Heliothis* in Trinidad. Symp. on maize and peanut. Paramaribo, Suriname, Nov. 13-18, 1978. *Proc. Caribbean Food Crops Soc.* 15: 136-141.
- YASEEN, M., F. D. BENNETT, AND R. M. BARROW. 1981. Introduction of exotic parasites for control of *Spodoptera frugiperda* in Trinidad, the eastern Caribbean and Latin America, pp. 161-171. *In* C. W. D. Brathwaite and G. V. Pollard [eds.] *Urgent plant pest and disease problems in the Caribbean*. IICA Misc. Publ. 378.
- ZUCCHI, R. A., J. R. P. PARRA, AND S. SILVEIRA NETO. 1991. *Trichogramma* species associated with some lepidopterous pests in Brazil. *Coll. De-LINRA* 56: 131-134.