

Brachymeria spp. (Hymenoptera: Chalcididae) Parasitizing Pupae of Hersperidae and Nymphalidae (Lepidoptera) Pests of Oil Palm in the Brazilian Amazonian Region

Authors: Tinôco, Ricardo S., Ribeiro, Rafael C., Tavares, Marcelo T., Vilela, Evaldo F., Lemos, Walkymário De P., et al.

Source: Florida Entomologist, 95(3) : 788-789

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.095.0335>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

BRACHYMERIA spp. (HYMENOPTERA: CHALCIDIDAE) PARASITIZING PUPAE OF HERSPERIDAE AND NYMPHALIDAE (LEPIDOPTERA) PESTS OF OIL PALM IN THE BRAZILIAN AMAZONIAN REGION

RICARDO S. TINOCO¹, RAFAEL C. RIBEIRO², MARCELO T. TAVARES³, EVALDO F. VILELA⁴, WALKYMÁRIO DE P. LEMOS⁵ AND JOSÉ C. ZANUNCIO^{4*}

¹Grupo Agropalma S/A. Rodovia PA-150, km 74, Tailândia, Pará State, Brazil

²Departamento de Fitotecnia. Universidade Federal de Viçosa. Viçosa, Minas Gerais State, Brazil

³Departamento de Ciências Biológicas, Universidade Federal do Espírito Santo, Avenida Marechal Campos, Vitória, Espírito Santo State, Brazil

⁴Departamento de Entomologia, Universidade Federal de Viçosa. Viçosa, Minas Gerais State, Brazil

⁵Laboratório de Entomologia, Embrapa Amazônia Oriental, Tv. Dr. Enéas Pinheiro, sn, Marco, Belém, Pará State, Brazil

*Corresponding author; E-mail:zanuncio@ufv.br

The African oil palm (*Elaeis guineensis* Jacq.; Arecales: Arecaceae) is one of the most important oil producing plants yielding 4 to 6 tonnes of oil/ha/yr in Brazil and 8 to 10 tonnes of oil/ha/yr in Asia (Camillo et al. 2009). Plantations of this crop in Pará State, Brazil can reach up to 6 tonnes of oil/ha/yr with appropriate technologies; and Brazil can be a leading world producer of renewable fuels especially in scenarios of rising petroleum oil prices and concern over environmental pollution (Abdalla et al. 2008; Brokamp et al. 2011). This palm is well adapted to the ecological conditions of the Amazonian region, which has the largest area available to expand this crop in the world (Chia et al. 2009).

Lepidopteran pests can damage African oil palm plants in South America; control of such pests with chemical pesticides, can cause environmental and socioeconomic problems (Zeddam et al. 2003). The northern region of Brazil presents great diversity of lepidopteran defoliators of oil palm including *Opsiphanes invirae* Hübner and *Brassolis sophorae* L. (Nymphalidae); *Sibine* spp. and *Talima* sp. (Limacodidae); *Euprosterna* sp., *Automeris* spp. (Saturniidae) and *Saliana* sp. (Hesperiidae); and these pests have high damage capacities (Ribeiro et al. 2010).

The aim of this study was to evaluate the occurrence of parasitoids and hyperparasitoids obtained from pupae of lepidopteran defoliators of the oil palm cultivated in Pará State, Brazil.

This work was conducted in oil palm plantations of the Agropalma Complex in the Municipality of Thailândia, Southeast of Para State, Brazil. The total area of this company is 112,551 hectares with 39,700 hectares planted with oil palm. The geographical coordinates of the land of this company are S 24° 2'04" W 48° 08'02". The main cultivars of the oil palm grown in Brazil include 'Avros', 'Lamé', 'Ghana', 'Ekona', 'Embrapa-Lamé' and 'Kigoma'. Pupae of lepidopteran defoliators

were first collected from Aug to Oct 2007 (dry season) and subsequently in Jan and Feb 2008 (rainy season).

One hundred *Saliana* sp. pupae and 2,261 *O. invirae* pupae were collected in the field and held individually in the laboratory at 27 ± 1 °C, 75 ± 10% RH and 12:12 h L:D until emergence of adults of the parasitoids or the Lepidoptera. Adult parasitoids were mounted and/or maintained in 70% ethanol, sent for identification and deposited in the entomology collection of the EMBRAPA Eastern Amazon, Belém, Pará State, Brazil.

An average of 3 specimens of *Brachymeria pandora* was obtained from the each of the parasitized immatures of *Saliana* sp. and 1 individual each of the endoparasitoids *Brachymeria annulipes* and *Brachymeria koehleri* was obtained from each correspondingly parasitized *O. invirae* immature and from each parasitized pupa of the parasitoid *Chetogena scutellaris* (Wulp) (Diptera: Tachinidae). The genus *Brachymeria* Westwood (Chalcididae) has 45 species described in the Neotropical region (De Santis 1989; Tavares et al. 2006). *Brachymeria koehleri* Blanchard, 1935 has been registered in Venezuela, Brazil (Espírito Santo and Rio de Janeiro States) and Argentina; *Brachymeria pandora* (Crawford 1914) has been reported in Espírito Santo, Goiás and Rio de Janeiro States, Brazil and in Venezuela and Guyana; and *Brachymeria annulipes* (Costa Lima 1919) has been registered in Maranhão and Espírito Santo States, Brazil (Marchiori et al. 2003; Tavares & Araújo 2007).

Brachymeria koehleri was recorded as a hyperparasitoid from pupae of the parasitoid *Chetogena scutellaris* (Diptera: Tachinidae) that had been reared in pupae of *O. invirae* collected on oil palm plants. This species has also been reported as a hyperparasitoid of Tachinidae and Sarcophagidae (Diptera) from lepidopteran pupae, but it emerged

from *Alabama argillacea* (Hübner) and *Mocis latipes* (Guenée) (Noctuidae) pupae (Terán 1980). *Brachymeria annulipes* parasitized *Pectinophora gossypiella* (Saunders) (Gelechidae) (Noyes 2002), and the parasitoid *B. pandora* emerged from pupa of *Calpodesthlius* (Stoll), *Argon lota* Hewitson (Hesperiidae) and *Historis odius* (Fabricius) (Nymphalidae) (Marchiori et al. 2003; Gil-Santana & Tavares 2005; Salgado-Neto et al. 2010).

This is the first report of a species of the genus *Brachymeria* parasitizing pupae of *O. invirae* and *Saliana* sp., and as an hyperparasitoid of *C. scutellaris* in palm plantations in the Brazilian Amazonian region.

SUMMARY

The cultivation of the African oil palm is one of the main agricultural activity in humid areas of the world, such as Amazonia, but defoliating caterpillars can reduce the productivity of this crop in northern Brazil. This is the first report in Brazil of occurrence of the parasitoids *Brachymeria annulipes* (Costa Lima 1919) from pupae of *Opsiphantes invirae* (Hübner 1808) (Nymphalidae), *Brachymeria pandora* (Crawford 1919) (Chalcidiidae) from those of *Saliana* sp. (Hesperiidae) and *Brachymeria koehleri* Blanchard, 1935 as an hyperparasitoid of pupae of the natural enemy *Chetogena scutellaris* (Tachinidae) from *O. invirae* on oil palm cultivated in Pará State, Brazil. This is the first report of species of the genus *Brachymeria* parasitizing pupae of *O. invirae*, and *Saliana* sp., and as an hyperparasitoid of *C. scutellaris* in palm plantations in the Brazilian Amazonian region.

Key Words: *Elaeis guineensis*, natural enemies, new records.

RESUMO

O cultivo da palma de óleo é a principal atividade agrícola em áreas úmidas do mundo, como a Amazônia, mas lagartas desfolhadoras podem reduzir a produtividade desta cultura no Norte do Brasil. Este é o primeiro relato no Brasil da ocorrência dos parasitoides *Brachymeria annulipes* (Costa Lima 1919) de pupas de *Opsiphantes invirae* (Hübner 1808) (Lepidoptera: Nymphalidae) e *Brachymeria pandora* (Crawford 1919) (Hymenoptera: Chalcidiidae) de pupas de *Saliana* sp. (Lepidoptera: Hesperiidae) e de *Brachymeria koehleri* Blanchard, 1935 como hiperparasitoide de pupas do inimigo natural *Chetogena scutellaris* (Wulp) (Diptera: Tachinidae) de *O. invirae* em cultivos da palma do óleo no Estado do Pará, Brasil. Este é, também, o primeiro relato de espécies do gênero *Brachymeria* parasitando pupas de *O. invirae*, *Saliana* sp. e de *C. scutellaris* como hiperparasitoide em plantações de palmeiras na região Amazônica Brasileira.

Palavras-chave: *Elaeis guineensis*, inimigos naturais, novos registros.

ACKOWLEGMENTS

To “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)”, “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)” and “Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)” for financial support.

REFERENCES CITED

- ABDALLA, A. L., SILVA FILHO J. C., GODÓ, A. R., CARMO C. A., AND EDUARDO, J. L. P. 2008. Utilização de subprodutos da indústria de biodiesel na alimentação de rumiantes. Rev. Brasileira Zootec. 37: 260-258.
- BROKAMP, G., VALDERRAMA, N., MITTELBACH, M., GRANDEZ, C. A. R., BARFOD, A. S., AND WEIGEND, M. 2011. Trade in palm products in North-Western South America. Bot. Rev. 77: 571-606.
- CAMILLO, J., LUIS, Z. G., AND SCHERWINSKI-PEREIRA, J. E. 2009. Tolerância de sementes de dendzeiro à criopreservação. Pesqui. Agropecu. Brasileira 44: 11-15.
- CHIA, G. S., LOPEZ, R., CUNHA, R. N. V., ROCHA R. N. C., AND LOPEZ M. T. G. 2009. Repetibilidade da produção de cachos de híbridos interespécíficos entre o caiáue e o dendzeiro. Acta Amaz. 39: 249-254.
- DE SANTIS, L. 1989. Catalogo de los Himenopteros Calcidoideos (Hymenoptera) al sur de los Estados Unidos, segundo suplemento. Acta Entomol. Chil. 15: 9-90.
- GIL-SANTANA, H. R., AND TAVARES, M. T. 2005. *Brachymeria pandora* (Crawford) (Hymenoptera, Chalcidiidae): a new parasitoid of *Historis odius* (Fabricius) (Lepidoptera, Nymphalidae). Rev. Bras. Zool. 22: 1211-1212.
- MARCHIORI, C. H., PENTEADO-DIAS, A. M., AND TAVARES, M. T. 2003. Parasitoids of the family Chalcidiidae collected in pastures and forests using yellow traps, in Itumbiara, Goiás, Brasil. Braz. J. Biol. 63: 357-360.
- NOYES, J. S. 2002. Interactive catalogue of world Chalcidoidea. Taxapad, Vancouver. (Banco de Dados em CD-ROM).
- RIBEIRO, R. C., LEMOS, W. P., BERNARDINO, A. S., BUECKE, J., AND MÜLLER, A. A. 2010. Primeira ocorrência de *Alcaeorrhynchus grandis* (Dallas) (Hemiptera: Pentatomidae) predando lagartas desfolhadoras do dendzeiro no Estado do Pará. Neotrop. Entomol. 39: 131-132.
- SALGADO-NETO, G., MARÉ R. D., AND LOPEZ-DA-SILVA, M. 2010. Parasitismo de pupas de *Argon lota* Hewitson (Lepidoptera: Hesperiidae) por *Brachymeria pandora* (Crawford) (Hymenoptera: Chalcidiidae) no Rio Grande do Sul. Neotrop. Entomol. 39: 311-312.
- TAVARES, M. T., AND ARAUJO, B. C. 2007. Espécies de Chalcidiidae (Hymenoptera, Insecta) do Estado do Espírito Santo, Brasil. Biota Neotrop. 7: 1-8.
- TAVARES, M. T., NAVARRO-TAVARES, A. B., AND ALMEIDA, G. D. S. 2006. The species of Chalcidiidae (Hymenoptera) parasitoids of *Parides ascanius* (Cramer), an endangered Papilionidae (Lepidoptera) from restingas of Southeastern Brazil. Zootaxa 1197: 55-63.
- TERÁN, B. J. 1980. Lista preliminar de Hymenoptera parasitos de otros insectos em Venezuela. Rev. Fac. Agron. (LUZ) 11: 283-389.
- ZEDDAM, J., CRUZADO, J. A., RODRIGUEZ J. L., AND RAVALLEC M. 2003. A new nucleopolyhedrovirus from the oil-palm leaf-eater *Euproctis elaeasa* (Lepidoptera: Limacodidae): preliminary characterization and field assessment in Peruvian plantation. Agr. Exosyst. Environ. 96: 69-75.