



New Attractant Food for Catching Adult Rednecked Peanutworm (Lepidoptera: Gelechiidae) in Peanut

Authors: Rivero, Yoandry Rodríguez, Andrade, Daniel Junior de, Santos, Fabiano Aparecido dos, Melville, Cirano Cruz, and Leite, Gustavo Wallace Pacheco

Source: Florida Entomologist, 100(3) : 660-662

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0326>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

New attractant food for catching adult rednecked peanutworm (Lepidoptera: Gelechiidae) in peanut

Yoandry Rodríguez Rivero^{1,*}, Daniel Junior de Andrade¹, Fabiano Aparecido dos Santos¹, Cirano Cruz Melville¹, and Gustavo Wallace Pacheco Leite¹

Production of peanuts, *Arachis hypogaea* L. (Fabaceae), is a traditional agricultural activity in several regions of Brazil, especially in the state of São Paulo, which accounts for 80% of the national production (Santos & Todeschini 2012). The rednecked peanutworm, *Stegasta bosqueella* Chambers (Lepidoptera: Gelechiidae), is currently considered the most important lepidopteran pest of peanut crops due to the extent of damage and its occurrence in several regions (Boiça Junior et al. 2012). *Stegasta bosqueella* caterpillars attack shoots when the leaflets are still closed, reducing the leaf area and affecting development of the plant (Almeida 2013).

To control *S. bosqueella*, pyrethroid and carbamate insecticides are used, which contribute to environmental contamination (Almeida 2013; Scarpellini et al. 2014). As an alternative to chemical control, ethological control is recommended, which is a method based on studies of the physiology and behavior of insects for pest control (Altamar et al. 2016). Ethological control includes techniques such as trapping using attractants and baits, repellents, feeding inhibitors, and other products (Cisneros 1995).

Sugarcane molasses-baited traps are used to monitor several pests, mainly lepidopterans such as *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae), *Helicoverpa* species (Lepidoptera: Noctuidae), *Chrysodeixis includens* Walker (Lepidoptera: Noctuidae), *Manduca sexta* L. (Lepidoptera: Sphingidae), and *Keiferia lycopersicella* Walshingham (Lepidoptera: Gelechiidae) (Garza 2002).

Cisneros (2010) confirmed that several noctuid species and some other lepidopterans are attracted by the odor of sugarcane molasses fermentation. Jiménez (2010) also obtained satisfactory results in the capture of *Heliothis virescens* (F.) (Lepidoptera: Noctuidae) and *Mocis* sp. (Hübner) (Lepidoptera: Noctuidae) with attractant traps in areas of tobacco seed production in Cuba. In Brazil, traps with sugarcane molasses are used to capture the important fruit fly pests *Anastrepha fraterculus* (Wiedemann) (Diptera: Tephritidae) and *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae) (Raga et al. 2006).

To reduce risks and impacts on human health and the environment caused by the use of pesticides in peanut crops, alternative methods for monitoring and control of the rednecked peanutworm should be explored. Therefore, we sought to evaluate the potential of sugarcane molasses as an attractant food for the capture of *S. bosqueella* adults in peanut crops.

Evaluations were carried out in an experiment established to compare 2 peanut cultivation systems: intercropped with sorghum, *Sorghum bicolor* (L.) Moench (Poaceae), and sunflower, *Helianthus annuus* L. (Asteraceae), and the conventional production system (peanut only). The experiment was conducted in an agricultural area located at the School of Agricultural and Veterinary Sciences, Jaboticabal, São Paulo, Brazil, from Nov 2016 to Feb 2017 (average temperature of 24.2 °C, 75.4% RH, and total annual precipitation of 216.2 mm) and with a record of this insect being present in the area. The intercropping system occupied an area of 0.32 ha of peanut, intercropped with 0.16 ha of sorghum and 0.12 ha of sunflower, whereas the conventional system occupied an area of 0.49 ha. The latter was located 300 m from the intercropping system. The crops were planted on 28 Nov 2016 and the traps installed on 17 Jan 2017.

A total of 10 traps were installed, 5 in each system, using 2 L plastic bottles containing 100 mL of sugarcane molasses per bottle. The bottles had 2 “doors” or “flaps” cut into the sides, allowing entry of the moths. Traps were supported by wood rods at a height of approximately 20 cm from ground level, and sugarcane molasses was replaced twice per mo.

The traps were checked weekly and moths were removed and returned to the laboratory for identification. Voucher specimens were sent to EMBRAPA Cerrados in Brasília, Federal District, Brazil, for confirmation of *S. bosqueella* and other lepidopterans.

Moths of *S. bosqueella* (Fig. 1) were captured in traps containing sugarcane molasses in both cropping systems. In total, 310 *S. bosqueella* adults were captured, representing 60% of total moth capture in the conventional system and 41% of the moths in the intercropped system (Table 1). On average, 7.4 and 1.7 *S. bosqueella* adults per evaluation period were trapped in the conventional and intercropping systems, respectively.

Adult lepidopterans feed on pollen, nectar, and fermenting fruit juices. Females of several species orient to the host plant by olfactory stimuli and phototaxis, or by the combination of both (Bosa et al. 2011). Lorya et al. (2002) reported that adult *Opsiphanes cassina* Felder (Lepidoptera: Nymphalidae) females visited traps containing sugarcane molasses looking for sugars and nitrogen compounds that are important for the maturation of their eggs, and Parra et al. (2009) found that sugarcane molasses captured *O. cassina* adults.

¹UNESP - São Paulo State University, School of Agricultural and Veterinary Sciences (FCAV/UNESP). Via de Acesso Prof. Paulo Donato Castellane, s/nº, 14884-900 Jaboticabal, São Paulo State, Brazil; E-mail: yoandryrodriguez52@gmail.com (Y. R. R.), daniel.andrade@fcav.unesp.br (D. J. A.), fabiano.unesp@hotmail.com (F. A. S.), ciranomelville@outlook.com (C. C. M.), gustavowalacepl@outlook (G. W. P. L.)

*Corresponding author; E-mail: yoandryrodriguez52@gmail.com (Y. R. R.)



Fig. 1. Trap containing sugarcane molasses and *Stegasta bosqueella* adults (see white circles) captured in a peanut crop.

Particularly for the state of São Paulo, the use of traps baited with sugarcane molasses is an economically viable alternative due to the ready availability of the product coming from the sugarcane industry.

However, improvements in this technique are needed, including improved formulations, sugarcane molasses preservatives, and optimal distribution of traps in the crop.

Table 1. Weekly captures of *Stegasta bosqueella* and other lepidopterans captured by traps in 2 peanut cultivation systems.

		Number of lepidopteran adults per evaluation period											
		1		2		3		4		5		6	
		Sb	L	Sb	L	Sb	L	Sb	L	Sb	L	Sb	L
Intercropping system	1	0	0	3	1	0	3	0	2	0	0	0	5
	2	6	2	11	0	2	1	0	5	0	2	0	5
	3	2	2	6	4	3	0	0	1	0	0	0	3
	4	2	3	7	7	0	1	1	1	0	1	0	3
	5	0	0	0	0	0	3	0	2	0	0	0	3
Conventional system	1	3	9	10	1	18	2	9	4	4	9	5	7
	2	5	6	7	1	10	2	14	3	6	6	10	12
	3	12	7	7	6	6	5	9	7	11	8	6	12
	4	9	2	22	10	3	8	5	4	6	3	4	6
	5	4	4	14	5	12	4	21	2	7	13	8	9

Sb: *S. bosqueella*; L: other lepidopterans.

We thank Amábilio José Aires de Camargo from EMBRAPA Cerrados, Brasília, Federal District, Brazil, for the confirmation of *S. bosqueella* and other lepidopterans.

Summary

Stegasta bosqueella Chambers (Lepidoptera: Gelechiidae) is one of the principal pests of Brazilian peanut crops. Traps made from plastic bottles and baited with molasses sugar can be used to attract *S. bosqueella*. Molasses-baited traps are a promising tool for the monitoring and control of this insect.

Key Words: *Stegasta bosqueella*; sugarcane molasses; monitoring; attractant trap; action threshold

Sumario

Stegasta bosqueella Chambers (Lepidoptera: Gelechiidae) es una de las principales plagas de los cultivos de maní brasileños. Las trampas hechas con botellas de plástico y cebadas con melaza de caña de azúcar pueden ser usadas para atraer a *S. bosqueella*. Las trampas cebadas con melaza son una herramienta promisoría para el monitoreo y control de este insecto.

Palabras Clave: *Stegasta bosqueella*; melaza de caña de azúcar; moritoreo; trampa atractiva; umbral de acción

References Cited

- Almeida RP de. 2013. Manejo de insetos-praga na cultura de amendoim, pp. 333–390. In Pereira A. [ed.], O Agronegócio do Amendoim no Brasil, 2 ed. Embrapa Algodão, Brasília, Brazil.
- Altamar P, Pérez D, Rodríguez R, Guerrero C, Coy E. 2016. Chemical composition of the low-polar fraction of the *Copitarsia unclata* Burgos & Leiva (Lepidoptera: Noctuidae) eversible pheromone gland. Neotropical Entomology 45: 734–739.
- Boiça Junior AL, Chagas-Filho NR, Godoy IJ, Lourenção AL, Souza JR. 2012. Avaliação de resistência de cultivares de amendoim de hábito de crescimento rasteiro a *Enneothrips flavens* Moulton (Thysanoptera: Thripidae). Arquivos do Instituto Biológico 79: 33–38.
- Bosa CF, Clavijo A, Frida M, Marina A, Witzgall P. 2011. Respuesta de *Tecia solanivora* (Lepidoptera: Gelechiidae) a compuestos volátiles de papa, *Solanum tuberosum*. Revista Colombiana de Entomología 37: 1–7.
- Cisneros FH. 1995. Control Etológico. Control de Plagas Agrícolas. Lima: CIP, Horticulture International (AgriFoodGateway) Database, Department of Horticultural Science, North Carolina State University. Available online: <https://hortintl.cals.ncsu.edu/articles/control-etologico-de-plagas> (last accessed 17 Feb 2017).
- Cisneros FH. 2010. Caso 5. Programa-MIP de Espárrago en Chavimochic. Control de Plagas: MIP. Centro Internacional de la Papa (CIP), Lima, Peru. Available online: http://www.avocadosource.com/books/cisnerosfausto1995/cpa_caso_5.pdf (last accessed 17 Feb 2017).
- Garza E. 2002. Manejo integrado de plagas del jitomate en la planicie huasteca. Technical Bulletin 9. Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias-Centro de Investigación Regional Noreste (INIFAP-CIRNE), San Luis Potosí, Mexico.
- Jiménez SF. 2010. Empleo de trampas rústicas en el control de plagas de insectos, In Curso-taller nacional “Manejo Agroecológico de Plagas en la Agricultura Suburbana”. Programa de Agricultura Urbana y Suburbana. Instituto de Investigaciones de Sanidad Vegetal- Instituto de Investigaciones Fundamentales en Agricultura Tropical (INISAV-INIFAT), La Habana, Cuba.
- Lorya R, Chinchilla C, Domínguez J, Mexzón R. 2002. Una trampa efectiva para capturar adultos de *Opsiphanes cassina* Felder (Lepidoptera: Brassolidae) y observaciones sobre el comportamiento de la plaga en palma de aceite. PALMAS 23: 29–37.
- Parra E, Peña J, Esparza D, Labarca M. 2009. Evaluación de sustratos orgánicos y en combinación con insecticida para la captura de adultos de *Opsiphanes cassina* Felder en una plantación de palma aceitera (*Elaeis guineensis* Jacquin.) en el estado Zulia, Venezuela. Revista de la Facultad de Agronomía 26: 455–469.
- Raga A, Amaro R, Dinardo W, Strikis PC. 2006. Eficácia de atrativos alimentares na captura de mosca-das-frutas em pomar de citros. Bragantia 65: 337–345.
- Santos RF, Todeschini A. 2012. Evolução e perspectiva da cultura do amendoim para biocombustível no Brasil. Revista Brasileira de Energias Renováveis 1: 197–220.
- Scarpellini JR, Tavares P, Gentilin O. 2014. Controle da lagarta-do-pescoço-vermelho *Stegasta bosqueella* (Chambers, 1875) (Lepidoptera: Gelechiidae) na cultura do amendoim com produtos em pulverização no sulco de plantio ou foliares, In VIII Workshop Agroenergia: Matérias Primas. Ribeirão Preto, São Paulo, Brazil.