

Genetic Differentiation of Populations of a Hydrothermal Vent-Endemic Gastropod, Ifremeria nautilei, between the North Fiji Basin and the Manus Basin revealed by Nucleotide Sequences of Mitochondrial DNA

Authors: Kojima, Shigeaki, Segawa, Ryoko, Fujiwara, Yoshihiro, Hashimoto, Jun, and Ohta, Suguru

Source: Zoological Science, 17(8): 1167-1174

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zsj.17.1167

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Genetic Differentiation of Populations of a Hydrothermal Vent-Endemic Gastropod, *Ifremeria nautilei*, between the North Fiji Basin and the Manus Basin revealed by Nucleotide Sequences of Mitochondrial DNA

Shigeaki Kojima^{1*}, Ryoko Segawa², Yoshihiro Fujiwara³, Jun Hashimoto³ and Suguru Ohta¹

 ¹Ocean Research Institute, University of Tokyo, 1-15-1 Minamidai, Nakano, Tokyo 164-8639, Japan
²Department of Biology, Faculty of Science, Tokyo Metropolitan University, 1-1 Minami-Osawa, Hachioji, Tokyo 192-0364, Japan
³Japan Marine Science and Technology Center, 2-15 Natsushima, Yokosuka, Kanagawa 237-0061, Japan

ABSTRACT—The genetic differentiation of populations of a hydrothermal vent-endemic gastropod, *lfremeria nautilei*, between two back-arc basins in the south Western Pacific, namely the Manus Basin and the North Fiji Basin, was analyzed on the basis of nucleotide sequences of the mitochondrial gene for cytochrome oxidase I. The two populations of *I. nautilei* had no common haplotypes and appeared, therefore, to be isolated from one another. All haplotypes obtained from the North Fiji Basin formed a monophyletic group supported by a high bootstrap probability and the genetic diversity of the population in the North Fiji Basin was much smaller than that of the population in the Manus Basin. The present results suggest that the larval dispersal ability of *I. nautilei* might be lower than that of an undescribed species in the closely related genus *Alviniconcha*.

INTRODUCTION

Many hydrothermal vents with associated chemoautosynthesis-based faunal communities have been discovered in the south Western Pacific back-arc basins, namely, the Mariana Trough (Craig et al., 1987; Hessler and Lonsdale, 1991), the Manus Basin (Both et al., 1986; Tufar, 1990; Auzende et al., 1997), the North Fiji Basin (KAIYO 87 Shipboard Party, 1988; Desbruyères et al., 1994) and the Lau Basin (NAUTILAU Group, 1990; Desbruyères et al., 1994). The chemoautosynthesis-based faunal communities are composed of many endemic groups, such as vestimentiferan tube worms, large bivalves of the genera Calyptogena and Bathymodiolus, and crabs of the genus Bythograea (Tunnicliffe et al., 1998). Within such communities, those in the south Western Pacific are unique with respect to species composition (Tunnicliffe and Fowler, 1996). In the vent areas in the south Western Pacific, the communities are dominated by large endemic gastropods (Both et al., 1986; Okutani and Ohta, 1988), which

* Corresponding author: Tel. +81-3-5351-6469; FAX. +81-3-5351-6469. E-mail: kojima@ori.u-tokyo.ac.jp live in symbiosis with chemosynthetic bacteria (Stein *et al.*, 1988; Endow and Ohta, 1989; Windoffer and Giere, 1997). Furthermore, bivalves of the genus *Calyptogena*, which form a dominant group in the communities in both the Eastern Pacific and the north Western Pacific, have not been found in the south Western Pacific, with the exception of an undescribed species collected in the Desmos Cauldron of the Manus Basin (Ohta *et al.*, 1997) and the New Ireland Basin (Herzig *et al.*, 1994).

The dominant gastropods in the vent areas in the south Western Pacific have been classified into two genera in the family Provannidae, namely *Alviniconcha* and *Ifremeria*. Although *Alviniconcha* gastropods have been reported from the Mariana Trough, the Manus Basin, the North Fiji Basin and the Lau Basin, *Ifremeria* gastropods have not been discovered in the hydrothermal vent fields in the Mariana Trough (Bouchet and Warén, 1991; Warén and Bouchet, 1993). The absence of *Ifremeria* gastropods in the Mariana Trough might be attributable to lower larval dispersal ability of *Ifremeria* than *Alviniconcha*. The morphology of the larval shell suggests that *Alviniconcha* gastropods undergo planktotrophic development (Warén and Bouchet, 1993). The high-dispersal mode of



Fig. 1. Location of sites at which samples of *lfremeria nautilei* were collected: 1, Manus Basin; 2, North Fiji Basin. An open circle denotes the location of the Lau Basin.

development is thought to have been resulted in a wide geographical distribution of *Alviniconcha* gastropods. The degree of genetic differentiation between conspecific populations is expected to be closely related to the dispersal ability of the species. As expected from the high dispersal ability of the genus *Alviniconcha*, the absence of genetic differentiation between populations in the North Fiji Basin and the Manus Basin of an undescribed species in the genus *Alviniconcha*

Location	Site	Depth (m)	Submersible	Dive#	Ν
North Fiji Basin	White Lady site	1970	Nautile	D11	1
				D12	3
			Shinkai 6500	D77	9
				D78	1
				D80	3
	STARMER II site	1980	Nautile	D15	2
				D20	3
Manus Basin	PACMANUS site	1630	Shinkai 2000	D911	4
				D914	3
				D922	8
				D1062	9
				D1063	3
				D1067	3

Table 2. Primers used in the present study. Y, R, S and W denote T or C, A or G, G or C, and A or T, respectively. Positions refer to the corresponding amino acid residues encoded by the gene for mitochondrial cytochrome oxidase I from *Drosophila yakuba*.

Name	Sequence	Position	Direction
COI-B	5'-GGATGAACNGTNTAYCCNCC-3'	123–129	Forward
Gastro-3	5'-TTAGCTGGTGCTTCNTCNATYYTNGG-3'	150–158	Forward
COI-3	5'-GTNTGRGCNCAYCAYATRTTYACNGT-3'	285–293	Forward
TW-2	5'-ACTACRTARTANGTRTCRTG-3'	366–372	Reverse
COI-6	5'-GGRTARTCNSWRTANCGNCGNGGYAT-3'	434–442	Reverse

was reported (Kojima et al., 1998).

The larval type of *Ifremeria* gastropods remains unknown because of corrosion of the protoconch of all available specimens (Warén and Bouchet, 1993). In addition, no analysis of the genetic differentiation between populations of *Ifremeria* gastropods has been reported. In the present study, we analyzed the genetic differentiation of populations of *I. nautilei* between the Manus Basin and the North Fiji Basin (Fig. 1) by examining nucleotide sequences of mitochondrial DNA in an effort to characterize the larval dispersal ability of *Ifremeria*.

MATERIALS AND METHODS

During dives of the submersible "Nautile" of Institut français de recherche pour l'exploitation de la mer (IFREMER) and of the

submersibles "Shinkai 2000" and "Shinkai 6500" of the Japan Marine Science and Technology Center (JAMSTEC), a total of 52 specimens of *Ifremeria nautilei* was collected, as summarized in Table 1. One specimen of *Alviniconcha hessleri*, collected at the Forecast Vent site on the South Mariana Ridge (1470 m depth) during a dive of "Shinkai 6500" (No. D186), was used as the outgroup for phylogenetic analysis. Warén and Bouchet (1993) examined young specimens and the anatomy of *A. hessleri* and *I. nautilei*, and they stated that the genera *Alviniconcha* and *Ifremeria* are closely related and that these genera should be grouped into the family Provannidae. Beck (1991) also stated that *Alviniconcha* and *Olgaconcha* (*=Ifremeria*) are closely related to one another.

Mitochondrial DNA (mtDNA) was extracted from the deep-frozen head-foot region of each individual by a modified version of the method described by Komm *et al.* (1982). In the case of some damaged samples, which had been collected in the North Fiji Basin in 1990 and stored at -20° C, total DNA was extracted by grinding of

2 1 I TCT GTA AAC TTT ATT ACA ACG GTT ATT AAY ATR CGA TGA CGR GGA ATG CAA TTT GAG CGA TTR CCT YTA TTT 7 9 10 11 12 I GIW TGA TCT GTA AAA ATT ACY GCA RTT TTA TTG TTR CTT TCY CTY CCT GTG TTR GCC GGT GCT ATT ACA ATA 13 14 15 16 17 I CTT CTT ACT GAY CGA AAT TTT AAC ACT GCR TTC TTT GAY CCA GCT GGA GGT GGR GAY CCT GTG CTA TAT CAA 18 19 20 21 22 I CAT CTG TTY TGA TTY TTT GGA CAY CCA GAA GTA TAT ATY TTA ATT CTT CCT GGS TTT GGA ATG ATT TCT CAT 23 25 26 27 29 24 28 I ATY GTT AGA CAT TAT TCA GCT AAR AAA GAA ACT TTC GGA ACG YTR GGR ATG ATT TAT GCR ATG ATR GCA ATT A **T *** *** *** *** *** *** *** **G *** *** *** **T **G **A C*T **A **A **C *** **A *** **A *** *** 31 32 33 34 35 36 37 30 I GGA TTG YTA GGR TTT ATY GTT TGA GCT CAC CAT ATG TTT ACA GTR GGA ATR GAY GTT GAY ACY CGA GCT TAC COI-3 ----> 38 39 40 43 41 42 I TTY ACA GCT GCT ACT ATT ATT GCT GTT CCT ACA GGT ATT AAA GTT TTY AGT TGA CTT GCT ACR ATT YAT 44 45 47 I GGT TCA AAG ATT AAR TAT GAA ACT CCT ATR CTT TGA GCT TTA GGT TTT ATT TTY TTA TTY ACT GTA GGG GGA 48 49 50 I CTA ACT GER GTT GTT CTT GCA AAT TGT TCY TTA GAT ATT ATG ATG CAT GAT ACT TAY TAT GTA GTA GCT CAY ----- TW-2 52 53 55 56 57 54 I TTC CAT TAT GTT YTR TCT ATG GGA GCC GTG TTT GCC YTA TTC GCG GCT TTY AAT TAT TGA TTC CCT TTR YTA 58 I ACA GGT GTA ACT CTT CAC TCC CGG TGA ACA AAG GCC CAT TTC TAC ATC ATT TTT ATT GGT GTA AAT ATC ACT

Fig. 2. Nucleotide sequences of mitochondrial genes for cytochrome oxidase I from *Ifremeria nautilei* (I) and *Alviniconcha hessleri* (A). Asterisks indicate nucleotides that are identical to those in I. Numbers above the sequence from *I. nautilei* denote positions at which intraspecific variations were observed. Y, R, S and W denote T or C, A or G, G or C, and A or T, respectively. Underlining indicates sites at which amino acid substitutions were identified. Positions of primers COI-3 and TW-2 are shown. The nucleotide sequences will appear in the GSDB, DDBJ, EMBL and NCBI nucleotide sequence databases under accession numbers AB010097 (I) and AB010094 (A). samples, digestion with sodium dodecyl sulfate (SDS) and proteinase K, and extraction with phenol and chloroform. In the case of samples collected in the Manus Basin in 1998, total DNA was isolated from samples without digestion with proteinase K.

A fragment (about 450 bp) of the mitochondrial gene for cytochrome oxidase I (COI) was amplified by the polymerase chain reaction (PCR) with "universal" primers for amplification of most genes for metazoan COI, COI-3 and COI-6 (Shimayama et al., 1990). The conditions for PCR were as follows: 94°C for 60 s; then 30 to 40 cycles at 92°C for 40 s, 40°C for 60 s, and 72°C for 90 s. Genereleaser™ (BioVenture Inc., Murfreesboro, TN, USA) was used to sequester products of cell lysis that might have inhibited the polymerase. The nucleotide sequences of fragments (306 bp) were determined by the dideoxynucleotide chain-termination method using a Sequenase™

1

1

	11111111122222222333333333344444444445555555555	
Туре	1234567890123456789012345678901234567890123456789012345678	N
м1	CAGACATGGTTGTGTGTCCTTCTGTGAAGCATGGCTTCGACACGGCTGTCTCGTTACG	5
M2	***G**********************************	1
мз	**************************************	2
M4	**************************************	1
M5	** <u>A</u> **********************************	1
м6	******************C*C*C***************	1
M7	******A*******************************	1
M8	*G*G**********************************	1
м9	****TT************************G****C********	1
M10	****T************************G****C****C******	1
M11	****T************************G****C********	1
M12	T***T************************G****C****T********	1
M13	****TT************************G****C****T****T********	1
M14	****T*****A****************G****C**********	1
M15	****T*********************************	1
M16	****T***************************G****C*******	1
M17	****T***********T*********************	1
M18	****T************************G****C****C******	1
M1 9	**************************************	1
M20	T***T************************G****C****TG********	1
M21	****TT**A****A************G****C**********	1
M22	T***T***********T******G***G***C****C**	1
M23	T***T*C********T****T******G***G***C****C********	2
M24	****T****************C****CA****G****C****T*****C****A****	1
F1	***********C*C******G******G*****G*****G*****G******	1
F2	***********C*C******G******G*****G*****G*****G*****G****	1
F3	********C*C*C******G******G******G*****G******	1
F4	***********C*C******G*****G**G*********	4
F5	***********C*C******G*****TG*****G*****G********	1
F6	****T******C*C*****G*****G**G**********	1
F 7	***********C*C*****G*****GA*G******G*******G********	1
F8	***********C*C******G*****G**G****C**G******	1
F9	***********C*C******G*****G**G*********	1
F10	***********C*C******G*****G**G*********	1
F11	**********C*C*****G*****G**G*****G***G*	2
F12	**********C*C*****G*****G**G**********	1
F13	**********C*C*****G*****G**G**********	1
F14	***********C*C******G*****G**G*********	1
F15	T*********C*C******G*****G**G**********	1
F16	********C**C*C*C******G*****G**G********	1

1170

Fig. 3. Nucleotides at variable positions in the sequence of the gene for COI from Ifremeria nautilei. Numbers refer to the positions shown in Fig. 2. Asterisks indicate nucleotides that are identical to those in the uppermost sequences. M and F, as prefixes in names of haplotypes, denote haplotypes obtained from specimens from the Manus Basin and those from the North Fiji Basin, respectively.

Site

PCR product sequencing kit (United States Biochemical Coop., Cleveland, OH, USA). A longer fragment (about 960 bp) of genes for COI was amplified by PCR using primers COI-B (Hasegawa *et al.*, 1996) and COI-6. The conditions for PCR were as follows: 94°C for 60 s; then 30 to 40 cycles at 92°C for 40 s, 50°C for 60 s, and 72°C for 90 s. Nucleotide sequences within the upper region of this fragment were determined with a sequencer (DSQ-2000L; Shimazu Corp., Kyoto, Japan) using primers TW-2 (Kojima *et al.*, 1997) and Gastro-3. The sequence of primer Gastro-3 was based on sequences from some specimens of *I. nautilei* that had been determined using primer COI-B (data not shown). Nucleotide sequences of primers used in the present study are summarized in Table 2. Amino acid sequences of COI were deduced by reference to the modified genetic code of molluscan mtDNA (Shimayama *et al.*, 1990; Hoffmann *et al.*, 1992).

The genetic distance between sequences was calculated by Kimura's two-parameter method (Kimura, 1980). Phylogenetic trees were constructed by the neighbour-joining method (Saitoh and Nei, 1987) with the program from the MEGA package, Version 1.0 (Kumar *et al.*, 1993) and by the maximum parsimony method with the computer program Parsimony, which was provided by Dr. K. Tamura (Tokyo Metropolitan University). Number of net nucleotide substitution between populations (Nei, 1987) was estimated on the basis of numbers of nucleotide substitution between haplotypes calculated by

Kimura's two-parameter method (Kimura, 1980). Genetic diversity of populations was estimated in terms of two indices: gene diversity (Nei, 1987) and nucleotide diversity (Tajima 1983; Nei, 1987). Estimations were made with the computer program Arlequin (Schneider *et al.*, 1996). Gene diversity is the probability that two randomly chosen haplotypes are different. Nucleotide diversity is the probability that two randomly chosen homologous nucleotides are different.

RESULTS

Partial sequences (792 bp) of mitochondrial genes for COI were determined for 52 specimens of *Ifremeria nautilei* and one specimen of *Alviniconcha hessleri* (Figs. 2 and 3). Two intraspecific amino acid substitutions were detected in the sequences from the specimens of *I. nautilei*. Each amino acid substitution was detected in the case of only one specimen collected in the Manus Basin. Six amino acid substitutions were detected when we compared the dominant sequence from *I. nautilei* and the sequence from *A. hessleri* (Fig. 2).

Within the sequences of *I. nautilei*, 58 sites were found to



Fig. 4. Phylogenetic tree for haplotypes of *Ifremeria nautilei*. The tree was constructed by the neighbour-joining method with *Alviniconcha hessleri* as the outgroup. Bootstrap values are shown below branches of clades that are supported by bootstrap values of more than 50%.



Fig. 5. A 50% majority rule consensus phylogenetic tree for haplotypes of *Ifremeria nautilei*. The tree was constructed by the maximum parsimony method with *Alviniconcha hessleri* as the outgroup. Bootstrap values are shown above branches of clades that are supported by bootstrap values of more than 50%.

Table 3. Genetic diversity of populations of *Ifremeria nautilei*, as estimated from two indices, with standard deviations.

Diversity index	Population		
	Manus Basin	North Fiji Basin	
Gene diversity Nucleotide diversity	0.97±0.02 0.0075±0.0041	0.97±0.03 0.0032±0.0020	

be polymorphic and 42 haplotypes were obtained from 52 individuals (Fig. 3). Between the population of *I. nautilei* in the North Fiji Basin and that in the Manus Basin, five nucleotide substitutions were fixed (Fig. 3) and the two populations had no common haplotypes.

The phylogenetic relationships among the deduced haplotypes of *I. nautilei* were analyzed by the neighbour-joining (NJ) method and the maximum parsimony (MP) method. Individuals collected in both basins formed reciprocally monophyletic clusters on the NJ tree (Fig. 4). While monophyly of haplotypes from the North Fiji Basin was supported by a high bootstrap probability (88%), that of haplotypes from the Manus Basin was supported by a very low bootstrap probability (26%). On the MP tree (Fig. 5), haplotypes from the North Fiji Basin formed a monophyletic cluster supported by a high bootstrap probability (82%) and a bootstrap probability supporting the monophyly of haplotypes from the Manus Basin was low (45%).

The number of net nucleotide substitution between two

populations was estimated to be 0.0082. The genetic diversity of the two populations is summarized in Table 3. While gene diversity values were almost the same in the Manus Basin and the North Fiji Basin, the genetic diversity in the Manus Basin, as estimated in terms of nucleotide diversity, was about twice that in the North Fiji Basin. These results mean that diversity of haplotypes was the same in both populations but the average extent of genetic divergence among individuals was much greater in the Manus Basin than in the North Fiji Basin.

DISCUSSION

In the present study, we analyzed the genetic differentiation between the Manus Basin and the North Fiji Basin of populations of *Ifremeria nautilei*, one of the dominant species in the hydrothermal areas in the south Western Pacific. We found that these populations had no common haplotypes and that five nucleotide substitutions were fixed between them (Figs. 2 and 3). Thus, they appeared to be geographically isolated from one another.

For 20 and 12 individuals of an undescribed species in the genus *Alviniconcha* in the Manus Basin and the North Fiji Basin, respectively, Kojima *et al.* (1998) determined nucleotide sequences of the same region of the mitochondrial gene for COI as the region analyzed for *I. nautilei* in the present study. Both the exact test of population differentiation (Raymond and Rousset, 1995) and the randomized chisquared test of independence (Roff and Bentzen, 1989) showed the absence of genetic differentiation between populations of *Alviniconcha* sp. in the Manus Basin and the North Fiji Basin (Kojima *et al.*, 1998).

The undescribed species in the genus *Alviniconcha* and *I. nautilei* live close to the same hydrothermal vents in the both basins. Thus, the present results suggest that the larval dispersal ability of *I. nautilei* might be lower than that of *Alviniconcha* sp. While *Alviniconcha* gastropods were shown to undergo planktotrophic development, the larval type of *Ifremeria* gastropods remains unknown (Warén and Bouchet, 1993). To determine the larval type of *Ifremeria* gastropods, it is necessary to collect larvae or juveniles at early development stages for direct observations using a submersible with a sampling system such as a slurpgun. The analysis of DNA sequences should be very useful for identification of such samples (Dixon *et al.*, 1995).

On the neighbour-joining tree (Fig. 4) and the maximum parsimony tree (Fig. 5), all haplotypes of specimens of *I. nautilei* obtained from the North Fiji Basin formed a monophyletic cluster that was supported by high bootstrap probabilities. The cluster from the Manus Basin was supported only by very low bootstrap probabilities and, therefore, the monophyly of individuals from the Manus Basin is open to question. They might be paraphyletic with exclusion of individuals from the North Fiji Basin.

The genetic diversity of the population of I. nautilei in the North Fiji Basin was much smaller than that of the population in the Manus Basin. The limited genetic divergence and the clear monophyly of haplotypes from the North Fiji Basin can be explained if we postulate that *I. nautilei* in the North Fiji Basin is derived from a relatively recent dispersion from another basin. If the population in the North Fiji Basin was founded by a few immigrants and if the time since their arrival has been insufficient for creation of high genetic diversity via the introduction of genetic mutations, the genetic diversity of this population would be expected to remain at a low level. Since the environments around hydrothermal vents are unstable, it is also likely that large fluctuations in population size have occurred very frequently. If the population of *I. nautilei* in the North Fiji Basin were to have recently experienced a severe decrease, such a decrease might also have reduced the genetic diversity of the population. As I. nautilei and Alviniconcha sp. inhabit the same vent areas in the both basins, the genetic diversity of the population of Alviniconcha sp. in the North Fiji Basin should be lower than that in the Manus Basin, if low genetic diversity of the population of *I. nautilei* in the North Fiji Basin is attributed to the bottleneck effect caused by the change of the environments in the North Fiji Basin. Between the population of Alviniconcha sp. in the Manus Basin and that in the North Fiji Basin, little difference in the genetic diversity was shown in terms of both gene diversity (0.86±0.06 for the Manus Basin and 0.85±0.10 for the North Fiji Basin) and nucleotide diversity (0.0035±0.0022 for the Manus Basin and 0.0030±0.0020 for the North Fiji Basin) (unpublished data). The decrease of population size in the past seems unlikely to explain the low genetic diversity of the population of *I. nautilei* in the North Fiji Basin.

The hydrothermal areas in the Manus Basin is the westernmost habitat of I. nautilei and the nearest habitat is situated in the North Fiji Basin. The St. Georges Undercurrent and the Equatorial Undercurrent flow from the Manus Basin to the North Fiji Basin (Butt and Lindstrom, 1994). If individuals of I. nautilei in the Manus Basin are paraphyletic, with exclusion of individuals in the North Fiji Basin, it seems likely that the population in the North Fiji Basin was founded by immigrants from the Manus Basin and that the two populations have not yet become reciprocally monophyletic, even though each population has been accumulating genetic differences. If the rate of sequence divergence of the COI gene of snapping shrimps, namely 0.014 per million years (Knowlton and Weigt, 1998) is used, we can estimate that the divergence between the population in the North Fiji Basin and that in the Manus Basin occurred 0.59 million years ago. Such a time frame suggests that the recruitment of *I. nautilei* to the North Fiji Basin was a relatively recent event.

ACKNOWLEDGMENTS

The authors thank the shipboard parties of the STARMER cruises, the BIOACCESS-Manus 96 cruise and the BIOACCESS-Manus 98 cruise; the operation teams of the submersibles "Nautile", "Shinkai 2000" and "Shinkai 6500"; and the officers and crew of the tender ships "Nadier", "Natsushima" and "Yokosuka" for their help in collecting samples. Dr. K. Tamura graciously provided the computer program Parsimony. Part of this study was supported by grants from the University of Tokyo and from the Science Technology Agency of Japan.

REFERENCES

- Auzende J-M, Hashimoto J, Fiala-Médioni A, Ohta S, l'équipe Bioaccess (1997) In situ geological and biological study of two hydrothermal zones in the Manus Basin (Papua New Guinea). C R Acad Sci Paris 325: 585–591 (in French with English abstract and abridgment)
- Beck LA (1991) Olgaconcha tufari n. gen. et n. sp. a new mesogastropod (Gastropoda: Prosobranchia) from hydrothermal vents in the Manus back-arc basin (Bismarck Sea, Papua New Guinea). Ann Naturhist Mus Wien 92B: 277–287
- Both R, Crook K, Taylor B, Brogan S, Chappell B, Frankel E, Lui L, Sinton J, Tiffin D (1986) Hydrothermal chimneys and associated fauna in the Manus back-arc basin, Papua New Guinea. EOS Trans AGU 67: 489–490
- Bouchet P and Warén A (1991) *Ifremeria nautilei*, a new gastropod from hydrothermal vents, probably associated with symbiotic bacteria. C R Acad Sci Paris 612 Série III: 495–501
- Butt J, Lindstrom E (1994) Currents off the east coast of New Ireland, Papua New Guinea, and their relevance to regional undercurrents in the Western Equatorial Pacific Ocean. J Geophys Res 99: 12503–12514
- Craig H, Horibe Y, Farley KA, Welham JA, Kim K-R, Hey RN (1987) Hydrothermal vents in the Mariana Trough: results of the first Alvin dives. EOS Trans AGU 68: 1531
- Dixon DR, Jollivet DASB, Dixon LRJ, Nott JA, Holland PWH (1995) The molecular identification of early life-history stages of hydrotheraml vent organisms. Geol Soc Sp Pub 87: 343–350
- Desbruyéres D, Alayse-Danet A-M, Ohta S, Scientific parties of

BIOLAU and STARMER cruises (1994) Deep-sea hydrothermal communities in Southwestern Pacific back-arc basins (the North Fiji and Lau Basins): composition, microdistribution and food web. Mar Geol 116: 227–242

- Endow K, Ohta S (1989) The symbiotic relationship between bacteria and a mesogastropod snail, *Alviniconcha hessleri*, collected from hydrothermal vents of the Mariana back-arc Basin. Bull Jap Soc Microbial Ecol 3: 73–82
- Hasegawa T, Yamaguchi T, Kojima S, Ohta S (1996) Phylogenetic analysis among three species of the genus *Tetraclita* (Cirripedia: Balanomorpha) by nucleotide sequences of a mitochondrial gene. Benthos Res 51: 33–39 (in Japanese with English abstract)
- Herzig P, Hannington M, McInnes B, Stoffers P, Villinger H, Seifert B, Binns R, Liebe T (1994) Submarine volcanism and hydrothermal venting studied in Papua, New Guinea. EOS Trans AGU 75: 513– 516
- Hessler RR, Lonsdale P (1991) Biogeography of Mariana Trough hydrothermal vent communities. Deep-Sea Res 38: 185–199
- Hoffmann RJ, Boore JL, Brown WM (1992) A novel mitochondrial genome organization for the blue mussel, *Mytilus edulus*. Genetics 131: 397–412
- KAIYO 87 Shipboard Party (1988) Rift system in the North Fiji Basin: results of Japan-France cooperative research on board KAIYO 87. La mer 26: 36–46 (in Japanese with English abstract)
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequence. J Mol Evol 16: 111–120
- Knowlton N, Weigt LA (1998) New dates and new rates for divergence across the Isthmus of Panama. Proc R Soc Lond B265: 2257–2263
- Kojima S, Ohta S, Fujiwara Y, Fujikura K, Hashimoto J (1998) Speciation of gastropods of the genus *Alviniconcha* in the South Western Pacific. JAMSTEC J Deep Sea Res 14: 501–505 (in Japanese with English abstract)
- Kojima S, Segawa R, Hashimoto J, Ohta S (1997) Molecular phylogeny of vestimentiferans collected around Japan revealed by the nucleotide sequences of mitochondrial DNA. Mar Biol 127: 505– 513
- Komm B, Michaels A, Linton J (1982) Isolation and characterization of the mitochondrial DNA from the Florida spiny lobster, *Panulirus argus*. Comp Biochem Physiol 73B: 923–929
- Kumar S, Tamura K, Nei M (1993) MEGA: molecular evolutionary genetics analysis Version 1.0. The Pennsylvania State University, University Park, PA
- NAUTILAU Group (1990) Hydrothermal activity in the Lau Basin. First results from the NAUTILAU Cruise. EOS Trans AUG 71: 678– 679

- Nei M (1987) Molecular Evolutionary Genetics. Columbia Univ Press, New York
- Ohta S, Hashimoto J, Participants of BIOACCESS-Manus 96 Cruise (1997) Hydrothermal vent fields and vent-associated biological communities in the DESMOS Cauldron, eastern Manus Basin. JAMSTEC Deep Sea Res 13: 233–242 (in Japanese with English abstract)
- Okutani T, Ohta S (1988) A new gastropod mollusc associated with hydrothremal vents in the Mariana back-arc basin, Western Pacific. Venus (Jap Jour Malac) 47: 1–9
- Roff DA, Bentzen P (1989) The statistical analysis of mitochondrial DNA polymorphisms and the problem of small samples. Mol Biol Evol 6: 539–545
- Raymond M, Rousset F (1995) An exact test for population differentiation. Evolution 49: 1280–1283
- Saitoh N, Nei M (1987) The neighbour-joining method: a new method for reconstructing phylogenetic trees. Mol Biol Evol 10: 471–483.
- Schneider S, Kueffer J-M, Roessli D, Excoffer L (1996) Arlquin, a software package for population genetics. Genetics and Biometry Lab., Dept. of Anthropology, Univ. of Geneva, Geneva
- Shimayama T, Himeno H, Sasuga J. Yokobori S, Ueda T, Watanabe K (1990) The genetic code of a squid mitochondrial gene. Nucleic Acid Symp Ser 22: 77–78
- Stein JL, Cary SC, Hessler RR, Ohta S, Vetter RD, Childress JJ, Felbeck H (1988) Chemoautotrophic symbiosis in a hydrothermal vent gastropod. Biol Bull 174: 373–378
- Tajima F (1983) Evolutionary relationship of DNA sequences in finite populations. Genetics 105: 437–460
- Tufar W (1990) Modern hydrothermal activity, formation of complex massive sulfide deposits and associated vent communities in the Manus back-arc basin (Bismark Sea, Papua New Guinea). Mitt Österr Geol Ges 82: 183–210
- Tunnicliffe V, Fowler CMR (1996) Influence of sea-floor spreading on the global hydrothermal vent fauna. Nature 379: 531–533
- Tunnicliffe V, McArthur AG, McHugh D (1998) A biogeographical perspective of the deep-sea hydrothermal vent fauna. Adv Mar Biol 34: 353–442
- Warén A, Bouchet P (1993) New records, species, genera, and a new family of gastropods from hydrothermal vents and hydrocarbon seeps. Zool Scripta 22: 1–90
- Windoffer R, Giere O (1997) Symbiosis of the hydrothermal vent gastropod *Ifremeria nautilei* (Provannidae) with endobacteria - structural analyses and ecological considerations. Biol Bull 193: 381– 392

(Received September 22, 1999 / Accepted May 24, 2000)