



Vitamin A Insufficiency Accelerates the Decrease in the Number of Sperm Induced by an Environmental Disruptor, Bisphenol A, in Neonatal Mice

Authors: Nakahashi, Keiko, Matsuda, Manabu, and Mori, Takao

Source: Zoological Science, 18(6) : 819-821

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.18.819>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

[SHORT COMMUNICATION]

Vitamin A Insufficiency Accelerates the Decrease in the Number of Sperm Induced by an Environmental Disruptor, Bisphenol A, in Neonatal Mice

Keiko Nakahashi*, Manabu Matsuda and Takao Mori

*Department of Biological Sciences, Graduate School of Science,
The University of Tokyo, Tokyo 113-0033, Japan*

ABSTRACT—Exposure of neonatal mice to an estrogenic endocrine disruptor, bisphenol A, resulted in a malfunction of the testes when the animals became adults. The effect of bisphenol A was cancelled out by concurrent administration of retinol acetate, a naturally occurring metabolite of vitamin A. In contrast, the effect of endocrine disruption became more severe in mice neonatally exposed to bisphenol A and nursed by mothers fed a vitamin A-deficient diet only a few days before and after parturition. These results clearly show that maternal vitamin A is important for relieving in a baby the effect of endocrine disruption caused by environmental xenoestrogens, and suggest that the changes in the content of vitamin A and similar physiological factors in the habitat may be worth considering in studies on environmental disruptors.

INTRODUCTION

Recently, endocrine disruptors that cause a hormonal imbalance in animals have received a great deal of attention (Carson, 1962; Colborn *et al.*, 1993, 1996; Cadbury, 1997), with reports of infertility in female sheep induced by a phytoestrogen (Obst *et al.*, 1971), androgyny in conch caused by organic tin (Gibbs *et al.*, 1987), and small penis in alligators exposed to polychlorinated biphenyls (Guillette *et al.*, 1999), etc. Human health has also been affected by endocrine disruptors; diethylstilbestrol which was used to prevent abortion in the 1960's induced abnormalities of the reproductive organs, such as vaginal cancer, prostatic hyperplasia, and testicular tumor (Herbst, 1981). And, it is claimed that the number of fertile sperm decreases in the young as a result of exposure to endocrine disruptors during fetal life (Sharpe and Skekkebaek, 1993), although there are conflicting reports (Safe, 1995). Most of the environmental endocrine disruptors that affect vertebrates are categorized as estrogenic substances interacting with estrogen receptors. This is the reason why most efforts have concentrated on determining the estrogenic potency of all sorts of chemicals (Soto *et al.*, 1995). However, the action of estrogen may be modified by many factors. Here, we introduce bisphenol A (BPA) as an endo-

crine disruptor and vitamin A (VA) as an inhibitor of the effect of BPA on the mouse testis during the neonatal period.

Exogenous estrogen or estrogenic substances cause much more serious problems in the fetus and neonate than in adults. The genital tracts are particularly fragile during their development, the so-called critical period, and are often subjected to irreversible and fatal injuries even after short-term exposure to a low dose of estrogenic substances (Mori and Nagasawa, 1988). The critical period in the developing genital tracts corresponds to 3–4 months in the human fetus and to a few days pre- and postpartum in mice and rats. As the sensitivity differs between mother and child, mothers often keep taking estrogenic chemicals unaware of the effect to their children. BPA is one of the most widespread artificial estrogenic chemicals. For instance, it has been used in coating for the inside of cans, baby bottles, dental sealant, plastics for automobiles and many other products (Krishnan *et al.*, 1993; Olea *et al.*, 1996). Although BPA is easy to resolve and possesses rather weak estrogenic activity (10^2 – 10^4 times weaker than estradiol-17 β) (Nagel *et al.*, 1997), the concentration in certain environments is thought to be enough to evoke endocrine disruption in animals during the critical period.

VA is involved in regulating the function and differentiation of various epithelia, but is also a key factor inducing organogenesis in the embryo and fetus (Conlon, 1995). VA plays a role in spermatogenesis (van Pelt and Rooij, 1991) as well as oogenesis (Morita and Tilly, 1999). Estrogen, meanwhile, plays a crucial role in the reproductive activities of male as

* Corresponding author: Tel. +81-3-5841-4436;
FAX. +81-3-5841-4439.
E-mail: ss06334@mail.ecc.u-tokyo.ac.jp

well as female animals (Robertson *et al.*, 1999). Mahato *et al.* (2000) claimed that male germ cells do not require estrogen to develop or to function in fertilization, and suggested that estrogen acts within somatic cells of the male reproductive system, resulting in disruption of the excretion system of sperm. Previously, it was demonstrated that the concurrent injection of VA with estrogen into neonatal female mice inhibits the irreversible proliferation and cornification in the vaginal epithelium that is induced by neonatal treatment with estrogen alone (Mori, 1968). Therefore, we investigated the effects of BPA plus retinol acetate (RA) as VA on the reproductive function in neonatal male mice. Furthermore, the involvement of VA in the effect of BPA on the function was studied in neonatal mice nursed by mothers fed a VA-deficient diet before and after parturition.

MATERIALS AND METHODS

Male mice of the SHN strain were divided into 4 groups and given daily injections of 0.5 or 50 μ g of BPA, 50 μ g of BPA plus 100 IU of RA, or the vehicle only for 5 days from the day of birth. The vehicle consisted of 16 μ l of sesame oil used for BPA and 4 μ l of dimethylsulfoxide (DMSO) used for RA. The animals were weaned at 20 days of age and provided with a commercial diet (CE-7; Clea Japan Inc., Tokyo). They were killed at 14 weeks of age, and the sperm in the left caput epididymis were enumerated using a hemocytometer chamber under a microscope.

In addition, pregnant mice were fed a VA-deficient (VAD) diet (Low vitamin A diet; Clea Japan) from 3 days before gestation to postnatal day 5 of their pups. The pups nursed by the mothers taking the VAD diet were given 5 daily injections of BPA (0.5 μ g) or the vehicle only from the day of birth (VAD/BPA or VAD mice). After 5 days, the mothers were fed a normal VA-containing diet (CE-7). After being weaned at 20 days of age, the pups were also given the normal diet until autopsy at 14 weeks of age. At autopsy, the number of sperm was determined as mentioned above.

RESULTS AND DISCUSSION

In the present study, mice treated neonatally with a high dose (50 μ g for 5 days) of BPA showed a 35% reduction in the number of sperm in the caput epididymis as compared with unexposed controls when the animals became adults, while the treatment with a lower dose (0.5 μ g for 5 days) affected little the number of sperm (Fig. 1).

Interestingly, neonatal treatment with high-dose BPA failed to induce a reduction in the number of sperm if RA was injected concurrently (Fig. 1). Neonatal treatment with RA alone had no effect on the sperm count (data not shown). Thus, RA could nullify the effect of BPA on the testis to some extent in neonatal mice.

The VAD mice nursed by mothers fed a VAD diet only 8 days before and after parturition, showed no decrease in the number of sperm when they became adults. Surprisingly, in the VAD mice, neonatal administration of a low dose of BPA (VAD/BPA mice) caused a marked decrease in the number of sperm, although the same dose hardly affected the sperm count at all in mice nursed by normal mothers (Fig. 1).

Although the interaction of VA with BPA was not clarified,




Fig. 1. Changes in the number of sperm in mice receiving neonatal BPA in combination with or without RA. Cont, mice were given the vehicle only for 5 days from the day of birth; BPA 0.5 or BPA 50, mice were treated with 0.5 or 50 μ g BPA for 5 days from the day of birth; BPA 50+VA, mice received 50 μ g BPA in combination with 100IU RA as VA for 5 days from the day of birth; VAD/oil or VAD/BPA0.5, mice received vehicle or 0.5 μ g BPA for 5 days from the day of birth and were nursed by mothers taking a VAD diet for 8 days before and after parturition. The number in parentheses represents the number of mice examined.

* $P < 0.05$, ** $P < 0.01$

there are two possible explanation for the decrease in the number of sperm in epididymis. First, BPA has an irreversible effect on the spermatogenesis and/or the excretion of sperm by disrupting the action of VA. Second, VA has the ability to make the developing testis and/or epididymis insensitive to BPA. However, a possible effect of BPA and VA on the hypothalamo-hypophyseal system could not be excluded because BPA could affect the secretion of gonadotropic hormones in adult mice (Tohei *et al.*, 2001). The preventive effect of VA on the action of BPA in the developing male reproductive organs is similar to the previous finding that VA inhibits the effect of estrogen on the developing vaginal epithelium (Mori, 1968). In all events, the development of the reproductive organs including gonads may be controlled by a fine balance between two biologically active substances, estrogen and VA. A few papers concerning the interactions between VA and endocrine disruptors have been published: poultry chicks receiving a low VA diet and exposed to 3, 4, 3', 4'-tetrachlorobiphenyl become hypothyroid in comparison with unexposed controls (Spear and Moon, 1986) and hydroxylated metabolites of polychlorinated biphenyls inhibit VA transport in mice and turtles (Kramer *et al.*, 1997).

In conclusion, depletion of VA in pregnant or lactating mothers may predispose the fetus and pups to the adverse effects of BPA as well as many other estrogenic chemicals in the environment. Alternatively, the presence of relatively large amounts of xenoestrogens in combination with a lack of vitamin A or the presence of chemicals disrupting the actions of VA would severely affect the development of gonads as well as genital tracts. The intake of plenty of VA by a mother who has been accidentally exposed to an excess amount of estro-

gen may be effective in protecting her fetus or pups from endocrine disruption. Therefore, assessment of the anti-VA activity of all kinds of artificial chemicals in the environment is important in the research of endocrine disruptors. It should be stressed that the *in vivo* data of the threshold for estrogenic action of endocrine disruptors were obtained under nutritionally favorable conditions.

ACKNOWLEDGEMENTS

This work was supported by Grants-in-Aid for Scientific Research (C), Scientific Research on Priority Areas (A) and the Encouragement of Young Scientists from the Ministry of Education, Science, Sports and Culture, Japan.

REFERENCES

- Cadbury D (1997) The feminization of nature, our future at risk, Hamish Hamilton Ltd., London
- Carson R (1962) Silent Spring, Houghton Mifflin Company, Boston
- Colborn T, vom Saal FS, Soto AM (1993) Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ Health Perspect* 101: 378–384
- Colborn T, Dumanoski D, Myers JP (1996) Our Stolen Future, Dutton, New York
- Conlon RA (1995) Retinoic acid and pattern formation in vertebrates. *Trends Genet* 11: 314–319
- Gibbs PE, Bryan GW, Pascoe PL, Burt GR (1987) The use of the dog-whelk, *Nucella lapillus*, as an indicator of tributyltin (TBT) contamination. *J Mar Biol Ass U.K.* 67: 507–523
- Guillette LJ Jr, Woodward AR, Crain DA, Pickford DB, Rooney AA, Percival HF (1999) Plasma steroid concentrations and male phallus size in juvenile alligators from seven Florida lakes. *Gen Comp Endocrinol* 116: 356–372
- Herbst AL (1981) Developmental Effects of Diethylstilbestrol (DES) in Pregnancy, Thieme Stration, New York
- Kramer VJ, Helferich WG, Bergman A, Klasson-Wehler E, Giesy JP (1997) Hydroxylated polychlorinated biphenyl metabolites are anti-estrogenic in a stably transfected human breast adenocarcinoma (MCF-7) cell line. *Toxicol Appl Pharmacol* 144: 363–376
- Krishnan AV, Stathis P, Permuth SF, Tokes L, Feldman D (1993) Bisphenol-A: an estrogenic substance is released from polycarbonate flakes during autoclaving. *Endocrinol* 132: 2279–2286
- Mahato D, Goulding EH, Korach KS, Eddy EM (2000) Spermatogenic cells do not require estrogen receptor-alpha for development or function. *Endocrinol* 141: 1273–1276
- Mori T (1968) Effects of neonatal injections of estrogen in combination with vitamin A on the vaginal epithelium of adult mice. *Annot Zool Japan* 41: 113–118
- Mori T and Nagasawa H (1988) Toxicity of Hormones in Perinatal Life, CRC Press, Boca Raton
- Morita Y and Tilly JL (1999) Segregation of retinoic acid effects on fetal ovarian germ cell mitosis versus apoptosis by requirement for new macromolecular synthesis. *Endocrinol* 140: 2696–2703
- Nagel SC, vom Saal FS, Thayer KA, Dhar MG, Boechler M, Welshons WV (1997) Relative binding affinity-serum modified access (RBA-SMA) assay predicts the relative *in vivo* bioactivity of the xenoestrogens bisphenol A and octylphenol. *Environ Health Perspect* 105: 70–76
- Obst JM, Seamark RF, McGowan CJ (1971) Haemoglobin type and fertility of Merino ewes grazing oestrogenic (Yarloop clover) pastures. *Nature* 13: 497–498
- Olea N, Pulgar R, Perez P, Olea-Serrano F, Rivas A, Novillo-Fertrell A, Pedraza V, Soto AM, Sonnenschein C (1996) Estrogenicity of resin-based composites and sealants used in dentistry. *Environ Health Perspect* 104: 298–305
- Robertson KM, O'Donnell L, Jones ME, Meachem SJ, Boon WC, Fisher CR, Graves KH, McLachlan RI, Simpson ER (1999) Impairment of spermatogenesis in mice lacking a functional aromatase (cyp 19) gene. *Proc Natl Acad Sci USA* 96: 7986–7991
- Safe SH (1995) Environmental and dietary estrogen and human health: is there a problem? *Environ Health Perspect* 103: 346–351
- Sharpe RM and Skakkebaek NE (1993) Are oestrogens involved in falling sperm counts and disorders of the male reproductive tract? *Lancet* 341: 1392–1395
- Soto AM, Sonnenschein C, Chung KL, Fernandez MF, Olea N, Serrano O (1995) The E-SCREEN assay as a tool to identify estrogens: an update on estrogenic environmental pollutants. *Environ Health Perspect* 103: 113–122
- Spear PA and Moon TW (1986) Thyroid-vitamin A interactions in chicks exposed to 3,4,3',4'-tetrachlorobiphenyl: influence of low dietary vitamin A and iodine. *Environ Res* 40: 188–198
- Tohei A, Suda S, Taya K, Hashimoto T, Kogo H (2001) Bisphenol A inhibits testicular functions and increases luteinizing hormone secretion in adult male rats. *Exp Biol Med* 226: 216–221
- van Pelt AM and de Rooij DG (1991) Retinoic acid is able to reinitiate spermatogenesis in vitamin A-deficient rats and high replicate doses support the full development of spermatogenic cells. *Endocrinol* 128: 697–704

(Received May 21, 2001 / Accepted June 10, 2001)