

## **Establishing a monitoring baseline for threatened large ungulates in eastern Cambodia**

Authors: Gray, Thomas N.E., Phan, Channa, Pin, Chanrattana, and Prum, Sovanna

Source: Wildlife Biology, 18(4) : 406-413

Published By: Nordic Board for Wildlife Research

URL: <https://doi.org/10.2981/11-107>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Establishing a monitoring baseline for threatened large ungulates in eastern Cambodia

Thomas N.E. Gray, Channa Phan, Chanrattana Pin & Sovanna Prum

Monitoring ungulate populations is an essential part of wildlife management with ungulates performing essential ecosystem roles including structuring populations of large carnivores. A number of ungulate species in Southeast Asia are also globally threatened and are therefore important conservation targets in their own right. We estimated large ( $> 15$  kg) ungulate densities in two protected areas, i.e. Mondulkiri Protected Forest and Phnom Prich Wildlife Sanctuary, in eastern Cambodia using distance-based line transect sampling. During the 2009/2010 and 2010/2011 dry seasons, we surveyed 110 line transects (randomly distributed across 3,406 km<sup>2</sup>) for a total of 1,310 km. We used DISTANCE 6.0 to model detection functions from observations of banteng *Bos javanicus*, wild pig *Sus scrofa* and red muntjac *Muntiacus muntjak* generating estimates of group density, cluster size and individual density. Estimated densities  $\pm$  SE were  $1.1 \pm 0.2$  individual banteng/km<sup>2</sup>,  $1.4 \pm 0.4$  individual wild pig/km<sup>2</sup> and  $2.2 \pm 0.2$  individual red muntjac/km<sup>2</sup> giving an overall density of approximately 4.7 large ungulates/km<sup>2</sup>. Although wild pig and red muntjac densities were within the range of estimates reported from ecologically similar protected areas in tropical Asia, overall large ungulate density is much lower than the intrinsic carrying capacity of deciduous dipterocarp forest. This appears largely to be due to the scarcity of large deer (i.e. hog deer *Axis porcinus*, sambar *Cervus unicolor* and Eld's deer *Cervus eldi*) as a result of extensive historic hunting. Current large ungulate densities appear too low to support a viable tiger *Panthera tigris* population in the long term, and ungulate recovery, driven by strong protected area management, needs to be achieved before tiger populations can be restored.

**Please note that the supplementary information, including Appendix SI mentioned in this article, is available in the online version of this article, which can be viewed at [www.wildlifebiology.com](http://www.wildlifebiology.com).**

*Key words:* deciduous dipterocarp forest, distance sampling, Indochina, tiger prey species, wild cattle

Thomas N.E. Gray, Channa Phan & Chanrattana Pin, World Wide Fund for Nature Greater Mekong Cambodia Country Program, House #54, Street 352, Boeung Keng Kang I, Phnom Penh, Cambodia - e-mail addresses: [tomnegray@hotmail.com](mailto:tomnegray@hotmail.com) (Thomas N.E. Gray); [phanchannal@yahoo.com](mailto:phanchannal@yahoo.com) (Channa Phan); [chanratana.pin@gmail.com](mailto:chanratana.pin@gmail.com) (Chanrattana Pin) Sovanna Prum, World Wide Fund for Nature Greater Mekong Cambodia Country Program, House #54, Street 352, Boeung Keng Kang I, Phnom Penh, Cambodia, and Mondulkiri Forestry Administration Cantonment, Forestry Administration, Ministry of Agriculture Forestry and Fisheries, Phnom Penh, Cambodia - e-mail: [prumsovanna@gmail.com](mailto:prumsovanna@gmail.com)

*Corresponding author:* Thomas N.E. Gray

*Received 1 November 2011, accepted 29 July 2012*

*Associate Editor:* Christophe Bonenfant

Mainland Southeast Asia forms part of the Indo-Burma biodiversity hotspot and is thus a top priority for conservation and wildlife management (Myers et al. 2000). However, Southeast Asian biodiversity is chronically threatened due to the most rapid global rate of forest conversion and intense pressures on natural resources from high human population den-

sities and a huge market for medicinal wildlife products (Sodhi et al. 2004, 2009, Nijman 2010). Subsequently, the region supports the highest global concentration of threatened terrestrial mammals (Schipper et al. 2008) and active conservation management is necessary for stemming regional biodiversity losses (Koh & Sodhi 2010). A critical requirement of all

conservation projects is to have measures of effectiveness concerning the achievement of conservation goals. The importance of evidence-based conservation, with the success or failure of conservation interventions supported by robust monitoring data, is now widely accepted within the conservation community (Sutherland et al. 2004, Pullin & Knight 2009). However, in Southeast Asia, few examples exist of robust monitoring systems for assessing the health of populations of focal species and the impact of conservation activities upon them (but see O'Kelly & Nut 2010, Ryan et al. 2011).

Monitoring large ( $> 15$  kg) ungulate populations is an essential part of wildlife management with ungulates performing key ecosystem roles including structuring populations of large carnivores, dispersing seeds and influencing vegetation patterns (Adler et al. 2001, Karanth et al. 2004, Prasad et al. 2006). With the global focus on tiger *Panthera tigris* conservation and recovery (Seidensticker 2010), monitoring large ungulates in Asia is essential given that prey densities appear to be a key determinant of tiger population dynamics (Karanth et al. 2004). A number of regionally endemic, or near-endemic, large ungulates that exist in Southeast Asia are also globally threatened, and they are thus key conservation targets in their own right. These species include saola *Pseudoryx nghetinhensis*, wild water buffalo *Bubalus arnee*, banteng *Bos javanicus*, Eld's deer *Cervus eldii*, hog deer *Axis porcinus* and large-antlered muntjac *Muntiacus vuquangensis*. Despite the acknowledged importance of monitoring ungulate populations, there are no published large ungulate densities from anywhere in mainland Southeast Asia obtained using statistically robust survey methodologies.

The plains of northern and eastern Cambodia were formerly described as one of the 'great game-lands of the world; a Serengeti of Asia', and they supported a diverse and abundant megafauna of ungulates, predators and scavengers (Wharton 1957, Tordoff et al. 2005). However, Cambodia suffered considerable political instability and conflict throughout the 20th century intensifying during the Lon Nol (1970-1975) and Pol Pot (1975-1979) regimes (Chandler 2000). During this period, there is evidence of large declines in the regional population and distribution of large mammal species including tiger, leopard *P. pardus*, Asian elephant *Elephas maximus*, wild cattle and Eld's deer and hog deer (Duckworth & Hedges 1998, Loucks et al. 2008). These declines were associated with a proliferation of firearms, the develop-


ment of an external market for wildlife products and, particularly during the Khmer Rouge era, government-sponsored hunting (Loucks et al. 2008). This hunting pressure possibly led to the global extinction of one large ungulate species endemic to Indochina: the kouprey *Bos sauveli*. In this paper, we report the first robust ungulate density estimates from anywhere in mainland Southeast Asia in order to provide a monitoring baseline for two protected areas in eastern Cambodia. We also test the ecological prediction that large ungulate densities in eastern Cambodia are lower than ecologically similar protected areas in the Indian subcontinent due to intense historic hunting pressure, and we use our results to assess the potential for tiger population recovery in the landscape.

## Material and methods

### Study area

We estimated large ungulate densities in the core areas of two adjacent protected areas in eastern Cambodia: Mondulkiri Protected Forest (MPF; 3,500 km<sup>2</sup> approximately centered on 12.08°N, 106.05°E) and Phnom Prich Wildlife Sanctuary (PPWS; 2,700 km<sup>2</sup> approximately centered on 12.40°N, 107.00°E; Fig. 1). These lowland (approximately 100-300 m a.s.l.) protected areas are dominated by deciduous dipterocarp forest with smaller areas of mixed deciduous and semi-evergreen forest (Phan & Gray 2010). Seven species of large native ungulates are known to be present within MPF and PPWS: wild water buffalo, gaur *Bos gaurus*, banteng, sambar *Rusa unicolor*, Eld's deer, wild pig *Sus scrofa* and red muntjac *Muntiacus muntjak* (Phan et al. 2010, Gray et al. 2012). Since 2005, law enforcement efforts in both protected areas have been initiated to limit hunting pressure on these ungulate species, all of which are protected under Cambodian forestry law. The study area forms part of a level-one tiger conservation landscape (Sanderson et al. 2010) and has been identified as one of 12 critical tiger conservation and recovery landscapes by the World Wide Fund for Nature (Wikramanayake et al. 2011). Despite extremely low numbers, Lynham (2010) considered the landscape irreplaceable for Indochinese tiger *P. t. corbetti* conservation, representing the only large block of dry forest habitat in Southeast Asia with a reintroduction programme recommended.

Figure 1. Location of Mondulkiri Protected Forest (MPF) and Phnom Prich Wildlife Sanctuary (PPWS), eastern Cambodia, showing the core zones of both protected areas and the locations of line transects (MPF-outer core surveyed in 2009/2010, black lines; MPF-inner core surveyed in 2010/2011, white lines; PPWS surveyed in both years, grey lines).



## Line transect surveys

We estimated ungulate densities using distance-based line transect sampling. This methodology is standard for estimating ungulate tiger prey densities in protected areas in the Indian subcontinent (Karanth & Nichols 2002) and addresses two of the most problematic aspects of animal abundance estimation: spatial sampling and detectability (Williams et al. 2002, Thomas et al. 2010). We used the Survey design function in DISTANCE 6.0 (Thomas et al. 2010) to plot between 34 and 38 line transects in each of 1) the core zone of PPWS (1,670 km<sup>2</sup>), 2) MPF outer core (1,276 km<sup>2</sup>) and 3) MPF inner core (460 km<sup>2</sup>). The latter roughly corresponds to an area identified and proposed by the Cambodian government to be a strictly protected tiger recovery zone. Although hunting of all ungulates is prohibited

within the protected areas, illegal hunting occurs and, based on ranger-patrol data, appears higher in PPWS and MPF outer core than in MPF inner core.

Each line transect was 1–4 km long and each was surveyed between one and 10 times during the dry season of 2009/2010 (PPWS: 33 transects with a total of 155 km, and MPF outer core: 38 transects with a total of 273 km) and the dry season of 2010/2011 (PPWS: 34 transects with a total of 467 km, and MPF inner core: 38 transects with a total of 415 km). In PPWS, the same transects were surveyed in both 2009/2010 and 2010/2011 whilst in MPF, the outer core was surveyed in 2009/2010 and inner core in 2010/2011. Figure 1 indicates the locations of transects within the study area and indicates which transects were surveyed during which years.

Surveys followed the protocols of Karanth &

Nichols (2002) for line-transect sampling of ungulate tiger prey species with two observers slowly walking the line transects at dawn (start at 06:30-07:00) and dusk (finish at 17:30-18:00). All large ungulate observations were recorded with the species, number of animals (cluster size), distance between the animal or centre of a group of animals and the observers on the line (with a laser rangefinder), compass bearing to the animal or to the centre of a group of animals and compass bearing of the transect line noted.

Line transect data analysis

We used the Conventional Distance Sampling (CDS) engine in software DISTANCE 6.0 (Thomas et al. 2010) to estimate species density. In order to meet the recommendation of a minimum of 40-60 observations for fitting detection functions (Buckland et al. 2001), we estimated densities only for ungulate species with > 50 observations. Prior to modelling, data were right-truncated to prevent the inclusion of additional adjustment terms which fit a long tail to the detection function but reduce precision for little gain (Thomas et al. 2010). The model which best described the detection process was selected on the basis of Akaike Information Criteria values corrected for small sample size ( $AIC_c$ ). Due to the limited number of encounters of each species in each sampling stratum (i.e. PPWS, MPF outer core and MPF inner core), a single 'global' detection function was fitted to all detections of each species and this was used to calculate stratum-specific and landscape-wide densities. Using the selected model, we derived estimates of group density, cluster size and individual




Figure 2. Density estimates (number of individuals/km<sup>2</sup>) for red muntjac, banteng and wild pig in Mondulkiri Protected Forest and Phnom Prich Wildlife Sanctuary based on distance-based line transect sampling. Boxes indicate mean  $\pm$  one Standard Error; whiskers indicate upper and lower 95% confidence intervals.

density for each species. We estimated expected cluster size by regressing log-cluster size against the estimated probability of detection except for banteng where checking for size bias in detection of animal clusters led to a non-significant regression equation at  $\alpha=0.10$  (Drummer & McDonald 1987); therefore, we used the mean observed cluster size ( $5.1 \pm SE\ 0.6$ ) for analysis. In order for our density estimates to be comparable with those from ecologically similar landscapes in South Asia (e.g. Karanth & Nichols 2000, Bagchi et al. 2003, Wegge & Storaas 2009), and to assess the potential carrying capacity for tiger in our study area, we present, and discuss in the text, individual (i.e. number of individuals of each species estimated/km<sup>2</sup>) rather than group densities.

Table 1. Estimated large ungulate densities in Mondulkiri Protected Forest inner core (MPF-core), Mondulkiri Protected Forest outer core (MPF-outer), Phnom Prich Wildlife Sanctuary (PPWS) and across all three stratum (Landscape) based on distance-based line transect sampling. The table shows the number of observations included in models (N), density of groups ( $D_g$ ), mean cluster size (Y), density of individuals ( $D_i$ ) and population size (95% confidence interval range rounded to nearest 10) for banteng, wild pig and red muntjac.

Species	Stratum	N	$D_g \pm SE\ (km^{-2})$	Y	$D_i \pm SE\ (km^{-2})$	Population size
Banteng	MPF-core	31	$0.4 \pm 0.05$	5.1	$1.9 \pm 0.4$	600-1280
	MPF-outer	8	$0.2 \pm 0.05$		$0.8 \pm 0.3$	500-2200
	PPWS	12	$0.1 \pm 0.03$		$0.7 \pm 0.2$	600-2020
	Landscape	51	$0.2 \pm 0.03$		$1.1 \pm 0.2$	2700-5690
Wild pig	MPF-core	19	$0.4 \pm 0.07$	5	$1.9 \pm 0.5$	500-1470
	MPF-outer	14	$0.4 \pm 0.08$		$1.9 \pm 0.6$	1400-4330
	PPWS	15	$0.2 \pm 0.04$		$1.0 \pm 0.3$	1595-2860
	Landscape	48	$0.3 \pm 0.04$		$1.4 \pm 0.4$	890-7970
Red muntjac	MPF-core	82	$2.6 \pm 0.3$	1.1	$2.8 \pm 0.3$	1020-1660
	MPF-outer	43	$2.3 \pm 0.4$		$2.5 \pm 0.5$	2160-4590
	PPWS	57	$1.4 \pm 0.3$		$1.5 \pm 0.3$	1740-3710
	Landscape	182	$2.0 \pm 0.2$		$2.2 \pm 0.2$	6060-9030

## Results

We recorded six large ungulate species, gaur (three encounters), banteng (63 encounters), Eld's deer (two encounters), sambar (three encounters), wild pig (58 encounters) and red muntjac (198 encounters) during line transect surveys. However, sufficient encounters to model detection functions, and hence estimate density, were only obtained for banteng, wild pig and red muntjac. Details of the best-fitting models and detection curves for each species are given in the online only, supplementary information (Appendix SI). Landscape-wide mean densities  $\pm$  SE were  $1.1 \pm 0.2$  individual banteng/km<sup>2</sup>,  $2.2 \pm 0.2$  individual red muntjac/km<sup>2</sup> and  $1.4 \pm 0.4$  individual wild pig/km<sup>2</sup> giving an overall density of approximately 4.7 large ungulates/km<sup>2</sup> (Fig. 2). For all three species, stratum-specific density estimates were higher in MPF inner core than the MPF outer core and PPWS (Table 1). This difference was more pronounced for banteng than for wild pig and red muntjac. Estimated population sizes across the entire 3,406 km<sup>2</sup> study area were between 2,700 and 5,700 banteng (with an estimated population of 600–1,300 in MPF inner core), 3,000–8,000 wild pig (500–1,500 in MPF inner core) and 6,000–9,000 red muntjac (1,000–1,600 in MPF inner core; see Table 1).

## Discussion

Despite the importance of the region for globally threatened large ungulates (Tordoff et al. 2005) and tiger conservation (Simcharoen et al. 2007, Lynham 2010), no published estimates of ungulate densities, based on robust distance-based line transect sampling, exist for mainland Southeast Asia. We provide the first large ungulate density estimate for the region and report densities of approximately 4.7 large ungulates (banteng, wild pig and red muntjac combined)/km<sup>2</sup> in the core areas of MPF and PPWS, eastern Cambodia. However, this estimate excludes the three ungulate species detected during surveys which were encountered insufficiently often to reliably model detection functions and hence estimate density (i.e. gaur, sambar and Eld's deer), and it may therefore underestimate total ungulate density in the landscape. Fitting a single detection function to all encounters of large deer and wild cattle (i.e. banteng, gaur, sambar and Eld's deer; 66 observations) gives a landscape-wide density estimate  $\pm$  SE of  $1.7 \pm 0.4$  individuals/km<sup>2</sup>. Whilst within the 95% confidence

intervals of the density estimate based solely on banteng encounters, this is 30% higher than the mean banteng density estimate. This suggests that excluding gaur, sambar and Eld's deer encounters slightly negatively biased the overall density estimates for ungulates in the landscape.

The inner core of MPF supported higher densities of all three species of ungulates for which density could be estimated than the other two survey strata. The higher ungulate densities in MPF inner core are probably due to the area's remoteness and inaccessibility, and hence reduced hunting pressure, in comparison with MPF outer core and PPWS. Our data therefore provide strong and consistent support for the importance of the inner core area of MPF for large ungulate populations and this provides strong evidence in support of the proposal of the site as Cambodia's first strictly protected tiger recovery zone.

Published total (i.e. across all species sampled) wild large ungulate densities within South Asian protected areas range between seven (Jigme Singye Wangchuck National Park, Bhutan; Wang 2010) and > 250 (Bardia National Park, Bhutan; Wegge & Storaas 2009) individuals/km<sup>2</sup> with > 50 individual ungulates/km<sup>2</sup> being the norm in most Indian tiger reserves (Karanth & Nichols 2000). Our estimates, of < 5 individual ungulates/km<sup>2</sup> are thus lower than any published estimates for large ungulate densities in tropical Asia. This is despite the potential for high ungulate densities in deciduous dipterocarp forest. In ecologically similar lowland deciduous *Sal Shorea robusta* forest in Ranthambore Tiger Reserve, India, ungulate density is approximately 75 animals/km<sup>2</sup> (Bagchi et al. 2003). This suggests that ungulate populations in MPF and PPWS remain severely depressed by hunting. The low ungulate density, in comparison with ecologically similar South Asian protected areas, seems to result from the extremely low densities of large deer species i.e. hog deer, sambar and Eld's deer. Our estimates of red muntjac density are similar to those from published studies in India: red muntjac mean density  $2.1 \pm$  SD 1.5 individuals/km<sup>2</sup>; N = 6 sites: Wegge & Storaas (2009), Karanth & Sunquist (1992), Karanth & Nichols (2000), Jathanna et al. (2003) and Wang (2010). In contrast, large deer densities are much higher in South Asian protected areas e.g. mean sambar density  $7.0 \pm$  SD 6.3 individuals/km<sup>2</sup>; N = 8 sites: Harihar et al. (2008), Karanth & Nichols (2000), Karanth & Sunquist (1992), Bagchi et al. (2003), Jathanna et al. (2003) and Wang (2010) and Chittal *Axis axis*

mean density  $65.5 \pm \text{SD } 6.3$  individuals/km<sup>2</sup>; N = 8 sites: Wegge & Storaas 2009, Harihar et al. (2008), Karanth & Nichols (2000), Karanth & Sunquist (1992), Bagchi et al. (2003) and Jathanna et al. (2003).

The limited number of encounters with larger deer in MPF and PPWS suggests that populations of these species have been severely depressed by hunting; Wharton (1957) reported an abundance of Eld's deer throughout northern and eastern Cambodia in the 1950s. Whilst the low encounter rates of sambar and Eld's deer may be due to a behavioural effect, for example species-specific responses to illegal hunting, and it is inadvisable to assume that low encounter rates on line transects equates to low density, an independent sampling method in both protected areas, i.e. camera-trapping, has also produced few encounters of large deer. In > 7,000 camera-trap nights, in both MPF and PPWS, just five photographs of sambar were obtained compared with 160 of banteng and 330 of wild pig (Phan et al. 2010). Eld's deer have only been photographed in the landscape when camera-trapping has targeted seasonal waterholes known to be used by the species. Low densities and slow recoveries of large deer, even when other ungulate species are increasing, has been noted elsewhere in Southeast Asia (Aung et al. 2001, Steinmetz et al. 2010), and it may be that these species are slower to recover from hunting-induced population declines than other ungulates. In terms of tiger recovery, low densities of large deer are worrying as deer make up > ¼ of prey consumed by tiger across most of the species' range (Karanth & Nichols 2002).

Based on our estimates of ungulate density, it is possible to extrapolate the number of tigers which MPF and PPWS could currently support. Studies across a number of tiger range countries have demonstrated a positive relationship between tiger abundance and prey density (Karanth & Sunquist 1995, Karanth et al. 2004, Smith et al. 2011). At low ungulate prey densities, the evidence suggests a linear relationship with female home-range size calibrated so as to provide a similar prey biomass, approximately 120,000 kg/home range (Smith et al. 2011). Converting our mean estimates of landscape-wide ungulate density into biomass (assuming an average individual biomasses of banteng 300 kg, wild pig 50 kg and red muntjac 20 kg; Karanth & Sunquist 1992) suggests approximately 440 kg of ungulate biomass/km<sup>2</sup>. Using the relationship in Smith et al. (2011), and the model of Karanth et al. (2004), which applies an average kill rate of 50 ungulates/tiger/year from a

base population of 500 ungulates (i.e. off-take of 10%/year), suggests a potential female tiger home-range size in our eastern Cambodia study area of approximately 260 km<sup>2</sup>. This equates to a population of approximately 13 breeding female tigers across the 3,400 km<sup>2</sup> study area.

This is less than the 25 breeding female tigers (ca 75 individuals), identified as the threshold for a source site by Walston et al. (2010), and is well below the levels of prey required to support 50 breeding females (ca 150 tigers), which was suggested for long-term population viability (Walston et al. 2010). Therefore, whilst current ungulate prey levels in MPF and PPWS could support a small number of tigers, such a population would unlikely be viable in the long term. Based on estimates of intrinsic growth rates of Southeast Asian large ungulates ( $R_{\text{max}}$  0.3-0.5; Steinmetz et al. 2010), recovery to prey densities sufficient to support 25-50 breeding female tigers would take 6-12 years. However, for such a recovery of prey species to occur, strong protected area management, to reduce ongoing threats of both habitat loss and hunting, is required. Regular monitoring of large ungulates, through distance-based line transect surveys, is also essential to ensure that ungulate populations increase to sufficient numbers to support a viable tiger population prior to any reintroductions.

*Acknowledgements* - our study forms part of WWF Cambodia's Eastern Plains Landscape Project. Major funding comes from WWF-US, WWF-Sweden, WWF-Germany and Humanscale. Our work in MPF was with permission of the Forestry Administration of the Ministry of Agriculture Forestry and Fisheries with support from His Excellency Cheng Kimsun, Men Phymean, Song Keang, Keo Omaliss and Keo Sopheak. Our work in PPWS was with the permission of the Ministry of the Environment and support from His Excellency Chay Samith, Sanrangdy Vicheth and Han Sakhan. Khaev Oudom, Ing Seangnirithy, Lien Nor, Van Sonny (MPF) and Sary Tre, Sin Somoan, Se Noun and Chan Touy (PPWS) assisted in data collection, whilst Barney Long, Nick Cox, Bivash Pandav, Hannah O'Kelly, Craig Bruce, Keith Metzner and Seng Teak provided logistical and statistical support. Christophe Bonenfant, Heiko Wittmer and an anonymous reviewer provided comments which greatly improved the manuscript.

## References

- Adler, P.B., Raff, D.A. & Laurenroth, W.K. 2001: The effect of grazing on the spatial heterogeneity of vegetation. - *Oecologia* 128 (2): 465-479.
- Aung, M., McShea, W.J., Htung, S., Than, A., Soe, T.M., Monfort, S. & Wemmer, C. 2001: Ecology and social

- organization of a tropical deer (*Cervus eldi thamin*). - Journal of Mammalogy 82 (3): 836-847.
- Bagchi, S., Goyal, S.P. & Sankar, K. 2003: Prey abundance and prey selection by tigers (*Panthera tigris*) in a semi-arid, dry deciduous forest in western India. - Journal of the Zoological Society London 260 (3): 285-290.
- Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L. & Thomas, L. 2001: Introduction to distance sampling: estimating abundance of biological populations. - Oxford University Press, New York, New York, USA, 448 pp.
- Chandler, D. 2000: A history of Cambodia. 3rd edition. - Westview Press, Boulder, Colorado, USA, 232 pp.
- Drummer, T.D. & McDonald, L.L. 1987: Size bias in line transect sampling. - Biometrics 43 (1): 13-21.
- Duckworth, J.W. & Hedges, S. 1998: Tracking Tigers: A review of the status of tiger, Asian elephant, gaur and banteng in Vietnam, Lao, Cambodia and Yunnan (China). - WWF Indochina, Hanoi, Vietnam, 282 pp.
- Gray, T.N.E., Ou, R., Huy, K., Pin, C. & Maxwell, A.L. 2012. The status of large mammals in eastern Cambodia: a review of camera-trapping data 1999-2007. - Cambodian Journal of Natural History, pp. 42-55.
- Harihar, A., Pandav, B. & Goyal, S.P. 2008: Responses of tiger (*Panthera tigris*) and their prey to removal of anthropogenic influences in Rajaji National Park, India. - European Journal of Wildlife Research 55 (2): 97-105.
- Jathanna, D., Karanth, K.U. & Johnsingh, A.J.T. 2003: Estimation of large herbivore densities in the tropical forests of southern India using distance sampling. - Journal of Zoology (London) 261 (3): 285-290.
- Karanth, K.U. & Nichols, J.D. 2000: Ecological status and conservation of tigers in India. Final technical report to the Division of International Conservation, US Fish and Wildlife Service, Washington DC and Wildlife Conservation Society, New York. - Centre for Wildlife Studies, Bangalore, India, 115 pp.
- Karanth, K.U. & Nichols, J.D. 2002: Monitoring tigers and their prey: a manual for researchers, managers and conservationists in tropical Asia. - Centre for Wildlife Studies, Bangalore, India, pp. 139-152.
- Karanth, K.U., Nichols, J.D., Kumar, N.S., Link, W.A. & Hines, J.E. 2004: Tigers and their prey: predicting carnivore densities from prey abundance. - Proceedings of the National Academy of Science, USA 101 (14): 4854-4858.
- Karanth, K.U. & Sunquist, M.E. 1992: Population structure, density and biomass of large herbivores in the tropical forests of Nagarhole, India. - Journal of Tropical Ecology 8 (1): 21-35.
- Karanth, K.U. & Sunquist, M.E. 1995: Prey selection by Tiger, Leopard, and Dhole in tropical forests. - Journal of Animal Ecology 64 (4): 439-450.
- Koh, L.P. & Sodhi, N.S. 2010: Conserving south-east Asia's imperilled biodiversity: scientific, management and policy challenges. - Biodiversity & Conservation 19 (4): 913-917.
- Loucks, L., Mascia, M.B., Maxwell, K.A., Huy, K., Duong, N., Chea, L.B., Cox, N. & Seng, T. 2008: Wildlife decline in Cambodia, 1953-2005: exploring the legacy of armed conflict. - Conservation Letters 2 (2): 82-92.
- Lynham, A.J. 2010: Securing a future for wild Indochinese tigers: transforming tiger vacuums into tiger source sites. - Integrative Zoology 5 (4): 324-334.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, A.B. & Kent, J. 2000: Biodiversity hotspots for conservation priorities. - Nature 403: 853-858.
- Nijman, V. 2010: An overview of international wildlife trade from Southeast Asia. - Biodiversity & Conservation 19 (8): 1101-1114.
- O'Kelly, H. & Nut, M.H. 2010: Monitoring of ungulate, primate and peafowl populations using line transect surveys in Seima Protection Forest, Cambodia 2005-2010. - Wildlife Conservation Society Cambodia, Phnom Penh, Cambodia, 98 pp.
- Phan, C. & Gray, T.N.E. 2010: Ecology and natural history of Banteng in eastern Cambodia: evidence from camera-trapping in Mondulkiri Protected Forest and Phnom Prich Wildlife Sanctuary. - Cambodian Journal of Natural History 2 (2): 118-126.
- Phan, C., Prum, S. & Gray, T.N.E. 2010: Recent camera-trap records of globally threatened species from the Eastern Plains Landscape, Cambodia. - Cambodian Journal of Natural History 2 (2): 89-93.
- Prasad, S., Krishnaswamy, J., Chellam, R. & Goyal, S.P. 2006: Ruminant-mediated seed dispersal of an economically valuable tree in Indian dry forests. - Biotropica 38 (5): 679-682.
- Pullin, A.S. & Knight, T.M. 2009: Doing more good than harm - Building an evidence-base for conservation and environmental management. - Biological Conservation 142 (5): 931-934.
- Ryan, G.E., Dove, V., Trujillo, F. & Doherty, P.F. 2011: Irrawaddy dolphin demography in the Mekong River: an application of mark-resight models. - Ecosphere 2 (1): 58.
- Sanderson, E., Forrest, J., Loucks, C., Ginsberg, J., Dinerstein, E., Seidensticker, J., Leimgruber, P., Melissa, A., Heydlauff, A., O'Brien, T., Bryia, G., Klenzendorf, S. & Wikramanayake, E. 2010: Setting priorities for Conservation and Recovery of Wild Tigers: 2005-2015. The Technical Assessment. - In: Tilson, R. & Nythus, P.J. (Eds.); Tigers of the world: the science, politics, and conservation of *Panthera tigris*. Elsevier, New York, New York, USA, pp. 143-161.
- Schipper, J., Chanson, J.S., Chiozza, F., Cox, N.A., Hoffmann, M., Katariya, V., Lamoreaux, J., Rodrigues, A.S.L., Cooke, J.G., Duckworth, J.W., Brooks, T.M., Foster, M., Williamson, E.A., Wyatt, S.A., Yan, X. & Young, B.E. 2008: The status of the world's land and marine mammals: diversity, threat and knowledge. - Science 322: 225-230.
- Seidensticker, J. 2010: Saving wild tigers: A case study in biodiversity loss and challenges to be met for recovery beyond 2010. - Integrative Zoology 5 (4): 285-299.
- Simcharoen, S., Pattanavibool, A., Karanth K.U., Nichols,

- J.D. & Sambar Kumar, N. 2007: How many Tigers *Panthera tigris* are there in Huai Kha Khaeng Wildlife Sanctuary, Thailand? An estimate using photographic capture-recapture sampling. - *Oryx* 41 (4): 447-453.
- Smith, J.L.D., Saksit, S., Simchareon, A., Cutter, P., Gurung, B., Chundawat, R., McDougal, C. & Seidensticker, J. 2011: Seasonally dry forest is essential tiger habitat. - In: McShea, W.J., Davies, S.J. & Bhumpakphan, N. (Eds.); The ecology and conservation of seasonally dry forests in Asia. - Smithsonian Institution Scholarly Press, Washington D.C., USA, pp. 1-8.
- Sodhi, N.S., Koh, L.P., Brook, B.W. & Ng, P.K.L. 2004: Southeast Asian biodiversity: an impending disaster. - *Trends in Ecology & Evolution* 19 (12): 654-660.
- Sodhi, N.S., Posa, M.R.C., Lee, T.M., Bickford, D., Koh, L.P. & Brook, B.W. 2009: The state and conservation of Southeast Asian biodiversity. - *Biodiversity & Conservation* 19 (3): 317-328.
- Steinmetz, R., Chutipong, W., Seuaturien, N., Chirngsaard, W. & Khaengkhetkarn, M. 2010: Population recovery patterns of Southeast Asian ungulates after poaching. - *Biological Conservation* 143 (1): 42-51.
- Sutherland, W.J., Pullin, P.M. & Knight, T.M. 2004: The need for evidence-based conservation. - *Trends in Ecology & Evolution* 19 (7): 305-308.
- Thomas, L., Buckland, S.T., Rexstad, E.A., Laake, J.L., Strindberg, S., Hedlet, S.L., Bishop, J.R.B., Marques, T.A. & Burnham, K.P. 2010: Distance software: design and analysis of distance sampling surveys for estimating population size. - *Journal of Applied Ecology* 2010 (1): 5-14.
- Tordoff, A.W., Timmins, R.J., Maxwell, A., Huy, K., Lic, V. & Khou, E.H. 2005: Biological assessment of the Lower Mekong Dry Forests Ecoregion. - WWF Greater Mekong, Phnom Penh, Cambodia, 171 pp.
- Walston, J., Robinson, J.G., Bennett, E.L., Breitenmoser, U., da Fonseca, G.A.B., Goodrich, J., Gumal, M., Hunter, L., Johnson, A., Karanth, K.U., Leader-Williams, N., Mackinnon, K., Miquelle, D., Pattanavibool, A., Poole, C., Rabinowitz, A., Smith, J.L.D., Stokes, E.J., Stuart, S.N., Vongkhamheng, C. & Wibisono, H. 2010: Bringing the Tiger Back from the Brink - The Six Percent Solution. - *PLOS Biology*, 8, e1000485.
- Wang, S.W. 2010: Estimating population densities and biomass of ungulates in the temperate ecosystem of Bhutan. - *Oryx* 44 (3): 376-382.
- Wegge, P. & Storaas, T. 2009: Sampling tiger ungulate prey by the distance method: lessons learned in Bardia National Park, Nepal. - *Animal Conservation* 12 (1): 78-84.
- Wharton, C.H. 1957: An Ecological Study of the Kouprey (*Novibos sauveli* Urbain). - Monographs of the Institute of Science and Technology, Monograph 5, Manila, Philippines, 111 pp.
- Wikramanayake, E., Dinerstein, E., Seidensticker, J., Lumpkin, S., Pandav, B., Shrestha, M., Mishra, H., Ballou, J., Johnsingh, A.J.T., Chestin, I., Sunarto, S., Thinley, P., Thapa, K., Jiang, G., Elagupillay, S., Kafley, H., Pradhan, N.H.B., Jigme, K., Seng, T., Cutter, P., Aziz, M.A. & Than, U. 2011: A landscape-based conservation strategy to double the wild tiger population. - *Conservation Letters* 3 (3): 219-227.
- Williams, B.K., Nichols, J.D. & Conroy, M.J. 2002: Analysis and Management of Animal Populations. 1st edition. - Academic Press, New York, New York, USA, 817 pp.